

Evidence of hydrolyzed traditional Korean red ginseng by malted barley on activation of receptor interacting proteins 2 and I κ B kinase-beta in mouse peritoneal macrophages

Hong-Kun Rim[†], Kyu-Yeob Kim[†], Phil-Dong Moon^{*}

Department of Pharmacology, College of Oriental Medicine, Institute of Oriental Medicine, Oriental Medical Science Center, Kyung Hee University, 1 Hoegi-dong, Dongdaemun-gu, Seoul 130-701, Republic of Korea

ABSTRACT

Red ginseng, which has a variety of biological and pharmacological activities including antioxidant, anti-inflammatory, antimutagenic and anticarcinogenic effects, has been used for thousands of years as a general tonic in traditional oriental medicine. Here, we tested the immune regulatory activities of hydrolyzed red ginseng by malted barley (HRG) on the expressions of receptor interacting proteins (Rip) 2 and I κ B kinase- β (IKK- β) in mouse peritoneal macrophages. We show that HRG increased the activations of Rip 2 and IKK- β for the first time. When HRG was used in combination with recombinant interferon- γ (rIFN- γ), there was a marked cooperative induction of nitric oxide (NO) production. The increased expression of inducible NO synthase from rIFN- γ plus HRG-stimulated cells was almost completely inhibited by pre-treatment with pyrrolidine dithiocarbamate (PDTC), an inhibitor of nuclear factor- κ B (NF- κ B). In addition, the treatment of peritoneal macrophages with rIFN- γ plus HRG caused significant increases in tumor necrosis factor (TNF)- α mRNA expression and production. Because NO and TNF- α play an important role in the immune function and host defense, HRG treatment can modulate several aspects of the host defense mechanisms as a result of the stimulations of the inducible nitric oxide synthase and NF- κ B. In conclusion, our findings demonstrate that HRG increases the productions of NO and TNF- α from rIFN- γ -primed macrophages and suggest that Rip2/IKK- β plays a critical role in mediating these immune regulatory effects of HRG.

Keywords red ginseng, receptor interacting proteins 2, I κ B kinase- β , nuclear factor- κ B, tumor necrosis factor- α , nitric oxide

INTRODUCTION

Ginseng has been used for thousands of years in Asian countries, particularly in Korea, China, and Japan, due to its wide medicinal effects, such as tonic, immunomodulatory, and anti-aging activities (Lee, 2005). In oriental medicine, the production of red ginseng is considered a form of processing medicinal herbs under a certain traditional theory. In addition, related to the theory of how medicine works within the body, it is regarded that the minor cold character is transformed into a warm character, and that the effect of protecting the vitality is improved. Korean red ginseng has been developed for long-term storage and distribution. It has been reported that red ginseng has more powerful pharmacological activities than white ginseng (Nam, 2005; Takagu et al., 1990). The differences in the biological activities of red and white ginseng may result from a change in the chemical constituents that occur during the steaming process (Ha et al., 2005; Park et al., 2005).

Products of partially or totally hydrolyzed materials, including rice or guar gum, by enzymatic hydrolysis with

various enzymes are known to have improved beneficial effects in humans (Stewart and Slavin, 2006). One of the examples of enzymatic hydrolysis is hydrolysis by malted barley in the Korean traditional method. It is well recognized that germinated barley seeds contain a variety of enzymes such as aspartic peptidases, serine peptidases and metallopeptidases, and products hydrolyzed by these enzymes exert many physiological functions (Fontanini and Jones, 2001; Gorjanovic et al., 2002).

Nuclear factor (NF)- κ B plays a critical role in the expression of many genes involved in immune and inflammatory responses (Ghosh et al., 1998; Thanos and Maniatis, 1995). In unstimulated cells, Rel protein dimers, mainly p50 and p65 subunits, are sequestered in the cytoplasm in complex with one of the several inhibitors of NF- κ B. The activation of NF- κ B is the consequence of the phosphorylation of two specific serines near the N terminus of I κ B- α and its degradation. The phosphorylation of I κ B- α leads to ubiquitination, resulting in the degradation, which targets the protein for degradation through the 26S proteasome and the translocation of NF- κ B to the nucleus (Scherer et al., 1995).

The I κ B kinase (IKK)- β enzyme complex is part of the upstream NF- κ B signal transduction cascade (Karin, 1999). After activation, IKK phosphorylates I κ B at two conserved serine residues in the N-terminus, which then triggers the degradation of this inhibitor and allows the rapid translocation of NF κ B into the nuclei where it avidly binds to DNA (Jeong et

*Correspondence: Phil-Dong Moon

E-mail: pdmoon@khu.ac.kr

[†]H.K. Rim and K.Y. Kim contributed equally to this work.

Received May 21, 2012; Accepted August 20, 2012; Published August 31, 2012

doi: <http://dx.doi.org/10.5667/tang.2012.0020>

© 2012 by Association of Humanitas Medicine

TANG / www.e-tang.org

al., 2002). Thus, NF- κ B's translocation and transactivation induce the inducible nitric oxide synthase (iNOS) and tumor necrosis factor (TNF)- α expression in macrophages. Receptor interacting proteins 2 (Rip 2), a serine/threonine kinase, is also essential for NF- κ B activation through the TNF receptor and Toll-like receptors (Kobayashi et al., 2002; Inohara et al., 1998).

In the present study, we show that hydrolyzed red ginseng by malted barley (HRG) synergistically induces Nitric oxide (NO) and TNF- α production by peritoneal macrophages when the cells are treated by recombinant IFN- γ (rIFN- γ). To investigate the mechanisms of HRG-induced NO and TNF- α production, we examined the effects of HRG on Rip2, IKK- β , and NF- κ B activation. These findings may explain the influence of LS on NO and TNF- α production via the NF- κ B signaling pathway.

MATERIALS AND METHODS

Reagents

Korean red ginseng was obtained from the Korea Ginseng Corp. (Daejeon, Republic of Korea). Murine rIFN- γ , recombinant TNF- α (rTNF- α), biotinylated TNF- α , and anti-murine TNF- α was purchased from R&D Systems (Minneapolis, Minnesota, USA). N-(1-naphthyl)- ethylenediamine dihydrochloride, lipopolysaccharide (LPS), sodium nitrite, and PDTC were purchased from Sigma (St. Louis, Missouri, USA). Rabbit polyclonal antisera to iNOS were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, California, USA). N^GMMA was purchased from Calbiochem (San Diego, California, USA). Thioglycollate (TG) was purchased from Difco Laboratories (Detroit, Michigan, USA). Dulbecco's Modified Eagle's Medium (DMEM) containing L-arginine (84 mg/l), Hank's balanced salt solution (HBSS), fetal bovine serum (FBS), and other tissue culture reagents were purchased from Gibco BRL (Grand Island, New York, USA).

Animals

The original stock of male C57BL/6J mice were purchased from the Daehan Biolink Co., (Daejeon, Republic of Korea), and were housed at a temperature of $23 \pm 1^\circ\text{C}$ with a 12/12 h light dark cycle. Food and water were available *ad libitum*. All protocols were approved by the institutional animal care and use committee of Kyung Hee University.

Preparation of HRG

The 90 g of raw rice and 10 g of red ginseng were washed and air-dried. These were precipitated in 200 ml distilled water for 2 h, and then boiled at 121°C for 15 min. After 1 kg of malted barley was extracted with 1 l distilled water at 60°C for 2 h, the resulted extracts were added and hydrolyzed the mixture of raw rice and red ginseng. The mixture was reacted at 60°C for 3 h, and then boiled at 90°C for 5 min. The reaction mixture was filtered and kept at 4°C .

Assay for endotoxin determination

The HRG used in this experiment was found to be free of endotoxin as determined within the limits of an assay E-TOXATE kit (Sigma, USA), preformed according to the manufacturers protocol. In this assay, saturation occurred at 40 EU/ml and the resolution limit was > 0.1 EU/ml.

Peritoneal macrophage cultures

TG-elicited macrophages were harvested 3 - 4 days after i.p. injection of 2.5 ml of TG to the mice and isolated, as reported

previously (An et al., 2008). Using 8 ml of HBSS containing 10 U/ml heparin, peritoneal lavage was performed. Then, the cells were distributed in DMEM, which was supplemented with 10% heat-inactivated FBS, in culture plates incubated for 3 h at 37°C in an atmosphere of 5% CO_2 , washed three times with HBSS to remove non-adherent cells, and equilibrated with DMEM that contained 10% FBS before treatment.

MTT assay

Peritoneal macrophages (2.5×10^5 cells/well) were cultured with with rIFN- γ (10 U/ml) for 6 h. The cells were then stimulated with various concentrations of HRG (0.01 - 1 mg/ml). Cell aliquots were incubated with 20 μl of a MTT solution (5 mg/ml) for 4 h at 37°C under 5% CO_2 and 95% air. Consecutively, 250 μl of DMSO was added to extract the MTT formazan and the absorbance of each well at 540 nm was read by an automatic microplate reader (Molecular Devices Corp., Sunwayle, California, USA).

Measurement of nitrite concentration

Peritoneal macrophages (2.5×10^5 cells/well) were cultured with rIFN- γ (10 U/ml) for 6 h. The cells were then stimulated with various concentrations of HRG. After 48 h of culture, NO synthesis in cell cultures was measured by a microplate assay method, as previously described (An et al., 2008). To measure nitrite, 100 μl aliquots were removed from conditioned medium and incubated with an equal volume of Griess reagent (1% sulfanilamide/ 0.1% N-(1-naphthyl)-ethylenediamine dihydrochloride/ 2.5% H_3PO_4) at room temperature for 10 min. The absorbance at 540 nm was determined by an automatic microplate reader (Molecular Devices Corp., Sunwayle, California, USA). NO_2^- was determined by using sodium nitrite as a standard. The cell-free medium alone contained 5 to 8 μM of NO_2^- . This value was determined in each experiment and subtracted from the value obtained from the medium with peritoneal macrophages.

Assay of TNF- α release

Peritoneal macrophages (2.5×10^5 cells/well) were incubated with rIFN- γ (10 U/ml), HRG, rIFN- γ plus LPS (10 $\mu\text{g}/\text{ml}$), and rIFN- γ plus various concentrations of HRG (0.01 - 1 mg/ml) for 24 h. Then the amount of TNF- α secreted by the cells was measured by a modified enzyme-linked immunosorbent assay (ELISA), as described previously (An et al., 2008). The ELISA was devised by coating 96-well plates of murine monoclonal antibody with specificity for TNF- α . Before use and between subsequent steps in the assay, coated plates were washed with phosphate-buffered saline (PBS) containing 0.05% Tween-20. All reagents used in this assay were incubated for 2 h at 37°C . The rTNF- α was diluted and used as a standard. Serial dilutions starting from 10 ng/ml were used to establish the standard curve. After 2 h incubation at 37°C , the wells were washed and then each of 0.2 $\mu\text{g}/\text{ml}$ of biotinylated anti-mouse TNF- α was added and the plates were incubated at 37°C for 2 h. After washing the wells, avidin-peroxidase was added and the plates were incubated for 20 min at 37°C . Wells were again washed and ABTS was added. Color development was measured at 405 nm using an automated microplate ELISA reader (Molecular Devices Corp., Sunwayle, California, USA).

Nuclear protein extraction

Nuclear extracts were prepared as described previously (Jeong et al., 2002). Briefly, after cell activation for the times indicated cells were washed with ice-cold PBS and resuspended in 60 μl of buffer A (10 mM HEPES/KOH, 2 mM MgCl_2 , 0.1 mM EDTA, 10 mM KCl, 1 mM DTT, and 0.5 mM PMSF, pH 7.9).

The cells were allowed to swell on ice for 15 min, lysed gently with 2.5 μ l of 10% Nonidet P-40, and centrifuged at 2000 g for 10 min at 4°C. The nuclei pellet was resuspended in 40 μ l of buffer B (50 mM HEPES/KOH, 50 mM KCl, 300 mM NaCl, 0.1 mM EDTA, 10% glycerol, 1 mM DTT, and 0.5 mM PMSF, pH 7.9), left on ice for 20 min, and inverted. The nuclear debris was then spun down at 15,000 g for 15 min. The supernatant (nuclear extract) was collected, frozen in liquid nitrogen and stored at -70°C until conducting the analysis.

Western blot analysis

Peritoneal macrophages (5×10^6 cells/well) were incubated for 6 h with rIFN- γ (R&D Systems, Minneapolis, MN, USA; 10 U/ml). The cells were then stimulated with HRG (1 mg/ml) or LPS (10 μ g/ml). Whole cell lysates were made by boiling peritoneal macrophages in sample buffer [62.5 mM Tris-HCl, pH 6.8, 2% sodium dodecyl sulphate (SDS), 20% glycerol, and 10% 2-mercaptoethanol]. Proteins in the cell lysates were then separated by 10% SDS-polyacrylamide gel electrophoresis and transferred to nitrocellulose paper. The membrane was then blocked with 5% skim milk in PBS-tween-20 and then incubated with antibodies (Santa Cruz Biotechnology, Inc., CA, USA). After washing in PBS-tween-20 three times, the blot was incubated with 1:3000 diluted anti-rabbit or anti-mouse IgG, horseradsh peroxidase linked whole antibody (Amersham Corp. Newark, NJ, USA), and the antibody-specific proteins were visualized using the enhanced chemiluminescence detection system according to the recommended procedure (Amersham Corp. Newark, NJ, USA).

Reverse transcription-polymerase chain reaction (RT-PCR) analysis

Total RNA was isolated from macrophages according to the manufacturer's specification using an easy-BLUE RNA extraction kit (iNtRON Biotech, Korea). Total RNA (2.0 μ g) was heated at 75°C for 5 min and then chilled on ice. Each sample was reverse-transcribed to cDNA for 60 min at 42°C using a cDNA synthesis kit (iNtRON Biotech). PCR was performed with the following primers for TNF- α (5'-ATGAGAACAGAAAGCATGATC-3'; 5'-TACAGGCTTG TCACTCGAATT-3'), iNOS (5'-TCACTGGGACAGCAC AGAAT-3'; 5'-TGTGTCTGCAGATGTGCTGA-3'), and glyceraldehydes-3-phosphate dehydrogenase (GAPDH) (5'-GGCATGGACTGTGGTCATGA-3'; 5'-TTCACCACCAT GGAGAAGGC-3') which were used to verify if equal amounts of RNA were used for reverse transcription and PCR amplification from different experimental conditions. The annealing temperature was 58°C for iNOS, 60°C for TNF- α , and 62°C for GAPDH, respectively. Products were electrophoresed on a 1.5% agarose gel and visualized by staining with ethidium bromide. GAPDH mRNA was used as a control of mRNA loading.

Statistical analysis

Results were expressed as the mean \pm S.E.M. of independent experiments, and statistical analyses were performed by one-way analysis of variance (ANOVA) with Tukey, and Duncan post hoc test to express the difference between groups. All statistical analyses were performed using SPSS v11.0 statistical analysis software (SPSS INC., Chicago, Illinois, USA). A value of $p < 0.05$ was considered to indicate statistical significance.

RESULTS

Effect of HRG on activation of Rip2, IKK- β , and NF- κ B

Macrophages play an important role in immune-enhancement. When the macrophage was activated, the expressions of iNOS and TNF- α increased. Rip2, IKK- β , and NF- κ B are upstream proteins for the expressions of iNOS and TNF- α . To investigate the immunostimulating activities of HRG, we investigated the effects of HRG on the activations of IKK- β , Rip2, and NF- κ B. As shown in Fig. 1A, stimulation of peritoneal macrophages with LPS after the treatment of rIFN- γ resulted in the increased protein synthesis of Rip2 and IKK- β . When HRG was used in combination with rIFN- γ , there was a marked co-operative activation of Rip2 and IKK- β . Since NF- κ B activation requires the nuclear translocation of the *Rel A/p65* subunit of NF- κ B, we examined the effect of HRG on the nuclear pool of *Rel A/p65* protein by Western blot analysis. As shown in Fig. 1B, rIFN- γ plus HRG treatment considerably increased the nuclear *Rel A/p65* protein level, indicating the nuclear translocation of *Rel A/p65*. However, HRG had no effect on the activations of Rip2, IKK- β , and NF- κ B by itself.

Effects of HRG on NO production in peritoneal macrophages

The downstream mediators of the pathways involving Rip2, IKK- β , and NF- κ B were investigated. NO is a highly reactive molecule produced from the guanidino nitrogen of arginine by iNOS (Nathan, 1992). Over the past decade, NO as a potent




Fig. 1. Effect of HRG on the activations of Rip 2, IKK- β , and NF- κ B in peritoneal macrophages. (A) The cells (5×10^6 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml). The cells were then stimulated with HRG (1 mg/ml) or LPS (10 μ g/ml). The protein extracts were prepared, and then the samples were analyzed for Rip2 and IKK- β expressions by Western blot analysis. (B) The cells (1×10^7 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml). The cells were then stimulated with HRG (1 mg/ml) or LPS (10 μ g/ml). The nuclear protein extracts were prepared, and then the samples were analyzed for NF- κ B expression by Western blot analysis. PARP, Poly ADP-ribose polymerase.




Fig. 2. Effect of HRG on NO synthesis in peritoneal macrophages. The cells (2.5×10^5 cells/well) were cultured with rIFN- γ (10 U/ml). (A) The cells were then stimulated with various concentrations of HRG for 6 h after incubation. After 48 h of culturing, NO release was measured. (B) The cells (2.5×10^5 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml) plus various concentrations of N^GMMA. The cells were then treated with HRG (1 mg/ml) and cultured for 48 h. NO (nitrite) released into the medium is presented as the mean \pm S.E.M. of three independent experiments duplicated in each run. Control, rIFN- γ + HRG (1 mg/ml). * $p < 0.05$ compared to rIFN- γ alone; # $p < 0.05$ compared to control (absence of N^GMMA).

macrophage-derived effector molecule against a variety of bacteria, parasites, and tumors has received increasing attention (Gantt et al., 2001). To determine the effect of HRG on the production of NO by mouse peritoneal macrophages, we treated non-primed (resting) and rIFN- γ -primed cells with HRG. The resultant NO production was determined by detecting the levels of nitrite concentration in the cell supernatants after 48 h treatment. As shown in Fig. 2A, HRG had no effect on NO production in resting mouse peritoneal macrophages. IFN- γ (10 U/ml) alone does not cause the induction of NO production in accordance with previous reports. However, when IFN- γ was used in conjunction with LPS, there was a marked augmentation of the production of NO. When mouse peritoneal macrophages were primed for 6 h with murine rIFN- γ and then treated with HRG, NO production increased compared to non-primed conditions. HRG didn't affect the viability of cells




Fig. 3. Effects of HRG on the expressions of iNOS mRNA and protein in peritoneal macrophages. (A) The cells (5×10^6 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml). The cells were then stimulated with HRG (1 mg/ml) or LPS (10 μ g/ml) for 24 h. The RNA extracts were prepared, and then the samples were analyzed for iNOS mRNA expression by RT-PCR. (B) The cells (5×10^6 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml) and then treated PDTC (100 μ M) for 1 h. The cells were then stimulated with HRG (1 mg/ml) or LPS (10 μ g/ml) for 24 h. The protein extracts were prepared and then the samples were analyzed for iNOS protein expression by Western blot analysis.

at those concentrations (data not shown). To determine if the signaling mechanism in HRG-induced NO production participates in the L-arginine-dependent pathway in mouse peritoneal macrophages, the cells were incubated for 6 h in the presence of rIFN- γ plus N^GMMA. The production of nitrites by rIFN- γ plus HRG in mouse peritoneal macrophages was progressively inhibited with the increasing amounts of N^GMMA. The HRG-induced accumulation of nitrites was significantly blocked by N^GMMA (0.01 - 1 μ M) (Fig. 2B, $p < 0.05$).

Effects of HRG on rIFN- γ -primed iNOS expression

Fig. 3 shows the effects of rIFN- γ plus HRG treatments on the expressions of iNOS mRNA and protein mouse peritoneal macrophages. HRG (1 mg/ml) increased synergistically the expressions of iNOS mRNA and protein in rIFN- γ -primed mouse peritoneal macrophages. It is known that PDTC, an anti-oxidant compound, inhibits NF- κ B activation (Chung et al., 2004). With this understanding, as an approach to define the signaling mechanism of HRG on NO production, we examined the influence of PDTC in rIFN- γ plus HRG-treated mouse peritoneal macrophages. Adding PDTC (100 μ M) to the rIFN- γ plus HRG-treated mouse peritoneal macrophages decreased the synergistic effects of HRG on iNOS expression (Fig. 3B).

Effects of HRG on rIFN- γ -primed TNF- α mRNA expression and production

We next examined the synergistic cooperative effects of HRG on rIFN- γ -induced TNF- α production. Mouse peritoneal macrophages secreted very low levels of TNF- α after 24 h incubation with a medium alone or rIFN- γ alone. However, HRG in combination with rIFN- γ markedly increased TNF- α mRNA expression and production (Fig. 4).




Fig. 4. Effects of HRG on TNF- α mRNA expression and production in peritoneal macrophages. (A) The cells (5×10^6 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml). The cells were then stimulated with HRG (1 mg/ml) or LPS (10 μ g/ml). The RNA extracts were prepared, and then the samples were analyzed for TNF- α expression by RT-PCR. (B) The cells (2.5×10^5 cells/well) were incubated for 6 h with rIFN- γ (10 U/ml). The cells were then stimulated with HRG (0.01 - 1 mg/ml) or LPS (10 μ g/ml). The amount of TNF- α secreted by peritoneal macrophages was measured by the ELISA method after 24 h of incubation. Values are the mean \pm S.E.M. of three independent experiments duplicated in each run. * $p < 0.05$ compared to rIFN- γ alone.

DISCUSSION

In this study, we showed that HRG induced the expressions of iNOS and TNF- α through the activation of NF- κ B, IKK- β , and Rip 2.

Red ginseng is produced by steaming fresh ginseng first, and then drying the steamed ginseng; on the other hand, white ginseng is made of fresh ginseng by the drying process only. Recently, several investigators have reported the discovery of new ginsenosides from red ginseng which are not usually found in raw ginseng. These ginsenosides are Rg3, Rg5, Rg6, Rh2, Rh4, Rs3, and Rf and their levels in red ginseng are relatively low (Kim et al., 2000; Park, 1996). Ginsenosides Rg3, Rg5, and Rh2 have shown anticancer effects (Yun, 2003). Among these three ginsenosides, Rg3 has been reported to exhibit neuroprotective (Tian et al., 2005), antinociceptive (Rhim et al., 2002), hematopoietic (Joo et al., 2004), anticarcinogenic, and antimetastatic effects (Tao et al., 2002; Yun et al., 2001). Choi et al., reported that red ginseng acidic polysaccharide in

combination with IFN- γ enhanced the macrophage function through activation of the NF- κ B pathway (2008). In this study, we demonstrated that NO production in mouse peritoneal macrophages by HRG could be highly stimulated in combination with rIFN- γ . HRG increased NO production to a higher level than red ginseng or ginsenosides. Therefore, we postulate that HRG may contain the many active compounds. However, further studies to analyze the active compounds of HRG are necessary to support the present findings.

It has been previously reported that LPS stimulation of rIFN- γ -primed macrophages induces NF- κ B activation (Shin et al., 2004). Expressions of iNOS and the TNF- α gene are dependent on the activation of NF- κ B (Baldwin, 1996). We found that the addition of the NF- κ B modulator, PDTC, inhibits the synergistic effect of HRG with rIFN- γ on iNOS expression. These results suggest that HRG increases NO production through NF- κ B activation. The NF- κ B system may provide a future target in cancer therapy. Rip2 and IKK complexes may play an important role for NF- κ B activation (Inohara et al., 2000). Until now, the regulatory effect of drugs in Rip 2 activation has not been reported. In this study, we showed that HRG induced the activations of Rip 2 and IKK- β for the first time. The results of this study suggest that HRG activates NF- κ B via interaction with Rip 2 and IKK- β .

In conclusion, our results demonstrated that HRG acted as an accelerator of peritoneal macrophages activation by rIFN- γ via a process involving L-arginine-dependent NO production and that it increased the production of TNF- α significantly via NF- κ B activation by interaction with Rip 2 and IKK- β . These results suggest that the processes of HRG might explain its regulatory effect in the immune system.

ACKNOWLEDGEMENTS

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2005-0049404).

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

REFERENCES

- An HJ, Seo MJ, Choi IY, Park RK, Jeong S, Lee JY, Kim HM, Um JY, Hong SH. Induction of nitric oxide & tumour necrosis factor- α by *Psoralea corylifolia*. *Indian J Med Res.* 2008;128:752-758.
- Baldwin AS Jr. The NF-kappa B and I kappa B proteins: new discoveries and insights. *Annu Rev Immunol.* 1996;14:649-683.
- Choi HS, Kim KH, Sohn E, Park JD, Kim BO, Moon EY, Rhee DK, Pyo S. Red ginseng acidic polysaccharide (RGAP) in combination with IFN-gamma results in enhanced macrophage function through activation of the NF-kappaB pathway. *Biosci Biotechnol Biochem.* 2008;72:1817-1825.
- Chung HS, An HJ, Jeong HJ, Won JH, Hong SH, Kim HM. Water extract isolated from *Chelidonium majus* enhances nitric oxide and tumour necrosis factor- α production via nuclear

- factor-kappaB activation in mouse peritoneal macrophages. *J Pharm Pharmacol.* 2004;56:129-134.
- Fontanini D, Jones BL. Study of metallopeptidase isozymes from malted barley (*Hordeum vulgare* cv. Morex). *J Agr Food Chem.* 2001;49:4903-4911.
- Gantt KR, Goldman TL, McCormick ML, Miller MA, Jeronimo SM, Nascimento ET, Britigan BE, Wilson ME. Oxidative responses of human and murine macrophages during phagocytosis of *Leishmania chagasi*. *J Immunol.* 2001;167:893-901.
- Ghosh S, May MJ, Kopp EB. NF- κ B and Rel proteins: evolutionarily conserved mediators of immune responses. *Ann Rev Immunol.* 1998;16:225-260.
- Gorjanovic S, Cvetkovic A, Suznjevic D, Beljanski M, Vrvic M, Hranisavljevic J. Isolation and characterization of highly liganded protein from brewer's barley grain. *Biosci Biotech Bioch.* 2002;66:1940-1944.
- Ha DC, Lee JW, Ryu GH. Change in ginsenosides and maltol in dried raw ginseng during extrusion process. *Food Sci Biotechnol.* 2005;14:363-367.
- Inohara N, del Peso L, Koseki T, Chen S, Nunez G. RICK, a novel protein kinase containing a caspase recruitment domain, interacts with CLARP and regulates CD95-mediated apoptosis. *J Biol Chem.* 1998;273:12296-12300.
- Inohara N, Koseki T, Lin J, del Peso L, Lucas PC, Chen FF, Ogura Y, Núñez G. An induced proximity model for NF-kappa B activation in the Nod1/RICK and RIP signaling pathways. *J Biol Chem.* 2000;275:27823-27831.
- Jeong HJ, Hong SH, Lee DJ, Park JH, Kim KS, Kim HM. Role of Ca(2+) on TNF-alpha and IL-6 secretion from RBL-2H3 mast cells. *Cell Signal.* 2002;14:633-639.
- Joo SS, Won TJ, Kim MS, Lee DI. Hematopoietic effect of ginsenoside Rg3 in ICR mouse primary cultures and its application to a biological response modifier. *Fitoterapia.* 2004;75:337-341.
- Karin M. The beginning of the end: IkappaB kinase (IKK) and NF-kappaB activation. *J Biol Chem.* 1999;274:27339-27342.
- Kim WY, Kim JM, Han SB, Lee SK, Kim ND, Park MK, Kim CK, Park JH. Steaming of ginseng at high temperature enhances biological activity. *J Nat Prod.* 2000;63:1702-1704.
- Kobayashi K, Inohara N, Hernandez LD, Galan JE, Nunez G, Janeway CA, Medzhitov R, Flavell RA. RICK/Rip2/CARDIAK mediates signalling for receptors of the innate and adaptive immune systems. *Nature.* 2002;416:194-199.
- Lee TK. Radioprotective potential of ginseng. *Mutagenesis.* 2005;20:237-239.
- Nam KY. The comparative understanding between red ginseng and white ginseng, processed ginseng (*Panax ginseng* C.A. Meyer). *J Ginseng Res.* 2005;29:1-18.
- Nathan C. Nitric oxide as a secretory product of mammalian cells. *Faseb J.* 1992;6:3051-3064.
- Park JD. Recent studies on the chemical constituents of Korean ginseng (*Panax ginseng* C.A. Meyer). *Korean J Ginseng Sci.* 1996;20:389-415.
- Park JH, Cha HY, Seo JJ, Hong JT, Han K, Oh KW. Anxiolytic-like effects of ginseng in the elevated plus-maze model: comparison of red ginseng and sun ginseng. *Prog Neuro-Psychoph.* 2005;29:895-900.
- Rhim H, Kim H, Lee DY, Oh TH, Nah SY. Ginseng and ginsenoside Rg3, a newly identified active ingredient of ginseng, modulate Ca²⁺ channel currents in rat sensory neurons. *Eur J Pharmacol.* 2002;436:151-158.
- Scherer DC, Brockman JA, Chen Z, Maniatis T, Ballard DW. Signal-induced degradation of I kappa B alpha requires site-specific ubiquitination. *Proc Natl Acad Sci USA.* 1995;92:11259-11263.
- Shin HM, Kim MH, Kim BH, Jung SH, Kim YS, Park HJ, Hong JT, Min KR, Kim Y. Inhibitory action of novel aromatic diamine compound on lipopolysaccharide-induced nuclear translocation of NF-kappaB without affecting IkappaB degradation. *FEBS Lett.* 2004;571:50-54.
- Stewart ML, Slavin JL. Molecular weight of guar gum affects short-chain fatty acid profile in model intestinal fermentation. *Mol Nutr Food Res.* 2006;50:971-976.
- Takagu T, Kameda K, Matsuura Y, Sekiya K, Okuda H. Studies on insulin-like substances in Korean red ginseng. *Planta Med.* 1990;56:27-30.
- Tao H, Yao M, Zou S, Zhao D, Qiu H. Effect of angiogenesis inhibitor Rg3 on the growth and metastasis of gastric cancer in SCID mice. *Zhounghua Wai Ke Za Zhi.* 2002;40:606-608.
- Thanos D, Maniatis T. NF-kB: a lesson in family values. *Cell.* 1995;80:529-532.
- Tian JW, Fu FH, Geng MY, Jiang YT, Yang JX, Jiang WL, Wang CY, Liu K. Neuroprotective effect of 20(S)-ginsenoside Rg3 on cerebral ischemia in rats. *Neurosci Lett.* 2005;374:92-97.
- Yun TK. Experimental and epidemiological evidence on non-organ specific cancer preventive effect of Korean ginseng and identification of active compounds. *Mutat Res.* 2003;523:63-74.
- Yun TK, Lee YS, Lee YH, Kim SI, Yun HY. Anticarcinogenic effect of *Panax ginseng* C.A. Meyer and identification of active compounds. *J Korean Med Sci.* 2001;16:6-18.