

〈 Review 〉

## Fire and Rangeland's Management in India

Paulsamy, S., S. Manian<sup>1</sup> and Bong-Seop Kil<sup>2\*</sup>

*Department of Botany, Kongunadu Arts and Science College, Coimbatore - 641 029, India*

<sup>1</sup> *Department of Botany, Bharathiar University, Coimbatore - 641 046, India*

<sup>2</sup> *Division of Life Science, Wonkwang University, Iksan 570-749, Korea*

**ABSTRACT:** Fire, usually a detrimental factor makes changes in commonly structure and functions when it is practiced properly. In countries like Australia, South Africa, Japan and Texas in USA etc., fire is used as a tool to manage the natural ecosystems in productive condition on the basis of ecological approach. However, in developing countries like India there is effective measure to use and control fire in rangelands. This review provides knowledge on changes in sociological characters of plants, biomass profile, cycling of materials and strategy to use fire for the shaping of communities in different rangelands located in India and other countries. Further, the informations furnished in this article will be more useful for the forest managers, planners of rangelands and administrators in countries like India and tropical countries with similar bioclimatic conditions to prepare appropriate management plans where the fire is an integral factor.

**Key words:** Biomass, Fire management, Forest fire, Nutrient cycling, Phytosociology

### INTRODUCTION

Fire, always a powerful factor, destroys / designs community structure in majority of rangelands. The effects of fire on communities depend on season of burn, frequency and intensity of fire, velocity of wind, moisture content of air, type of plant species in the community, whether annuals or perennials and the macro-climatic zone where the fire has happened. Besides, the changes in community composition made by individual species, all functional attributes like flow of energy, availability and rate of transfer of nutrients between the tropic levels, the sequence of food chain, complexity of food web, biodiversity status, rate and direction of community succession and finally the degree of homeostasis are vigorously influenced by fire in ecosystems (Daubenmire 1968). Management of communities in fire pruned rangelands is habitat specific and depends on many external and internal factors (Show and Clarke 1994). Many workers have been interested on fire concerning grazing land, nitrogen fixers, minerals, nutrient uptake, etc. This review deals with the aspects like fire regime, fire effects on phytosociology, biomass structure and nutrient cycling in the rangelands, role of fire as a management tool in shaping vegetations and ecology of fire influenced rangelands in and over India.

### FIRE REGIME

Fire in a rangeland may be either wild or prescribed. Wild fire

is often not desired due to difficulties in post-burn management. On the other hand, prescribed burning has a role to play for many rangeland managers. A number of steps viz., preburn planning, the burning operation and postburn management are required to conduct any prescribed burning (Bailey 1988). When developing a burning programme for a game ranch, the most important factors to consider are the reasons for burning and the appropriate fire regime to be applied (Trollope 1990). The fire regime refers to the type and intensity of fire, and the season and frequency of burning. The dry fuel load accumulated during summer in the grassland patches of forests is one of the origin places of fire and from there it can enter easily into the forests of deciduous and semi-evergreen types (Paulsamy 1992). Joshi (2003) also pointed out that forest fires of short-to medium-return intervals are quite common during summer seasons in Garhwal Himalaya.

Effect of fire on grasses is largely determined by season of burn, size of plant, amount of dead material, growth form, species, precipitation and whether it is an annual or perennial (Wright 1985). Fire season is likely to be the dominant feature of any prescribed fire regime and is directly amenable to management. Griffin and Friedel (1984, 1985) found that winter fire was effective in reducing fuel and was easily controlled. It maintained or improved the composition of pasture plants palatable to cattle. A summer fire significantly decreased the grass component while increasing the proportion of forbs, particularly unpalatable species. Rainfall, season of burning and reduction of cover appeared to be important factors in controlling the composition of post-fire herbage. There was no im-

---

\* Corresponding author; e-mail: bskil@wonkwang.ac.kr

portant effect of fire on nutrients in either season of burning. Pressland *et al.* (1986) favoured burning of native vegetation following good rainfall seasons for effectively decreasing the density of some useless woody weeds growing in the mulga (*Acacia aneura*) rangelands of Queensland. This judicious timing of fire also resulted in minimal damage to pasture species as well as post-fire loss of carrying capacity. Long term community responses to altered fire regimes should be predictable from a basic understanding of individual species response to fire (Hodgkinson 1986).

### PHYTOSOCIOLOGY

Most rangeland communities are resistant to fire but are not necessarily stable. Significant changes in structure and composition may occur. Mall and Mehta (1978) noted higher density value in the burned plot when compared to control plot. Further, burning resulted in enhanced tillering and increased percentage germination of the seeds. The response of plants to fire is species specific. Interestingly it has been noted that the dicot weeds in general and the Asteraceae member, *Anaphalis subdecurrens* in particular significantly enhanced its quantitative ecological characters in the post fire communities of subtropical grasslands of Anaimalais (Paulsamy *et al.* 1995, 1996, 1997a, b, c). Similarly, in the same grasslands, the fern, *Pteris aquilina* responded parallelly with the dominant grass, *Chrysopogon zeylanicus* to annual summer fire by improving its ecological characters and it is found to have some clues for the preparation of management plans (Paulsamy *et al.* 1995 a, 1995 b). On the other hand, Paulsamy *et al.* (2001) reported that the species like *Impatiens tomentosa*, *Drosera peltata*, *Osbeckia parviflora*, *Emelia sonchifolia*, *Lecanthus peduncularis* and *Lobelia nicotianifolia* are fire threatened medicinal herbs in the grasslands of Anaimalais and they need immediate attention for conservation.

Fire also affects the seed pool in soil and it varies according to species. Fire can enhance the number and diversity of forbs because they have hard seed coat that can be scarified by fire (Wright 1985). Shaw (1957) and Tothill (1969) cited a number of factors favouring the dominance of black spear grass in Queensland. The sharply pointed seeds become buried in the soil and are protected from fire. Fire reduces the basal cover of competing species and stimulates seed germination whereas the established black speargrass plants are resistant to fire. Hassan and West (1971) reported a significantly lower seed pools in some taxa such as *Bromus tectorum*, *Agropyron* spp. and *Poa* spp. on the burned plots six weeks after fire.

The sprouting response of the rhizomatous grasses generally increases immediately after fire (Wright 1985, Pandey 1988). In the early stages of post-fire growth and under certain conditions, individual plants of *Triodia wiseana* achieved a larger size than *T.*

*basedowii* due to its sprouting response (Casson and Fox 1987). Paulsamy (1992) observed that the sprouting of perennial grasses in general and the dominant grass of the community, *Chrysopogon zeylanicus* in particular is significant over the unburned grasslands in Anaimalais, India.

Fire is a natural phenomenon in the evolution of plants in the pastoral areas. Qualitative evolution of the grasses was more homogeneous in burned plots than in unburnt savanna of Ivory Coast (Abbadie 1985). Populations of most of the Queensland's native plants are capable of surviving in certain fire regimes and may benefit from burning (Anderson and Pressland 1987). In the semi-arid regions of western United States, burning reduced *Bromus* density, allowing forbs and *Sitanion* to successfully compete for growing season precipitation. Whisenant and Bulsiewicz (1985) observed that spring burning of mixed-grass prairie in the Badlands National Park at Western South Dakota, USA significantly reduced Japanese brome density for the first growing season, but returned to its original density by next year. Paulsamy *et al.* (2003b) reported that the grasses and legumes in the high level grasslands of Anaimalais, Western Ghats, Southern India improved their ecological importance significantly by annual summer fire. However, the 'other forbs' in the post fire communities have lost their importances. Fire is often used in combination with other range improvement techniques in the management of grasslands. In the Juniper communities of Texas, prescribed fire is generally used in combination with other range improvement techniques which reduced the Juniper canopy cover, allowing herbaceous vegetation to increase. Noble *et al.* (1986) reiterated that the changes in community structure are not due to fire alone because the interaction of grazing pressure and rainfall regime before and after fire determine the overall ecological impact. Andrew (1986) found that the pasture degradation was arrested if burning was used with grazing in the monsoon tall grass pasture. The species, *Calluna vulgaris* responded well to grazing and other types of management practices including fire (Gimingham 1989). Due to the discontinuity of such management practices, there was a replacement of *Calluna* by other species in the grasslands of Texas, USA.

### BIOMASS

The biomass of grasses, forbs and litter and the relative abundance of different life-forms were highly coupled with the burning cycle. Abrams *et al.* (1986) found that live biomass was greater on burned than unburned low land sites. Biomass of the graminoids was 40 percent lower whereas that of forbs and woody plants was 200-300 per cent greater in the unburned than in the annually burned site of Kansas tall grass prairie. Studies of Andariese and

Covington (1986) for more than two years on the understorey response following both wild fire and prescribed burning in Arizona ponderosa pine showed a general trend towards increased production. Harniss and Murray (1973) also reported a higher production from rhizomatous grasses on burned plots than that of controls for about 30 years. Pandey (1988) observed growth stimulation of both shoot and root components in a *Dichanthium annulatum* dominated grassland after burning. However, the belowground biomass showed a considerable decrease. Research shows that, when correctly practiced, burning can maintain the vegetation in a productive condition with minimal habitat deterioration (Gimingham 1985). Britton *et al.* (1987) stressed the importance of adequate soil moisture at the time of burning to achieve higher production. In south-western New South Wales, Australia, a three year sequence of above-average rainfall immediately following burning in either spring, autumn or winter resulted in a substantial increase in species richness and herbage dry matter production (Noble 1989). In the grasslands of Varanasi, India, the summer burning prior to rains resulted in higher production and thereby the greater recovery rates of biomass compared to winter burning (Pandey 1988). The burned area of Arapaho Prairie, Arthur County, Nebraska initially had more aerial phytomass than the control. However, in later stage, the burned area had statistically lower phytomass production (23% less) than the control (Morrison *et al.* 1986).

Garza and Blackburn (1985) discussed the effect of burning season on the production. In the Post Oak Savanna of Texas, burning in early winter favoured the growth of forbs, whereas spring burning tended to favour the production of grasses. Herbicide treatment and burn-herbicide combination were recognized as suitable management practice for an increased buffel grass (*Cenchrus ciliaris*) standing crop (Mayeux and Hamilton 1983).

### NUTRIENT CYCLING

The importance of various nutrients in determining pasture production or quality varies from soil to soil. Deficiencies of nitrogen or phosphate are probably most widespread, though other macronutrients such as magnesium, potassium and sulphur can be limiting in many soils (Snaydon 1987). One way of making good these deficiencies is through prescribed burning. Increase of mineral salts with burning has already been well documented (Daubenmire 1968, Viro 1974, Fox 1980). However, Paulsamy *et al.* (1993) reported that the minerals like phosphorous and potassium did not get change after fire in the grassland soils of Anaimalais, Western Ghats, India. Another effect of burning is that the nutrients are changed to simple, water soluble salts that are immediately available to plants. Leaching out of the easily soluble phosphate before the restoration of

equilibrium between different electrolytes in the soil is also possible (Viro 1974).

Hobbs and Schimel (1984) concluded that fire in the sagebrush-grassland and mountain brush communities increased the rate of nitrogen mineralization and presumably the availability of nitrogen to the vegetation. Manian *et al.* (2002) reported that the free N<sub>2</sub> fixer, *Azospirillum* increased its population considerably in post-fire grassland communities. They also noted the increase of nodulation in the legume, *Atylosia trinervia* after fire in the Kundah plateau of Nilgiris, southern India which consequently increased the N<sub>2</sub> status in post-fire community. Wells *et al.* (1979) reported that nitrogen fixation, both symbiotic and non-symbiotic was more prevalent following fire, with general increases in P, K, Ca and Mg. The increased root exudates of burned grasslands create a favourable environment for the growth of N<sub>2</sub> fixing and P-solubilizing microorganisms (Senthilkumar *et al.* 1995). Curl and Truelove (1986) pointed out that under the conditions of high temperature during fire with consequent low soil moisture followed by wetting, increased the root exudation which in turn enhanced the population of microbes like N<sub>2</sub> fixers and P-solubilizers. Jalaludin (1969) generalized that the decomposers like *Trichoderma* and *Penicillium* are the pioneers in tropic soil after fire. Lynch and Panting (1980) explained that the management practices like burning increased the total carbon accumulation in soil and in turn it increases the size and activities of soil microbial biomass. There was generally no direct loss of soil nutrients except for the volatilization of nitrogen and sulphur (Vogl 1974). With low fire intensities, however, these losses to atmosphere was probably minimal (Griffin and Friedel 1984). The nitrogen losses are often recovered through precipitation and increased actions of nitrogen fixing plants, particularly legumes (Burton 1972), soil algae, bacteria and certain fungi (Stewart 1967). The accumulation and rate of transfer of nutrients between the components of ecosystem is one of the most useful parameters to measure the community development (Odum 1971). Paulsamy *et al.* (1995a) found that the fire influenced grassland communities of Anaimalais, Western Ghats, India recorded significantly rapid uptake /release rate of certain nutrients like nitrogen, phosphorous and potassium.

The nutrient content of grassland plants on burned sites tended to be higher (Daubenmire 1968) or relatively unchanged (Lloyd 1971). Nielsen and Hole (1963) even noted symptoms of nitrogen and phosphorus deficiencies in unburned grassland plants. At pre-forest savanna of Ivory coast, Abbadie (1984) observed the maximum concentration of nitrogen in the aerial living parts of plants just after fires which decreased regularly thereafter upto the following fires. Shoot phosphorus also decreased in the second half of the growing season in the annual burned reed vegetation of Zuidelljk Fleveland polder which was coupled with an increase of

rhizome phosphorus (Linden and Van der 1986).

A few workers have questioned the increase of mineral pool in the rangelands after fire. In the Queensland grasslands, the stimulation of grass growth after burning was presumed to be due to higher soil temperatures rather to a release of available nutrients in the ash (Anderson and Pressland 1987). No evidence of any effect of fire on the phosphorus, potassium, calcium, magnesium, sodium and aluminium content of soils has been recorded after 28 years of veld burning at Ukulinga (Cass and Collins 1983). White and Grossman (1972) have reported a decline in calcium, magnesium, potassium and sodium concentrations and in the base saturation of soils in which the grassland was burnt annually for 37 years. According to Christensen (1977) there was no clear cut evidence that increased uptake of nutrients by plants occurred as a result of fire. Some increases of tissue concentration of nutrients were measured but they may simply have been a function of tissue age.

### FIRE A MANAGEMENT TOOL

Fire is primarily used on rangeland to manipulate plant populations, maintain habitat for animals, improve forage quality and production and increase fire proof vegetation preventing damaging wild-fires. Vegetation management which, while being of considerable antiquity, is still widely practiced at the present time. Burning of woody material, if properly done, has the added advantage of promoting rapid regeneration and encouraging the development of a new supply of young nutritious green shoots (Gimingham 1985).

The greatest potential use of fire will be in wildlife management since many wild-life preserves, refuges and hunting grounds still support native grasslands. Most of the national parks of Africa, for example, contain various grasslands and or savannas and have management programmes that include fire as an important and often essential tool (Hill 1971, Van Rensburg 1971). Komarek (1971) in summarizing the effects of fire on wildlife habitats in south-eastern United States pointed out that grasslands and early stages of brushland, maintained by controlled burning, provide the diversity of flora necessary for healthy wildlife populations. In Anaimalais of Western Ghats, Southern India, the forage yield of grasslands are increased drastically by fire which in turn is more useful for the benefit of wild animals (Paulsamy *et al.* 2003a). Hence, patch-wise rotational burning is advocated for sustained availability of fodder and habitat for wild animals in that areas. The use of fire for habitat manipulation for both wild and domestic animals is common, for example, deer and cattle in California (Hendricks 1968) and sheep and grouse in Britain (Ward 1972).

Fire has been shown to be an important tool in the management and control of bush encroachment in grazing rangelands (Bastin

1991). Major vegetative and edaphic changes are taking place in range ecosystems where fire has been excluded (Wright and Bailey 1982). Reduction in the occurrence of fire since European settlement is considered to be the major cause of shrub increase (Hodgkinson 1987). The tree and shrub invasion is occurring in the forest steppe area of Eurasia (Walter 1985), the aspen parkland of Canada (Bailey and Wroe 1974), the tall grass prairie of the Central United States (Bragg and Hulbert 1976) when prescribed fire is not regularly imposed. In the management of the grassland biome, fire is deliberately excluded only from those sites in which successional development to scrub or forest is being encouraged (Tainton and Mentis 1984).

The ecological status of Indian grasslands has been a debatable point. Champion (1936) considered that the climax vegetation of India would be either forests or deserts. He also expressed the opinion that nowhere in India is grassland of climax type and they are all existing due to biotic activities like lopping, burning, shifting, cultivation and grazing of the forests for the last several thousands of years. Singh and Krishnamurthy (1981) also concluded that the tropical grasslands of India are seral in nature, owing to their origin to deforestation or abandoned cultivation. They are maintained at various successional levels in different parts of the country mainly due to biotic factors like grazing, harvesting and man induced fire. The occurrence of annual summer fire every year in the Grass Hills of Anaimalais, Western Ghats maintained the ecosystem in a highly productive seral stage (Paulsamy *et al.* 2001). The differences in the habitat complex (including the edaphotope and macro- and micro-climate), age and mode of origin and intensity of biotic operations result in an array of very diverse grassland communities (Singh 1976), though, at places they may attain the status of a disclimax (Misra 1959).

Ranganathan (1938) studied the ecology of the shola grassland vegetation of the Nilgiris plateau and he believed that the Nilgiris grasslands are 'climatic climax' and these grasslands have been in possession of their territory for centuries and are entitled to be called stable association whatever the factors that ensure their stability might be. Bor (1938) opined that once we admitted the existence of grazing and burning in an area, we could not apply the term 'climatic climax' and so these grasslands should be considered as 'biotic climax'. Champion (1939) also felt that these grasslands are not climatic climaxes as they are caused by periodic fires.

Ecophysiological (ecothermotical) investigations of Indian grasslands are limited (Trivedi *et al.* 1981, Pandey 1988, Vinod Shankar and Trivedi 1988, Paulsamy 1992, Paulsamy *et al.* 1995a, b, 1996, 1997a, b, c, 2001). Working in a *Dichanthium annulatum* dominated grasslands, Pandey (1974a) found an increase in the live biomass of grasses and forbs as a result of burning. To this increase, he attri-

buted the fertilizing effect of nutrients released in ash. Some of the major consequences of fire were elimination or invasion of certain species, and alteration of the relative frequency and growth characteristics of certain species (Pandey 1974b). Summer burning prior to rains showed promising result for a higher and there by greater recovery rate of biomass compared to winter burning (Pandey 1988). The recurring fire during every summer in the grasslands of Anaimalais maintains the elevated biomass every year (Paulsamy *et al.* 1997b). Paulsamy *et al.* (1997c) further reported that the index of dominance was increased with concomitant decrease of species diversity index in the summer fire influencing *Chrysopogon zeylanicus* dominated grasslands of Anaimalais, India.

Mall and Mehta (1978) recorded a significant increase in the live green aboveground biomass following burning in a tropical grassland of Ujjain. On the other hand, the grazed and annually burnt *Sehima* grassland showed decreased annual average biomass, net community production and stability than that of the protected one (Vinod Shankar *et al.* 1975a). Nevertheless, in a *Sehima-Dichanthium* cover, the combination of burning and grazing proved to be a good tool for increasing the potentiality of range grasses as the burning removes the rank growth deposited near the sprouting buds and exposes those to environment while the grazing stimulates these buds and ultimately the vigour increases. The increase in plant height and tiller number of *S. nervosum* in every year grazing plot, may be the cause of stimulation of apical buds through animal trampling. The severe effect of these factors (burning and grazing) damages or destroys the buds resulting the decrease in plant vigour. Overall vigour of the species went down, if the grasslands were left unused as in control treatment (Trivedi *et al.* 1981). In Kundah rangelands of Nilgiris, Western Ghats, Senthilkumar *et al.* (1998) reported that the aboveground and litter productions and turnover rate of aboveground parts were increased significantly after summer fire. However, in their observation, belowground counter part remain unchanged after fire. In addition, they have noted the significant higher contribution of legume biomass to the total community after fire in the same grasslands.

In general, control of fire in rangelands used to improve the species diversity, well established food web and stability (Paulsamy 1992). Public involvement and awareness about the rangeland fire are more important for the effective control of fire. Netalkar (1997) in a case study at Bapeli region of Karnataka forests proved the effective control of fire by involving the local people. Paulsamy (2003) stated that ecological approaches must be necessary for the development of any fire control measures to protect the biodiversity and ecological balance as well. For the pre-suppression of fire, number of methods are practiced depending upon the environment. Show and Clarke (1994) explained the application of various fire breaks

like natural breaks, existing roads, trails and tracks, cleared fire breaks, tree covered breaks and fire lines for the pre-suppression of fire. Construction of vegetational fire breaks as fire barrier is getting vital importance in terms of biodiversity conservation and maintenance of ecological balance (Rasmussen *et al.* 1986, Perara 1992). In the high hills of Nilgiris of Southern India, vegetational fire breaks of 5 × 1.5 m size constructed by using certain evergreen shrubs like *Berberis tinctoria*, *Elaeagnus kolaga* and *Rhodomyrtus tomentosa* effectively control the spread of fire (Paulsamy *et al.* 2003c). In addition, they have evaluated that due to higher fire retardant capacity, the fire belt made by the species, *Rhodomyrtus tomentosa* was highly effective in comparison to other two species in the same area of Nilgiris, southern India.

## CONCLUSION

In tropical regions, the fire generally disturbs the natural development of the communities towards maturity, thus maintaining them at seral stage. These fires influenced disclimax vegetations support a rich fauna. However, uncontrolled wild fire often destroys vast tracts of grasslands within a short period, depriving fodder to the dependant herbivores. This can be averted by prescribed rotational burning, taking into account the phenology of the vegetation and the fodder needs of the fauna. Pre-seasonal controlled fire can also serve as an effective management tool to check the devastating wild fire. Suitable vegetational fire breaks may be employed to control and regulate the surface fire so as to preserve the seed pool, woody vegetation and biodiversity.

## LITERATURE CITED

- Abbadie, L. 1984. Evolution saisonniere du stock d' azote dans la strate herbacee d'une savane soumise au feu en Cote-d' Ivoire. *Acta Ecologica Ecol. Plant.* 5: 321-334.
- Abbadie, L. 1985. Evolution saisonniere du stock d'azote dans la strate herbacee d'une savane de CSte-d'Ivoire protegee des feux de brousse. *Acta Ecologica Ecol. Plant.* 6: 323-335.
- Abrams, M.D., A.K. Knapp and L.C. Hulbert. 1986. A ten-year record of above ground biomass in a Kansas tall grass prairie: Effects of fire and topographic position 73(10): 1509-1515.
- Andariese, S.W. and W.W. Covington. 1986. Changes in understory production for three prescribed burns of different ages in *Ponderosa* pine. *For. Ecol. Manage.* 14: 193-203.
- Anderson, E.R. and A.J. Pressland. 1987. Carefull burning usefull as a management tool. *Qld. Agric. J.* Jan-Feb: 40-44.
- Andrew, M.H. 1986. Use of fire for spelling monsoon tallgrass pasture grazed by cattle. *Trop. Grasslds.* 20: 69-78.
- Anderey, R. 1961. *African Genesis* Reprinted by Collins Fontana Books, London, 1970 (in West, 1971).
- Baily, A.W. 1988. Understanding fire ecology for range management. *In* Trueller, P.T.(ed.). *Vegetation science applications for rangeland analysis and management.* Kluwer Academic Publishers, Dordrecht, Boston, London.

- pp.527-557.
- Bailey, A.W. and R.A. Wroe. 1974. Aspen invasion in a portion of the Alberta parklands. *J. Range Manage.* 27: 263-266.
- Bastin, G. 1991. Rangeland reclamation at Ataritinga Station, Central Australia. *Aust. J. Soil Water Conserv.* 4: 18-25.
- Bor, N.L. 1938. The vegetation of the Nilgiris. *Ind. For.* 64: 600-609.
- Bragg, T.B. and L.C. Hulbert. 1976. Woody plant invasion of unburned Kansas bluestem prairie. *J. Range Manage.* 29: 19-24.
- Britton, C.M., H.A. Wright, B.E. Dahl and D.N. Ueckert. 1987. Management of tobosagrass rangeland with prescribed fire. Management Note 12, contribution No. T-9-481, College of Agricultural Sciences, Texas Tech. University.
- Burton, J.C. 1972. Nodulation and symbiotic nitrogen fixation by prairie legumes. *Midwest Prairie Conf.*, 2nd, 1970, pp. 116-121.
- Cass and Collins. 1983. Fire in grassland. *In* Boosen, P.deV. and N.M. Tainton (eds.). *Ecological effects of fire in South African ecosystems.* Springer-Verlag, Berlin. pp.116-147.
- Casson, N.E. and J.E.D. Fox. 1987. The post fire regeneration responses of *Triodia wiseana* and *T. basedowii*. *Aust. Rangel. J.* 9: 53-55.
- Champion, H.G. 1936. A preliminary survey of the forest types of India and Burma. *Indian For. Rec. (N.S.) Silv.* 236. p.
- Champion, H.G. 1939. The relative stability of Indian vegetational types. *J. Indian Bot. Soc.* 18: 1.
- Christensen, N.L. 1977. Fire and soil-plant nutrient relations in a pine-wire grass savanna on the coastal plains of North Carolina. *Oecologia* 31: 27-44.
- Curl, E.A. and B. Truelove. 1986. *The rhizosphere.* Springer-Verlag, Berlin.
- Daubenmire, R. 1968. Ecology of fire in grassland. *Adv. Ecol. Res.* 5: 209-266.
- Fox, J.E.D. 1980. Effects of fire on the mulga (*Acacia aneura*) community. *Mulga Res. Cent. J.* 3: 1-19.
- Garza, N.E. and W.H. Blackburn. 1985. The effect of early winter or spring burning on runoff, sediment, and vegetation in the post oak Savannah of Texas. *J. Range Manage.* 38: 283-287.
- Gimingham, C.H. 1985. *Muriburn.* BCPC Mono. No.30. Vegetation management in northern Britain.
- Gimingham, C.H. 1989. Heather and heathlands. *Bot. Linn. Soc.* 101: 263-268.
- Griffin, G.F. and M.H. Friedel. 1984. Effects of fire on central Australian rangelands. I. Fire and fuel characteristics and changes in herbage and nutrients. *Aust. J. Ecol.* 9: 381-393.
- Griffin, G.F. and M.H. Friedel. 1985. Discontinuous change in central Australia: some implications of major ecological events for land management. *J. Arid. Environ.* 9: 63-80.
- Harniss, R.O. and R.S. Murray. 1973. Thirty years of vegetal change following burning of sagebrush-grass range. *J. Range Manage.* 26: 322-325.
- Hassan, M.A. and N.E. West. 1986. Dynamics of soil seed pools in burned and unburned sagebrush semi-deserts. *Ecology* 67: 269-272.
- Hendricks, J.H. 1968. Control burning for deer management in chaparral in California. *Proc. 8th Annu. Tall Timbers Fire Ecol. Conf.* pp. 219-233.
- Hill, P. 1971. Grass forage, food for fauna or fuel for fire or both? *Proc. 11th Annu. Tall Timbers Fire Ecol. Conf.* pp. 337-375.
- Hobbs, N.T. and D.S. Schimel. 1984. Fire effects on nitrogen mineralization and fixation in mountain shrub and grassland communities. *J. Range Manage.* 37: 402-405.
- Hodgkinson, K.C. 1986. Response of rangeland plants to fire in water-limited environments. *In* Joss, P.J., P.W. Lynch and O.B. Williams (eds.) *Rangelands: A resource under siege.* Proc. 2nd Int. Rangeland Congr., Australian Academy of Science, Canberra, 1986.
- Hodgkinson, K.C. 1987. Shrub and herbage changes under prescribed burning. *In* Knight, F. (ed.). *Division of Wildlife and Rangelands Research (CSIRO) Report, 1984-86.* Commonwealth Scientific and Industrial Research Organization, Australia. pp.68.
- Jalaludin, M. 1969. Microorganic colonization of forest soil after burning. *Plant Soil* 30: 150-152.
- Joshi, S.C. 2003. Impact of forest fires on the regional climate. *Current Sci.* 85(1): 41-45.
- Komarek, E.V. 1971. Effects of fire on wildlife and range habitats. *Proc. Prescribed Burn. Symp.* pp. 46-52.
- Linden, M.J. and H.A. Van der. 1986. Phosphorus economy of reed vegetation in the Zuidelijk Flevoland polder (The Netherlands): seasonal distribution of phosphorus among shoots and rhizomes and availability of soil phosphorus. *Acta Ecologica Ecol. Plant* 7: 397-405.
- Lloyd, P.S. 1971. Effects of fire on the chemical status of herbaceous communities of the Derbyshire Dales. *J. Ecol.* 59: 261-273.
- Lynch, J.M. and L.M. Panting. 1980. Cultivation and the soil biomass. *Soil Biol. Biochem.* 12: 29-33.
- Mall, L.P. and S.C. Mehta. 1978. The grassland fire: an ecological study. *In* Singh, J.S. and B. Gopal (eds.) *Glimpses of Ecology (R. Misra commemoration volume).* International Scientific Publisher, Jaipur, India. pp.291-297.
- Manian, S., S. Paulsamy, K. Senthilkumar and B.S. Kil. 2002. Impact of surface fire on the dynamics of N<sub>2</sub>-fixing and P- solubilizing microbial population in natural grassland soils, southern India. *Korean J. Ecol.* 25: 227-234.
- Mayeux, Jr. H.S. and W.T. Hamilton. 1983. Response of Common Golden weed (*Isocoma coronopifolia*) and Buffelgrass (*Cenchrus ciliaris*) to fire and soil-applied herbicides. *Weed Sci.* 31: 355-360.
- Misra, R. 1959. The status of plant community in upper Gangetic plain. *J. Indian bot. Soc.* 38: 1-7.
- Morrison, L.C., J.D. DuBois and L.A. Kapustka. 1986. The vegetational response of a Nebraska Sandhills Grassland to a naturally occurring fall burn. *Prairie Nat.* 18: 179-184.
- Netalkar, P.S. 1997. A case study of fire protection in Bapeli village. *My Forest* 33(3): 577-579.
- Nielsen, G.A. and F.D. Hole. 1963. A study of the natural processes of incorporation of organic matter into soil in the University of Wisconsin Arboretum. *Trans. Mis. Acad. Sci. Arts Lett.* 52: 213-227.
- Noble, J.C. 1989. Fire studies in mallee (*Eucalyptus* spp.) communities of Western New South Wales: The effects of fires applied in different seasons on herbage productivity and their implications for management. *Aust. J. Ecol.* 14: 169-187.
- Noble, J.C., G.N. Harrington and K.C. Hodgkinson. 1986. The ecological significance of irregular fire in Australian rangelands. *In* P.J. Joss, P.W. Lynch and O.B. Williams (eds.). *Rangelands: A Resource under siege,* Aust. Acad. Sci., Canberra. pp.577-580.
- Odum, E.P. 1971. *Fundamentals of Ecology.* (3rd ed.) W.B. Saunders Co., Philadelphia and London, pp.574.
- Pandey, A.N. 1974a. Short-term effect burning on the above ground production of *Dichanthium annulatum* grassland stands at Varanasi. *Trop. Ecol.* 15: 152-154.
- Pandey, A.N. 1974b. Short-term effect of seasonal burning on the composition of *Dichanthium annulatum* grassland stands at Varanasi. *Proc. Indian Natn. Sci. Acad.* 40: 107-112.
- Pandey, A.N. 1988. Short-term study of recovery of a tropical grassland following seasonal burning. *Trop. Ecol.* 29: 159-170.
- Paulsamy, S. 1992. Role of fire in the maintenance of *Chrysopogon zeylanicus* The. dominated grassland of Grass Hills, Anaimalais, the Western Ghats, India. Ph.D Thesis, Bharathiar University, Coimbatore, India.
- Paulsamy, S. 2003. Forest fire: a socio-ecological problem. *SAJOSPS.* Jan-June: 101-105.
- Paulsamy, S., D. Senthilkumar, S. Manian and K. Arumugasamy. 1997a. Impact of annual summer fire on some sociological characters of *Anaphalis subdecurrens* Gamble in a *Chrysopogon zeylanicus* The dominated grassland. *Geobios New Reports* 16: 113-116.
- Paulsamy, S., K. Arumugasamy and T.N. Rangarajan. 1997b. Effect of annual summer fire on the primary production in Grass Hills ecosystem, Western Ghats. *Geobios* 24 (2-3): 151-154.

- Paulsamy, S., R. Sivakumar, V. Balasubramanian, K. Arumugasamy and N. Nagarajan. 2001. Fire control - a conservation tool for certain medicinal plants in Grass Hills ecosystem, the Western Ghats. *Ancient Science of Life* 20 (4): 89-92.
- Paulsamy, S., S. Manian and S. Padmavathy. 2003a. Rotational burning-a tool for sustainable forage production for wild life in Grass Hills, Western Ghats. *Range Mgmt. & Agroforestry* 24 (1): 14-17.
- Paulsamy, S., S. Manian and K. Udaiyan. 1995a. A strategy to evaluate community dynamics through uptake, release rate of nutrients. *Range Mgmt. Agroforestry* 16 (1): 15-20.
- Paulsamy, S., S. Manian and K. Udaiyan. 1995b. Relationship in the response of a grass, *Chrysopogon zeylanicus* and a fern, *Pteris aquilina* to annual summer fire on Grass Hills ecosystem, Western Ghats. *Indian J. Forestry* 18 (3): 22-225.
- Paulsamy, S., S. Manian, K. Udaiyan, K. Arumugasamy, N. Nagarajan and B.S. Kil. 2003b. Biomass structure and dry matter dynamics in a fire influencing montane subtropical humid grassland, Western Ghats, southern India. *Korean J. Ecol.* 24: 227-232.
- Paulsamy, S., S. Manian, T.N. Rangarajan, K. Udaiyan and K. Arumugasamy. 1997c. Impact of fire on certain structural and functional relationships in *Chrysopogon zeylanicus* Thw. dominated grassland, Western Ghats. *Range Mgmt. & Agroforestry* 18 (2): 133-141.
- Paulsamy, S., S. Manian and K. Udaiyan. 1993. Influence of annual summer fire on soil nutrient pool in *Chrysopogon zeylanicus* dominated grasslands of Western Ghats, India. *Indian Bot. Contactor* 10: 141-146.
- Paulsamy, S., T.N. Rangarajan and K. Arumugasamy. 1996. Short-term studies on the ecological status of weeds in an annual fire influenced *Chrysopogon zeylanicus* dominated grassland, Western Ghats. *World - Weeds* 3: 67-75.
- Paulsamy, S., T.N. Rangarajan, S. Manian and K. Udaiyan. 1995. Impact of annual summer fire on the sociological attributes of the fern, *Pteridium aquilinum* L. in the grass hills ecosystems, Western Ghats, India. *J. Econ. Tax. Bot.* 19 (3): 745-749.
- Paulsamy, S., B.-S. Kil, K. Arumugasamy and S. Padmavathy. 2003c. Strategy to control surface fire using vegetational fire breaks in rangelands of Nilgiris, southern India. *Korean J. Ecol.* 26(5) : 251-254.
- Perara, A.H. 1992. Ecological implication of establishing pine plantations in the Central Region of Sri Lanka. In K.P. Singh and J.S. Singh (eds.) *Tropical Ecosystem: Ecology and Management*, pp. 409-414.
- Pressland, A.J., D.C. Cowan, C.J. Evenson and P.S. Bowly. 1986. Benefits of infrequent fire in the mulga (*Acacia aneura*) rangelands of Queensland. In Joss, P.J., P.M. Lynch and O.B. Williams (eds.) *Rangelands: A resource under siege*, Proc. Rangeland Congr., Australian Academy of Science, Canberra. pp. 608-609.
- Ranganathan, C.R. 1938. Studies in the ecology of shola-grassland vegetation of the Nilgiri Plateau. *Ind. For.* 64 (9).
- Rasmussen, G.A., G.R. Mcpherson and H.A. Wright. 1986. Prescribed burning of juniper communities in Texas. Management Note 10. Range and Wild Life Management. Texas Tech. University, Lubbock, Texas.
- Senthilkumar, K., S. Manian, K. Udaiyan and S. Paulsamy. 1998. Elevated biomass production in burned natural grasslands in southern India. *Tropical Grasslands* 32: 50-63.
- Senthilkumar, K., S. Manian, K. Udaiyan and V. Sugavanam. 1995. Effect of burning on soil nutrient status and abundance of VA - mycorrhizal fungi in a savannah type grassland ecosystem in south India. *Tropics* 4(10): 173-186.
- Shaw, N.H. 1957. Bunch spear grass dominance in burnt pastures in south eastern Queensland. *Aust. J. Agric. Res.* 8 : 325-334.
- Show, S.B. and B. Clarke. 1994. Forest fire control. International Book Distributors, Dehradun, India. p.110.
- Singh, J.S. 1976. Structure and function of tropical grassland vegetation of India. *Pol. . Ecol. Stud.* 2: 17-34.
- Singh, J.S. and L. Krishnamurthy. 1981. Analysis of structure and function of tropical grassland vegetation of India. *Indian Rev. Life Sci.* 1: 225-270.
- Snaydon, R.W. 1987. Fertilizer inputs and botanical composition. In R.W.Snaydon, (ed.). *Managed grasslands, B. Analytical Studies.* Elsevier Science Publishers B.V., Amsterdam, pp.239-246.
- Stewart, W.D.P. 1967. Nitrogen fixing plants. *Science* 158: 1426-1432.
- Tainton, N.M and M.T. Mentis. 1984. Fire in grassland. In P. Boosen, dev. and N.M. Tainton (eds.). *Ecological effects of fire in South African ecosystems*, Springer-Verlag, Berlin. pp. 115-147.
- Tothill, J.C. 1969. Soil temperatures and seed burial in relation to the performance of *Heteropogon contortus* and *Themeda australis* in burnt native woodland pastures in eastern Queensland. *Aust. J. Bot.* 17: 269-275.
- Trivedi, B.K., K.C. Kanodia and Vinod Shanker. 1981. Effect of burning and grazing on the vigour of range grasses of *Schima-Dichanthium* cover. *Geobios* 8: 13-17.
- Trollope, W.S.W. 1990. Veld management with specific reference to game ranching in the grassland and Savanna areas of south Africa. *Koedoe* 33(2):77-86.
- Van Rensburg, H.J. 1971. Fire: Its effects on grasslands, including swamps-southern, central and eastern Africa. Proc. 11<sup>th</sup> Annu. Tall Timbers Fire Ecol. Conf. pp 175-199.
- Vinod Shankar and B.K. Trivedi. 1988. Grazing-burning rotation for the management of Schima-Heteropogon grasslands in Central India. In Panjab Singh *et al.* (eds.) *Third Int. Rangeland Congr. Abstr. Vol.1. RMSI, IGFR, Jhansi, India.* pp.382-385.
- Vinod Shankar., K.C. Velayudhan and B.K. Trivedi. 1975a. Pattern of interspecific association in *Schima-Dichanthium* grassland under three different system of management. *Ann. Arid Zone* 14(4): 358-366.
- Viro, P.J. 1974. Effects of forest fire on soil. In T.T. Kozlowski and C.E. Ahlgren (eds). *Fire and ecosystems.* Academic Press, New York, London. pp.7-72.
- Vogl, R.J. 1974. Effects of fire on grasslands. In T.T. Kozlowski and C.E. Ahlgren (eds). *Fire and ecosystems.* Academic press, New york, London. pp. 528.
- Walter, H. 1985. *Vegetation of the Earth*, 3<sup>rd</sup> ed. Springer-Verlag, Berlin.
- Ward, S.D. 1972. The controlled burning of heather, grass and gorse. *Nature (Wales)* 13: 24-32.
- Wells, C.G., R.E. Campbell, L.F. DeBano and five other authors. 1979. Effects of fire on soil. USDA For. Serv. Gen. Tech. Rep. WO-7.
- West, O. 1971. Fire, man and wildlife as interacting factors limiting the development of vegetation in Rhodesia- Proc. 11<sup>th</sup> Annu. Tall Timbers Fire Ecol. Conf- pp. 121-146.
- Whisenant, S.G. and W.R. Bulsiewicz. 1985. Effects of prescribed burning on Japanese brome population dynamics. Proc. XV Int. Grassland Congr., Kyoto, Japan. The Science Council of Japan and The Japanese Society of Grassland Science.
- White and Grossman, 1972. Fire in Grassland. In C. Boosen. P.deV and N.M. Tainton (eds.) *Ecological effects of fire in South African Ecosystems.* Springer-Verlag, Berlin. pp.116-147.
- Wright, H.A. and A.W. Bailey. 1982. *Fire ecology of the United States and southern Canada.* John Wiley & Sons, New York.
- Wright, H.A. 1985. Effects of fire on grasses and forbs in sagebrush grass communities. In Sanders, K. and J. Durham (eds.). *Rangeland Fire Effects.* Idaho State Office, USDI-Bureau of Land Management, Boise, Idaho. pp.12-21.

(Received July 30, 2004; Accepted December 29, 2004)