

## Bacterial Diversity in the Rhizosphere of Halophyte *Phragmites communis* at the Western Coastal Mudflats of Korea

Moon, Ho-Sang<sup>1</sup>, Suhk-Hwan Park<sup>1</sup>, Jong-Ok Ka<sup>2</sup>, Hong-Gyu Song<sup>3</sup> and Geon-Hyoung Lee<sup>1\*</sup>

<sup>1</sup>Department of Biology, Kunsan National University, Gunsan 573-701, Korea

<sup>2</sup>Department of Agricultural Biotechnology, Seoul National University, Seoul 151-742, Korea

<sup>3</sup>Division of Biological Sciences, Kangwon National University, Chuncheon 200-701, Korea

**ABSTRACT:** This study investigated the population densities and diversity of heterotrophic bacteria, and the rhizosphere-to-soil ratios (R/S) in the rhizosphere soil of halophyte *Phragmites communis* at the western coastal mudflats of Korea. The population densities of aerobic heterotrophic bacteria on the rhizosphere soil of *P. communis* were in the range of  $3.3 \pm 0.9 \times 10^7 \sim 1.2 \pm 0.5 \times 10^8$  cfu g<sup>-1</sup> dry weight (d. wt.). Population densities of amylolytic bacteria ranged from  $1.1 \pm 0.2 \times 10^6$  to  $3.0 \pm 1.2 \times 10^6$  cfu g<sup>-1</sup> d. wt., while those of cellulolytic bacteria and proteolytic bacteria ranged from  $5.6 \pm 2.3 \times 10^6$  to  $1.5 \pm 0.3 \times 10^7$  cfu g<sup>-1</sup> d. wt. and from  $1.4 \pm 0.3 \times 10^6$  to  $3.5 \pm 2.3 \times 10^7$  cfu g<sup>-1</sup> d. wt., respectively. The R/S ratios ranged from 2.26 to 6.89. Genetic (16S DNA) analysis of fifty-one isolates from the roots of *P. communis* suggested that the dominant species were closely related to the  $\gamma$ -proteobacteria group (18 clones) and the  $\alpha$ -proteobacteria group (14 clones). We found that halophyte species and mudflat environment both affected the rhizosphere bacterial communities.

**Key words:** 16S rDNA, Bacterial population density, Halophyte, *Phragmites communis*,  $\gamma$ -proteobacteria group, R/S ratio, Rhizosphere

### INTRODUCTION

Among the 2,815 km<sup>2</sup> of mudflats in South Korea, 2,393 km<sup>2</sup> are found on the western and southern coasts of Korea. Mudflats in coastal estuaries act as protective filters and final repositories for runoff pollutants (Teal and Howes 2000), pathogens (Grant et al. 2001), and nutrients (Howes et al. 1996). They also play a vital role in the development of many marine populations and are essential as breeding grounds or nursery areas for many species.

Halophytes in mudflat areas function as primary producers, and their roots and stalks stabilize the sediments, preventing erosion. They also provide habitats for various organisms and have the ability to remove contaminants from ecosystem (Choi and Lee 1996). However, because of recent large-scale land reclamation projects, halophyte habitats have been reduced rapidly, threatening the functioning of coastal ecosystems.

Rhizospheres, generally defined as the soil adjacent to and influenced by plant roots, are regarded as "hot spots" for microbial colonization and activity (Metting 1993). In contrast to bulk soil, where organic carbon is available only at low concentrations, rhizospheres are supplied with higher concentrations of nutrients generated during plant photosynthesis (Duineveld et al. 1998, 2001).

Rhizosphere microbial communities are mainly determined by the plant species (Marshner et al. 2001) and soil characteristics (Degens et al. 2000, Gelsomino et al. 1999), and rhizosphere microbial communi-

ties influence plant nutrition, growth, and disease (Assigbetse et al. 2005).

Microbiological studies of mudflat environments in Korea have mainly investigated microbe distribution (Lee et al. 1996), enzyme activities (Kim and Lee 1992, Choi and Lee 1996), and diversity (Lee et al. 2001, Kim et al. 2004, Kim et al. 2005). However, microbial diversity on the rhizosphere of halophytes in mudflat environments in Korea was also studied recently by Park and Lee (2006).

In this study, we investigated the R/S ratios, population densities, and diversities of bacteria in the rhizosphere of *Phragmites communis* communities on mudflats in western Korea as baseline data for a restoration project for halophyte habitats in Korea.

### MATERIALS AND METHODS

#### Sampling and Counting of Bacteria

Samples of rhizosphere soil (R) and soil remote from roots (S) of *P. communis* were collected from 3 stations on the western coastal mudflats of Korea during July and October, 2004 (Fig. 1). Soils were collected using a soil auger, stored at 5 °C, and processed within a few hours of collection.

The effects of the rhizosphere were measured by calculating at the ratio of the number of microorganisms in rhizosphere soil (R) to the number of microorganisms in soil remote from roots (S), or the R/S ratio.

\* Corresponding author; Phone: +82-63-469-4584, e-mail: ghlee@kunsan.ac.kr




Fig. 1. Map showing the sampling sites in the western mudflats of Korea (St. 1: Seocheon, St. 2: Mankyung R., St. 3: Muan bay).

We estimated of population densities following the methods of Paul and Clark (1988). To determine R/S ratios and the population densities of the aerobic heterotrophic bacteria, one gram of rhizosphere soil and one gram of soil remote from roots were suspended in 10-mL sterile saline solution (0.85% NaCl), and shaken for 5 min at 100 rpm. Then serial decade dilutions were made with sterile saline water and 0.1 mL of each was plated on Nutrient agar (Difco) and Marine agar 2216 (Difco). To determine the number of aerobic physiological groups of heterotrophic bacteria, soluble starch (0.2%) for amylolytic bacteria, carboxymethyl cellulose (0.5%) for cellulolytic bacteria, or gelatin (0.4%) for proteolytic bacteria was added as the sole carbon source to the Tryptic soy broth (Difco, USA) as the basal culture medium (Wollum 1982). After incubation at  $25 \pm 2^\circ\text{C}$  for 72 hrs, we counted the resulting colonies using the methods of Holding and Collee (1971). Final estimated population densities were expressed as  $\log_{10}$  colony-forming units (CFU)  $\text{g}^{-1}$  oven-dried sediment.

#### PCR Amplification of 16S rDNA

We amplified 16S rDNA fragments of isolates from the rhizosphere of the halophyte *P. communis* using PCR with the primers 27F (5'-AGAGTTTGTATCMTGGCTCAG-3') and 1522R (5'-AAGGAGGT

GWTCARCC-3'). The PCR reaction mixture contained 5  $\mu\text{L}$  of 10X PCR amplification buffer (final concentration: 50 mM KCl, 0.01% gelatin, 10 mM Tris-HCl, pH. 9.0), 4  $\mu\text{L}$  of 2.5 mM  $\text{MgCl}_2$ , 1  $\mu\text{L}$  of 10 mM dNTP, 1  $\mu\text{L}$  of each 10 pmol oligonucleotide primer, and 1 U of Taq polymerase (TaKaRa, Japan) in 50  $\mu\text{L}$  of PCR mixture. DNA was amplified with a GeneAmp PCR system 2700 (Applied Biosystems, USA) thermal cycler using the following program: initial denaturation at  $94^\circ\text{C}$  for 10 min, followed by 30 cycles of denaturation at  $94^\circ\text{C}$  for 30 sec, annealing at  $55^\circ\text{C}$  for 30 sec, extension at  $72^\circ\text{C}$  for 5 min, and a final extension at  $72^\circ\text{C}$  for 7 min. PCR products were either used immediately or stored at  $4^\circ\text{C}$  prior to subsequent analyses. Two replicate reactions were run for each sample.

#### 16S rDNA Sequencing and Phylogenetic Analysis

Ribosomal DNA sequences were determined by the Genotech Company (Daejeon, Korea) using an ABI PRISM 3700 DNA analyzer (Applied Biosystem, USA). We then compared the sequences directly to all known sequences in the GenBank database using the basic local alignment search tool (BLAST)(Altschul et al. 1997), and constructed phylogenetic trees with the neighbor-joining (NJ) method using the NEIGHBOR program (PLYLIP, version 3.5) (Saitou and Nei 1987).

## RESULTS AND DISCUSSION

#### Population Densities of Heterotrophic Bacteria

Estimated population densities of aerobic heterotrophic bacteria inhabiting the rhizosphere of *P. communis* ranged from  $3.3 \pm 0.9 \times 10^7$  to  $1.2 \pm 0.5 \times 10^8$  cfu  $\text{g}^{-1}$  dry weight (d. wt.) during the sampling periods (Fig. 2). The population densities of aerobic heterotrophic bacteria were the highest on the mudflats near Mankyung River, and lower near Seocheon and Muan. Although the differences in microbial population density among sampling stations were not explored in detail in this study, we suspect that bacterial population densities may be influenced by fine-scale environmental variables in the soil.

Previous studies have reported diversity in the microbial populations in the rhizospheres of different plants (Neal et al. 1970, 1973). In our study the population densities of aerobic heterotrophic bacteria were higher in the rhizosphere of *P. communis* than those reported for the *Suaeda japonica* rhizosphere (Park 2004), but were lower than those in the *Suaeda maritima* and *Salicornia herbacea* rhizospheres, which suggests that rhizosphere microbial communities are mainly determined by root exudates, which vary among plant species (Marshner et al. 2001). Soil microorganisms have been shown to respond to plant exudates and different plant species have different patterns of root exudation (Brimecombe et al. 2001).

Heterotrophic bacteria on the mudflats decompose organic matter derived from plant litter and transform pollutants (Benoit et al. 2003, Smith and Hollibaugh 1993). The population densities of physiological groups of heterotrophic bacteria on the roots of *P. communis* ranged




Fig. 2. Population densities of (a) aerobic heterotrophic bacteria, (b) amylolytic heterotrophic bacteria, (c) cellulolytic heterotrophic, and (d) proteolytic bacteria in the rhizosphere soil and soil remote from roots of the halophyte *Phragmites communis* from mudflats in western Korea in July and October, 2004.

from  $1.1 \pm 0.2 \times 10^6$  to  $3.0 \pm 1.2 \times 10^6$  cfu g<sup>-1</sup> d. wt. for amylolytic bacteria, from  $5.6 \pm 2.3 \times 10^6$  to  $1.5 \pm 0.3 \times 10^7$  cfu g<sup>-1</sup> d. wt. for cellulolytic bacteria, and from  $1.4 \pm 0.3 \times 10^6$  to  $3.5 \pm 2.3 \times 10^7$  cfu g<sup>-1</sup> d. wt. for proteolytic bacteria (Fig. 2). Population densities were similar among the different physiological groups of heterotrophic bacteria. However, during the sampling period, bacterial population densities of amylolytic and proteolytic bacteria were highest at Mankyung, while densities of cellulolytic bacteria were highest and densities of amylolytic bacteria were lowest at Muan. It is likely that the materials released by plants into the soil (Atlas and Bartha 1992) had a direct influence on the composition and density of the mudflat microbial community. Accordingly, root exudation is an important ecological phenomenon that affects succession in the plant and root microbial communities (Singh and Mukerji 2006).

### Rhizosphere Effects on Microbial Populations

The rhizosphere is chemically, physiologically and biologically complex due to the influence of plant roots on various types of microorganisms. The interaction of soil and rhizosphere microbes results in

stimulation of microorganisms known as the "rhizosphere effect".

The R/S ratios in the rhizospheres of *P. communis* ranged from 2.26 to 6.89 (Table 1), which are similar to the R/S values reported for rhizospheres of other halophytes (Park 2004), but lower than those of common terrestrial environments (Gray and Parkinson 1968, Woldendorp 1978). This result suggests that nutrient availability and soil texture have an important effect on the growth of rhizosphere bacteria

Table 1. Ratios of the number of microorganisms in the rhizosphere soil (R) to the number of corresponding microorganisms in soil remote from roots (S) (R/S ratios) of the halophyte *Phragmites communis* from mudflats in western Korea

Station	Halophyte <i>Phragmites communis</i> (R/S Ratio)
Seocheon	2.26
Mankyung	6.30
Muan	6.89




Fig. 3. Phylogenetic tree showing the affiliation of 16S rDNA sequences of heterotrophic bacteria sampled from the rhizosphere of *Phragmites communis* from Seocheon to selected reference sequences. The tree was constructed from a distance matrix by the neighbour-joining analysis. The bar represents 0.1% estimated sequence divergence.




Fig. 4. Phylogenetic tree showing the affiliation of 16S rDNA sequences of heterotrophic bacteria sampled from the rhizosphere of *Phragmites communis* from Mankyung to selected reference sequences. The tree was constructed from a distance matrix by the neighbour-joining analysis. The bar represents 0.1% estimated sequence divergence.




Fig. 5. Phylogenetic tree showing the affiliation of 16S rDNA sequences of the heterotrophic bacteria on the rhizosphere of *Phragmites communis* sampled from Muan to selected reference sequences. The tree was constructed from a distance matrix by the neighbour-joining analysis. The bar represents 0.1% estimated sequence divergence.

on mudflats, and therefore on R/S values.

### Sequencing and Phylogenetic Analysis

Bacterial 16S rDNA clones were characterized by partial sequencing and phylogenetic analysis. We identified thirteen isolates from the rhizosphere of *P. communis* from Seocheon: five clones, including *Burkholderia* spp., *Alcaligenes* sp., *Varovorax* sp., and *Ralstonia* sp., were determined to belong to the  $\beta$ -proteobacteria group and four clones, including *Pseudomonas* spp. and *Acinetobacter* sp. were determined to belong to the  $\gamma$ -proteobacteria group (Table 2). Among the twenty-six isolates from Mankyung, twelve clones were determined to belong to the  $\gamma$ -proteobacteria group, and ten were determined to belong to the  $\alpha$ -proteobacteria group (Table 2). We identified twelve isolates from Muan, six of which were determined to belong to the  $\gamma$ -proteobacteria

Table 2. List of heterotrophic bacteria isolated by 16S rDNA analysis from the rhizosphere of the halophyte *Phragmites communis* from mudflats in western Korea

Site	Closest Genebank library	Group
	<i>Flavobacterium</i> sp.	Flavobacteria
	<i>Bacillus</i> spp. (2 species)	Firmicutes
	<i>Varovorax</i> sp.	Betaproteobacteria
	<i>Pseudomonas</i> spp. (3 species)	Gammaproteobacteria
Seocheon	<i>Burkholderia</i> spp. (2 species)	Betaproteobacteria
	<i>Alcaligenes</i> sp.	Betaproteobacteria
	<i>Ralstonia</i> sp.	Betaproteobacteria
	<i>Micrococcus</i> sp.	Actinobacteria
	<i>Acinetobacter</i> sp.	Gammaproteobacteria
	<i>Cellulomonas</i> sp.	Actinobacteria
	<i>Sphingomonas</i> sp.	Alphaproteobacteria
	<i>Aeromonas</i> spp. (2 species)	Gammaproteobacteria
	<i>Pseudomonas</i> spp. (5 species)	Gammaproteobacteria
	<i>Streptomyces</i> sp.	Actinobacteria
	<i>Sphingomonas</i> sp.	Alphaproteobacteria
	<i>Microbacterium</i> sp.	Actinobacteria
Mankyung	<i>Paracoccus</i> spp. (4 species)	Alphaproteobacteria
	<i>Erythrobacter</i> spp. (2 species)	Alphaproteobacteria
	<i>Xanthomonas</i> sp.	Gammaproteobacteria
	<i>Citricella</i> sp.	Alphaproteobacteria
	<i>Sulfitobacter</i> sp.	Alphaproteobacteria
	<i>Microbulbifer</i> sp.	Actinobacteria
	<i>Oceanospirillum</i> sp.	Gammaproteobacteria
	<i>Shewanella</i> spp. (2 species)	Gammaproteobacteria
	<i>Acinetobacter</i> sp.	Gammaproteobacteria
	<i>Erwinia</i> spp. (4 species)	Gammaproteobacteria
	<i>Hyphomonas</i> spp. (3 species)	Alphaproteobacteria
	<i>Shewanella</i> sp.	Gammaproteobacteria
Muan	<i>Bacillus</i> sp.	Firmicutes
	<i>Arthrobacter</i> sp.	Actinobacteria
	<i>Pseudomonas</i> sp.	Gammaproteobacteria
	<i>Stappia</i> sp.	Alphaproteobacteria

group, and four to the  $\alpha$ -proteobacteria group (Table 2). Among the total of fifty-one isolates from the three sampling stations, the dominant group were the  $\gamma$ -proteobacteria (43.1%), followed by the  $\alpha$ -proteobacteria (27.5%), and the Actinobacteria (11.8%). According to Gray and Herwing (1996), the  $\gamma$ -proteobacteria group is dominant in marine sediments. The microorganisms prevailing in the rhizosphere originate in the soil. However, because of variation in root exudates characteristics between plant species, the specific organisms constituting the rhizosphere microbial community vary with both plant species and soil type (Grayston et al. 1998, Latour 1996, Maloney et al. 1997). In our study, most sequences had < 97% sequence similarity to previously cultivated microorganisms and phylogenetic analyses of isolated bacteria from the rhizosphere of *P. communis* revealed that the majority of bacteria detected were closely related to the  $\gamma$ -proteobacteria and the  $\alpha$ -proteobacteria. However, further study will be required to better understand the diversity, density, and functions of rhizosphere bacteria of halophytes on mudflat environments.

We found higher bacterial densities in the *P. communis* rhizosphere than in soils remote from the rhizosphere. Variation in rhizosphere bacterial populations among localities and plant communities suggests that halophyte species and soil type affects the rhizosphere bacterial communities in mudflat environments (Grayston et al. 1998, Latour 1996, Maloney et al. 1997).

#### ACKNOWLEDGMENTS

This study was partially supported by a grant from the Marine Development Research Center at Kunsan National University.

#### LITERATURE CITED

- Altschul SF, Madden TL, Schoëffer AA, Zhang J, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 5: 3389-3402.
- Assigbetse K, Gueye M, Thioulouse J, Duponnois R. 2005. Soil bacterial diversity response to root colonization by an ectomycorrhizal fungus are not root-growth-dependent. *Microb Eco* 50: 350-359.
- Atlas RM, Bartha R. 1992. *Microbial Ecology: Fundamentals and Application*, 2<sup>nd</sup> Ed. Benjamin/Cummings Publishing Co., Redwood City, CA. pp 69-74.
- Benoit JM, Gilmour CC, Heyes A, Mason RP, Miller CL. 2003. Geochemical and biological controls over methylmercury production and degradation in aquatic ecosystems. *ACS Symp Ser* 835: 262-297.
- Brimecombe MJ, de Leij FA, Lynch JM. 2001. The effect of root exudates on rhizosphere microbial communities. In: *The Rhizosphere: Biochemistry and Organic Substances at Soil-Plant Interface* (Pinton R, Varanini Z, Nannipieri P, eds). Marcel Dekker, New York, pp 95-140.
- Choi GK, Lee GH. 1996. Interaction between saprophytic bacterial distribution and their extracellular enzyme activities in the sediment of the Yellow sea near Seocheon. *The Microorganisms and Industry* 22: 119-126.
- Degens BP, Schipper LA, Sparling GP, Vojvodic-Vukovic M. 2000. Decreases in organic C reserves in soils can reduce the catabolic diversity of soil microbial communities. *Soil Biol Biochem* 32: 189-196.
- Duineveld BM, Rosado AS, van Elsas JD, van Veen JA. 1998. Analysis of the dynamics of bacterial communities in the rhizosphere of the chrysanthemum via denaturing gradient gel electrophoresis and substrate utilization patterns. *Appl Environ Microbiol* 64: 4950-4957.
- Duineveld BM, Kowalchuk GA, Keijzer A, van Elsas JD, van Veen JA. 2001. Analysis of bacterial communities in the rhizosphere of chrysanthemum via denaturing gradient gel electrophoresis of PCR-amplified 16S rRNA as well as DNA fragments coding for 16S rRNA. *Appl Environ Microbiol* 67: 172-178.
- Gelsomino A, Keijzer-Wolters AC, Cacco G, Van Elsas JD. 1999. Assessment of bacterial community structure in soil by polymerase chain reaction and denaturing gradient gel electrophoresis. *J Microbiol Methods* 38: 1-15.
- Grant, SB, Sanders BF, Boehm AB, Redman JA, Kim JH, Mrse RD, Chu AK, Gouldin M, McGee CD, Gardiner NA, Jones BH, Svejkovsky J, Leipzig GV. 2001. Generation of enterococci bacteria in a coastal saltwater marsh and impact on surf zone water quality. *Environ Sci Technol* 35: 2407-2416.
- Gray JP, Herwing RP. 1996. Phylogenetic analysis of the bacterial communities in marine sediments. *Appl Environ Microbiol* 62: 4049-4059.
- Gray TRG, Parkinson D (eds). 1968. *The Ecology of Soil Bacteria* 681. University of Toronto Press, Toronto, Canada.
- Grayston SJ, Vaughan D, Johnes D. 1996. Rhizosphere carbon flow in tree, in comparison with annual plants: the importance of root exudation and its impact on microbial activity and nutrient availability. *Appl Soil Ecol* 5: 29-56.
- Grayston SJ, Want SQ, Campbell CD, Edwards AC. 1998. Selective influence of plant species on microbial diversity in the rhizosphere. *Soil Biol Biochem* 30: 369-378.
- Holding AJ, Collee JG. 1971. Routine biochemical tests. In: *Methods in Microbiology* (Norris JR, Ribbons DW, eds) Vol 6A. Academic Press Inc Ltd, London and New York, pp 1-32.
- Howes BL, Weiskel PK, Geohringer DD, Teal JM. 1996. Interception of freshwater and nitrogen transport from uplands to coastal waters: the role of salt marshes. In: *Estuarine shores: evolution, environments and human alteration* (Nordstrom KF, Roman CT, eds). John Wiley & Sons, New York, N.Y. pp 287-310.
- Kim BS, Oh H-M, Kang H, Pack SS, Chun J. 2004. Remarkable bacterial diversity in the tidal flat sediment as revealed by 16S rDNA analysis. *J Microbiol Biotechnol* 14: 205-211.
- Kim BS, Oh H-M, Kang H, Chun J. 2005. Archaeal diversity in the tidal flat sediment as revealed by 16S rDNA analysis. *J Microbiol* 43: 144-151.
- Kim SJ, Lee GH. 1992. Distribution of heterotrophic bacteria and extracellular enzyme activities in the sediment of South Sea, Korea.

- Kor J Microbiol 30: 383-390.
- Latour X, Corberand TS, Laguerre G, Allard F, Lemanceau P. 1996. The composition of fluorescent pseudomonad populations associated with roots is influenced by plant and soil type. Appl Environ Microbiol 62: 2449-2456.
- Lee GH, Choi GG, Back CB. 1996. Distribution of aerobic/anaerobic saprophytic bacteria in the sediments of the Yellow Sea near Kunsan, Korea. Arch Hydrobiol Spec Issues Advanc Limnol 48: 227-232.
- Lee MS, Hong SG, Lee DH, Kim CK, Bae KS. 2001. Bacterial diversity in the mud flat of Suncheon bay, Chunnam Province, by 16S rRNA gene analysis. Kor J Microbiol 37: 137-144.
- Maloney PE, Vanbruggen AHC, Hu S. 1997. Bacterial community structure in relation to the carbon environments in lettuce and tomato rhizospheres and in bulk soils. Microb Eco 34: 109-117.
- Marshner P, Yang CH, Lieberei R, Crowley DE. 2001. Soil and plant specific effects on bacterial community composition in the rhizosphere. Soil Biol Biochem 33: 1437-1445.
- Metting BF. 1993. Soil Microbial ecology. Marcel Dekker, New York.
- Neal JL, Atkinson TG, Larson RI. 1970. Changes in the rhizosphere microflora of spring wheat induced by disomic substitution of a chromosome. Can J Microbiol 16: 153-158.
- Neal JL, Larson RI, Atkinson TG. 1970. Changes in rhizosphere populations of selected groups of physiological groups of bacteria related to substitution of specific pairs of chromosomes in spring wheat. Plant Soil 39: 209-212.
- Park SH. 2004. Taxonomical study of heterotrophic bacteria in the rhizosphere of halophytes in western and southern coastal area of Korea. 35. BS thesis. Graduate School, Kunsan National University, Korea.
- Park SH, Lee GH. 2006. Bacterial diversity in the rhizosphere of halophyte *Suaeda japonica* in western and Southern mudflats of Korea. J Ecol Biol 29: 399-404.
- Paul EA, Clark FE. 1988. Soil microbiology and biochemistry. Academic Press, San Diego.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstruction phylogenetic trees. Mol Biol Evol 4: 406-425.
- Singh G, Mukerji KG. 2006. Root exudates as determinant of rhizospheric microbial biodiversity. In: Microbial Activity in the Rhizosphere (Mukerji KG, Manoharachary C, Singh J, eds). Springer-Verlag, Berlin, Heidelberg, pp 39-53.
- Smith SV, Hollibaugh JT. 1993. Coastal metabolism and the oceanic organic carbon balance. Rev Geophys 31: 75-89.
- Teal JM, Howes BL. 2000. Salt marsh values; retrospection from the end of the century. In: Concepts and Controversies in Tidal Marsh Ecology (Weinstain MP, Kreeger DA eds). Kluwer Academic Publishers, Dordrecht, The Netherlands pp 3-7.
- Woldendorp JW. 1978. The rhizosphere as part of the plant-soil system. In Structure and Functioning of Plant Population. Verhandelingen der Koninklijke, Nederlandse Akademie van Wetenschappen, Afdeling Natuukunde, Twede Reeks, deel 70.
- Wollum AG. 1982. Cultural methods for soil microorganisms. In Method of Soil Analysis, Part 2: Chemical and Microbiological Properties, 2<sup>nd</sup> Ed, American Society of Agronomy, Inc., Soil Science Society of America, Inc. ed, Madison, Wisconsin pp 781-802.

(Received March 28, 2008; Accepted April 21, 2008)