



Biological assessment of streams and rivers in U.S. - design, methods, and analysis

Brenda Rashleigh^{1,*}, Steve Paulson², Joe Flotemersch³, Peg Pelletier¹

¹U.S. Environmental Protection Agency, Narragansett, Rhode Island, USA

²U.S. Environmental Protection Agency, Corvallis, Oregon, USA

³U.S. Environmental Protection Agency, Cincinnati, Ohio, USA

Abstract

Bioassessment is the use of biosurvey data, most commonly for benthic macroinvertebrates and fish, to obtain information about the health of waters in a region. In rivers, bioassessment results are used to evaluate biological condition and trends, to establish relationships between stressors and impairments, and to guide and evaluate management actions.

Keywords: bioassessment, monitoring, macroinvertebrates, fish, multimetric index

INTRODUCTION

Bioassessment is the use of biosurvey data to obtain information about the health of specific bodies of water. Biosurvey may measure the presence, condition, numbers, and types of fish, benthic macroinvertebrates, amphibians, algae, and plants. Biological endpoints provide the advantages of being easy to measure, responsive to stress in the watershed over time; and often ecologically important. Bioassessment is most commonly conducted in rivers and streams; it can also be applied in lakes, wetlands, and estuaries. There are several applications of the results of bioassessment: to evaluate the biological status, condition, and trends for a water body; to distinguish among potential stressors; to establish credible relationships between stressors and impairments; and to guide efforts and evaluate the effectiveness of management actions, including protection and restoration (Barbour et al. 1999). Bioassessment data are most effective when integrated with other measures of stream and river assessment, such as physical habitat, water chemistry, and landscape information (Angradi et al. 2009). We briefly re-

view bioassessment methods, analysis, and some recent developments in both areas.

BIOASSESSMENT METHODS

In the U.S., monitoring is typically conducted by federal and state governmental organizations, as well as non-governmental organizations (NGOs). A set of standardized methods has evolved at the federal level. National-level methods are provided by the U.S. Geological Survey (Moulton et al. 2002), and by the U.S. Environmental Protection Agency (EPA) (Barbour et al. 1999, U.S. EPA 2007). Most U.S. states also collect these data, and have established protocols that are similar to the national protocols. Most common methods for collecting algae include artificial substrates and hand collection (scraping of rocks). Most common methods for macroinvertebrates include collection nets (Slack sampler, kick net) and "grab" samplers. Most common methods for fish include

Open Access <http://dx.doi.org/10.5141/ecoenv.2013.010>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received 04 March 2013, Accepted 17 March 2013

***Corresponding Author**

E-mail: Rashleigh.Brenda@epa.gov

Tel: + 401-782-3014

www.kci.go.kr

electrofishing or seining. Additional methods have been developed for large rivers (Flotermersch et al. 2006). Seasonality is a consideration for methods development: the life cycles of algae and benthic macroinvertebrates, and the migration of fishes, can influence their distributions. Additional considerations include the sampling of representative habitat, rigorous sample tracking and preservation of samples for future identification, and preservation of archived samples of fish (Barbour et al. 1999).

Monitoring is conducted based on a sampling framework that describes the spatial distribution of sampling locations. In order to generate an unbiased, representative assessment, a randomized design is often used. For large regions, stratification may be used to generate sub-regions that are more homogeneous. For regional scale frameworks, Frimpong and Angermeier (2010) found that the best approach for stratification by zoogeographic and physiographic units. A challenge in the development of national-level assessment is that some regions are more degraded than others. Herlihy et al. (2008) suggest that region-specific models be developed, which addresses this issue, where reference site quality and assessment thresholds are set by region. However, it makes the comparison across regions more difficult. This could be addressed by assessing these reference conditions against one another. A recent review of monitoring design (Strobl and Robillard 2008) notes that monitoring network designs should consider input from policy and public sectors, as well as scientists and managers, and should be periodically reassessed.

Two future directions in bioassessment include recognizing the role of the stream network in influencing assemblages, and incorporating genetic and molecular techniques into bioassessment. Recent work (Hitt and Angermeier 2011) recognizes the importance of stream network position to bioassessment, where dispersal among streams influences fish community composition over small spatial scales (10 km), and may influence certain metrics often used in bioassessment programs. The development of molecular and genetic techniques is a future direction that is showing promise for early detection of stressor effects on individuals. A recent study by Pilgrim et al. (2011) hypothesized that DNA barcoding could provide greater discriminatory ability than current genus-level identifications of species, which could lead to more specific and sensitive assessments of water bodies, however, their analysis demonstrated limited advantages of this technique.

ANALYSIS

Most water resource agencies in the U.S. use a multi-metric approach for the analysis of bioassessment data (e.g., Karr and Chu 1999). A multimetric approach involves two steps: 1) selection and calibration of metrics, which can be combined into a multimetric index, and 2) the assessment and judgement of impairment. The selection of metrics is a well-documented activity (e.g., Barbour et al. 1999, Whittier et al. 2007). Metrics should be tested for performance across the full human disturbance gradient in a region, in order to improve statistical confidence, and to reduce bias (Diamond et al. 2012). A multivariate approach has also been used to analyze bioassessment data. Multivariate statistical analysis of biosurvey data can be used to assess stressor impairment (EPA 2000) and explore patterns in aggregations of taxa or taxon traits (Qian et al. 2012). A popular technique is the generation of the Observed/Expected (O/E) ratio as an indicator for each site – the Observed taxa richness (O) is obtained from the field sampling at a site, and the Expected taxa richness (E) is calculated as the sum of statistically generated probabilities of occurrence for each taxa (Hawkins 2012). Using this approach, a reference site is expected to have an O/E ratio of 1.0; a site impaired by stressors will have a value <1. A similar multivariate approach is used by water resource agencies in Europe and Australia (Wright 2000). Another direction is the use of a Biological Condition Gradient approach, which allows the assessment of directional trends, summary of different methods to a common scale, and easier communication to bioassessment results to non-scientific audiences (Davies and Jackson 2006).

Results from the bioassessment can be summarized at regional and national levels. For example, Paulson et al. (2008) reported that for the first U.S. Wadeable Streams Assessment, data collected at 1392 randomly selected sites yielded the result that 42% of the length of US streams was in poor condition compared to best available reference sites, 25% was in fair condition, and 28% was in good condition. A current challenge is the comparability of different bioassessment programs, Cao and Hawkins (2011) provided several recommendations for improving how different bioassessment programs can be integrated. Hughes and Peck (2008) describe the challenges of reaching a compromise between scientific rigor, consistent and practical implementation over large regions and many participants, and the realities of time and money; they emphasize the importance of advance planning, standardized methods, and cooperative research.

ACKNOWLEDGEMENTS

This is ORD tracking number ORD-003923 of the Atlantic Ecology Division, National Health and Environmental Effects Research Laboratory, Office of Research and Development, U.S. Environmental Protection Agency. Although the research described in this article has been funded by the U.S. Environmental Protection Agency, it has not been subjected to Agency review. Therefore, it does not necessarily reflect the views of the Agency.

LITERATURE CITED

- Angradi TR, Bolgrien DW, Jicha TM, Pearson MS, B.H. Hill, Taylor DL, Schweiger EW, Shepard L, Batterman AR and Moffett MF et al. 2009. A bioassessment approach for mid-continent great rivers: the Upper Mississippi, Missouri, and Ohio (USA). *Environmental Monitoring and Assessment* 152: 425-442.
- Barbour MT, Gerritsen J, Snyder BD, and Stribling JB. 1999. *Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish, Second Edition*. EPA 841-B-99-002. U.S. Environmental Protection Agency; Office of Water; Washington, D.C.
- Cao Y. and Hawkins CP. 2011. The comparability of bioassessments: a review of conceptual and methodological issues. *Journal of the North American Benthological Society* 30(3):680-701.
- Davies SP, and Jackson SK. 2006. The biological condition gradient: a descriptive model for interpreting change in aquatic ecosystems. *Ecological Applications* 16:1251-1266.
- Diamond J, Stribling JR, Huff L, Gilliam J. 2012. An approach for determining bioassessment performance and comparability *Environ Monit Assess* 184:2247–2260.
- Flotemersch JE, Stribling JB, and Paul MJ. 2006. *Concepts and Approaches for the Bioassessment of Non-wadeable Streams and Rivers*. EPA 600-R-06-127. US Environmental Protection Agency, Cincinnati, Ohio.
- Frimpong EA and Angermeier PL. 2010. Comparative Utility of Selected Frameworks for Regionalizing Fish-Based Bioassessments across the United States, *Transactions of the American Fisheries Society* 139(6):1872-1895
- Hawkins CP. 2012. Use of RIVPACS, a multi-taxon niche model, to assess the condition of macroinvertebrate assemblages in aquatic ecosystems. *Korean Journal of Limnology* (in press).
- Herlihy AT, Paulsen SG, Van Sickle J, Stoddard JL, Hawkins CP and Yuan LL. 2008. Striving for consistency in a national assessment: the challenges of applying a reference-condition approach at a continental scale. *Journal of the North American Benthological Society* 27(4):860-877.
- Hitt NP and Angermeier PL. 2011. Fish community and bioassessment responses to stream network position. *Journal of the North American Benthological Society* 30(1):296-309.
- Hughes RM and Peck DV. 2008. Acquiring data for large aquatic resource surveys: the art of compromise among science, logistics, and reality. *Journal of the North American Benthological Society* 27:837–859.
- Karr JR, and Chu EW. 1999. *Restoring life in running waters: better biological monitoring*. Island Press, Washington, DC.
- Moulton II, SR, Kennen JG, Goldstein RM, and Hambrook JA. 2002. *Revised Protocols for Sampling Algal, Invertebrate, and Fish Communities as Part of the National Water-Quality Assessment Program*. U.S. Geological Survey, Open-File Report 02-150, Reston, VA. <http://pubs.usgs.gov/of/2002/ofr-02-150/pdf/ofr02-150.pdf>
- Paulsen SG, Mayo A, Peck DV, Stoddard JL, Tarquinio E, Holdsworth SM, Van Sickle J, Yuan LL, Hawkins CP, Herlihy AT, Kaufmann PR, Barbour MT, Larsen DP and Olsen AR. 2008. Condition of stream ecosystems in the US: an overview of the first national assessment. *Journal of the North American Benthological Society* 27:812-821
- Pilgrim EM, Jackson SA, Swenson S, Turcsanyi I, Friedman E, Weigt L, and Bagley MJ. 2011. Incorporation of DNA barcoding into a large-scale biomonitoring program: opportunities and pitfalls. *Journal of the North American Benthological Society*. 30(1):217-231.
- Qian SS, Cuffney TF and McMahon G. 2012. Multinomial regression for analyzing macroinvertebrate assemblage composition data. *Freshwater Science* 31(3):681-694.
- Strobl RO and Robillard PD. 2008. Network design for water quality monitoring of surface freshwaters: A review. *Journal of Environmental Management* 87:639-648.
- U.S. EPA. 2000. *Stressor Identification Guidance Document*. U.S. Environmental Protection Agency. Office of Water, Washington, EPA-822-B-00-025
- U.S. EPA. 2007. *National Rivers and Streams Assessment: Field Operations Manual*. EPA-841-B-07-009. U.S. Environmental Protection Agency, Washington, DC.
- Whittier TR, Hughes RM, Stoddard JL, Lomnick GA, Peck DV and Herlihy AT. 2007. A structured approach to developing indices of biotic integrity: three examples from western USA streams and rivers. *Transactions of the American Fisheries Society* 136:718–735.

Wright JF. 2000. An introduction to RIVPACS. Pages 1–24 in J. F. Wright, D. W. Sutcliffe, and M. T. Furse (editors). Assessing the biological quality of fresh waters: RIVPACS and other techniques. Freshwater Biological Association, Ambleside, UK.