



Regional land cover patterns, changes and potential relationships with scaled quail (*Callipepla squamata*) abundance

Paikho Rho^{1,*}, X. Ben Wu², Fred E. Smeins², Nova J. Silvy³ and Markus J. Peterson³

¹Department of Environmental Planning, Keimyung University, Daegu 704-701, Korea

²Department of Ecosystem Science and Management, Texas A&M University, College Station, TX 77843-2138, USA

³Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, TX 77843-2258, USA

Abstract

A dramatic decline in the abundance of the scaled quail (*Callipepla squamata*) has been observed across most of its geographic range. In order to evaluate the influence of land cover patterns and their changes on scaled quail abundance, we examined landscape patterns and their changes from the 1970s to the 1990s in two large ecoregions with contrasting population trends: (1) the Rolling Plains ecoregion with a significantly decreased scaled quail population and (2) the South Texas Plains ecoregion with a relatively stable scaled quail population. The National Land Cover Database (NLCD) and the U.S. Geological Survey's (USGS) Land Use/Land Cover data were used to quantify landscape patterns and their changes based on 80 randomly located 20×20 km² windows in each of the ecoregions. We found that landscapes in the Rolling Plains and the South Texas Plains were considerably different in composition and spatial characteristics related to scaled quail habitats. The landscapes in the South Texas Plains had significantly more shrubland and less grassland-herbaceous rangeland; and except for shrublands, they were more fragmented, with greater interspersions among land cover classes. Correlation analysis between the landscape metrics and the quail-abundance-survey data showed that shrublands appeared to be more important for scaled quail in the South Texas Plains, while grassland-herbaceous rangelands and pasture-croplands were essential to scaled quail habitats in the Rolling Plains. The decrease in the amount of grassland-herbaceous rangeland and spatial aggregation of pasture-croplands has likely contributed to the population decline of scaled quails in the Rolling Plains ecoregion.

Key words: *Callipepla squamata*, landscape change, landscape pattern metrics, Rolling Plains, scaled quail, South Texas Plains

INTRODUCTION

The scaled quail (*Callipepla squamata*) is an upland game bird species whose population has declined 3.1% per year according to the North American Breeding Bird Survey 1966-2011 (Sauer et al. 2013). Although the overall population of scaled quails declined, population levels of scaled quails remained steady in some areas. Studies

on the population trends of scaled quails in several major ecological regions showed that in the Rolling Plains ecoregion, as in many other regions, scaled quail populations had decreased to a very low level with an average decline rate of 9.2% per year from 1978 to 2000; the rate of decline in the South Texas Plains, however, was only 0.2% per year

<http://dx.doi.org/10.5141/ecoenv.2015.020>



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received 9 March 2015, Accepted 3 April 2015

*Corresponding Author

E-mail: wildlife@kmu.ac.kr

Tel: +82-53-580-5917

www.kci.go.kr




Fig. 1. Location of 80 randomly located windows for the Rolling Plains and the South Texas Plains. ArcView script was used to extract random windows of $20 \times 20 \text{ km}^2$. Landscape metrics and scaled quail abundance were calculated based on these random windows.

(Peterson 2001, Bridges et al. 2002, Rho 2003, Silvy et al. 2007).

It is unclear why the population of scaled quails is relatively steady in the South Texas Plains, but declining rapidly in the Rolling Plains. Many factors, including rangeland management practices such as brush control (Rollins 2000), the Conservation Reserve Program (CRP; Schemnitz 1993), precipitation patterns (Bridges et al. 2002), and disease (Rollins 2000) have been suggested to cause discrepancies in the decline of scaled quail populations in the two ecoregions. Rollins (2000) found that the number of scaled quails had periodically increased and decreased in the Rolling Plains. Changes in landscape patterns of scaled quail habitats were also suspected to be a major factor causing the long-term population decline in areas such as the Rolling Plains ecoregion (Bridges et al. 2002). Although many studies have addressed the effects of landscape patterns and their changes on the abundance of species, landscape characteristics and scaled quail abundance has received little critical attention. Therefore, for the conservation of scaled quails, it is important to examine patterns of landscapes and their changes in regions with contrasting population trends. It is also important to explore how changes in a specific landscape relate to changes in scaled quail populations.

The objective of this study is to examine landscape patterns of the Rolling Plains and the South Texas Plains ecoregions, and landscape pattern changes that had occurred between the 1970s and the 1990s, and the possible relationship with the contrasting trends of the scaled

quail populations in these ecoregions. Scaled quails in the Rolling Plains and South Texas Plains are assumed to be different subspecies: the Arizona scaled quail (*C. s. pallida*) and the chestnut-bellied scaled quail (*C. s. castanogastris*) (Wallmo 1957, Silvy et al. 2007). They are known to use habitats differently (Guthery et al. 2001). But few studies have been conducted to quantify landscape structures related to wildlife habitats in the subspecies. Geographical Information Systems (GIS) and landscape analysis approaches were used in this study to determine (1) how landscape patterns differed in the Rolling Plains and the South Texas Plains, (2) how changes in these landscape patterns from the 1970s to the 1990s have affected the population of scaled quails, and (3) which components of landscape patterns and their changes likely contributed to the different habitat qualities of scaled quail subspecies in the Rolling Plains and the South Texas Plains.

MATERIALS AND METHODS

Study areas

The Rolling Plains and South Texas Plains ecoregions were selected as study areas (Fig. 1). The Rolling Plains, composed of approximately 9,700,000 ha of alternating shrublands and native prairies, has a gently rolling to moderately rough topography that ranges from 245-915 m in elevation. Annual rainfall ranges from about 55 cm in the west to 75 cm in the eastern region, and peaks in May and September. Typically, there is a dry summer period, with high temperatures and high rates of evaporation. Soils vary from coarse sands along outwash terraces adjacent to streams, to tight or compact clays or red-bed clays and shales (Wu et al. 2002).

The South Texas Plains are composed of about 8,000,000 ha of subtropical brushland with small trees, shrubs, cacti, forbs, and grasses, with level to rolling topography and elevation range of 305 m above sea level. The average annual precipitation range is 40-90 cm, and this increases from west to east with largest amounts of rainfall in May and September. Summer temperatures are high, with extremely high evaporation rates in the west. Periodic droughts are common in this ecoregion. Soils range from clays to sandy loams, and vary in chemical properties from calcareous to slightly acidic (Wu et al. 2002).

Spatial data development and analysis

The National Land Cover Database (NLCD, Riitters et

al. 2000) was acquired and used to quantify landscape patterns of the Rolling Plains and the South Texas Plains ecoregions in the 1990s. This fine (30 m) resolution data was developed based on Landsat 5 Thematic Mapper (TM) satellite imagery from the early 1990s by the Multi-Resolution Land Characteristics Consortium (MRLC), which is made up of multiple federal agencies. Quantification of landscape patterns for an entire ecoregion would be unfeasible with conventional computing power. Therefore, analyses based on randomly located landscapes (subsets of the ecoregion) with smaller spatial extent can be effective, and may even be necessary as to quantify the landscape pattern and statistical variation of each ecoregion. ArcView GIS Spatial Analyst (Environmental Systems and Research Institute, Inc., Redlands, CA, USA) was used to generate 160 random windows (80 for each ecoregion) of 40,000 ha ($20 \times 20 \text{ km}^2$) each in order to compare the landscape patterns between the Rolling Plains and the South Texas Plains (Fig. 1).

A wide variety of landscape metrics are available for evaluating spatial patterns of individual patches and patch types such as land-cover classes, and whole landscapes (Uuemaa et al. 2009). For this study, a set of metrics related to scaled quail habitats, including patch density (PD), mean patch size (MPS), edge density (ED), mean shape index (MSI), interspersed-juxtaposition index (IJI), and Shannon's evenness index (SEI), were used to quantify the pattern of forests, shrublands, grassland-herbaceous rangelands, and pasture-cropland cover types. PD expresses the number of patches within the entire landscape of each random window, and MPS is the average size of patches in the landscape. In order to quantify landscape heterogeneity, ED, the total edge length calculated per unit area, and MSI, the average perimeter-to-area ratio of all patches in the landscape are measured (McGarigal and Marks 1995). The value of IJI, a measure of the spatial configuration of patch types, increased when the patches were more evenly interspersed in a "salt and pepper" mixture. SEI quantifies evenness among landscapes. It is equal to zero when the observed patch distribution is low and approaches one when the distribution of patch types becomes more even at the landscape level (McGarigal and Marks 1995). These landscape metrics were selected based on the fact that scaled quails inhabit a variety of land-cover types throughout their life cycle; require interspersed habitats for breeding, escaping, food, and loafing; and more quails observed near the edges between different cover types (Schemnitz 1961, Silvy et

al. 2007). The Patch Analyst extension for ArcView (Elkie et al. 1999), based on the FRAGSTATS software (McGarigal and Marks 1995), was used to calculate the landscape metrics.

The USGS Land Use/Land Cover data (LULC; USGS 1990) was obtained and used to quantify landscape patterns for the 1970s. It was then compared to the NLCD of the 1990s to assess the regional landscape changes in the Rolling Plains and the South Texas Plains. Although these two datasets are standard, regional-scale land-cover data widely used in large-scale studies, they have considerable differences in source materials, classification schemes, and spatial resolution. GIS modeling approaches were used to standardize the two datasets in order to make comparisons between them as valid as possible. Raster maps of LULC with different spatial resolutions (30 m, 50 m, 100 m, and 300 m) were first generated from the vector data. Landscape metrics for each of these raster maps were calculated and compared to the landscape metrics of the original vector map in order to find the appropriate spatial resolution for resampling. Based on this analysis, 100 m was determined as an appropriate resolution for vector-to-raster conversion of the LULC data. The 30 m resolution NLCD data was converted into 90-m resolution data with a 3×3 majority filter, and then resampled to 100-m resolution data in order to compare to the rasterized LULC data. The resampled NLCD data was further processed by removing all patches smaller than the minimum map unit for LULC, and using the RegionGroup and Nibble function to better match with the LULC data (Environmental Systems Research Institute 1998).

Correlation analyses were conducted to examine the relationship between landscape pattern metrics and scaled quail abundance. Species abundance was calculated based on the Texas Parks and Wildlife Department (TPWD) quail abundance survey data (Wilson 1992). Due to the nature of scaled quail population dynamics, which is characterized by substantial annual fluctuations, expected levels of scaled quail abundance developed using route regression (Geissler and Sauer 1990) over the 1978-2000 period were used in the analysis. A raster map of species abundance, which represents the number of scaled quails seen in each 1.6-km transect segment, was developed by interpolation. The average value of the interpolated raster map for each random window was calculated and used to correlate scaled quail abundance to landscape pattern metrics of 1978 and 1992.




Fig. 2. Land cover classified into forest, shrub, herbaceous, and pasture-cropland types from 1970s to 1990s of the Rolling Plains and South Texas Plains.

RESULTS

Landscape patterns: Rolling Plains compared with South Texas Plains

Landscape composition, which is the proportion of total land area occupied by each land cover type, was significantly different between the Rolling Plains and the South Texas Plains (Fig. 2). Shrubland cover occupied the largest area in the South Texas Plains followed by pasture-croplands and grassland-herbaceous rangelands. In contrast, grassland-herbaceous rangelands were the most abundant cover type in the Rolling Plains, followed by substantial amounts of shrublands and pasture-croplands (Fig. 3).

Contrary to the landscape composition, analysis based on whole landscapes showed that spatial patterns of landscape elements in the Rolling Plains and the South Texas Plains were similar even though the South Texas Plains had significantly higher PD and interspersion among land cover classes. Examination of the spatial patterns of individual land cover classes, however, revealed significant differences between the Rolling Plains and the South Texas Plains (Fig. 2). For example, pasture-cropland




Fig. 3. Percentage of land cover type for the Rolling Plains and South Texas Plains in the 1990s. Meaningful t-test results between the two ecoregions of a 5% level are marked with an *. Each bar represents the mean of four land cover types: LC1, forest; LC2, shrubland; LC3, grassland-herbaceous; and LC4, pasture-cropland. The error bar shows standard error of the mean.

patches in the South Texas Plains had twice the PD and about 5 times smaller than those in the Rolling Plains (Table 1). The South Texas Plains had a greater amount of pasture-cropland edges, and were more interspersed with other land cover classes.

Table 1. Mean \pm S.E. of landscape pattern metrics at landscape and class level for the Rolling Plains and the South Texas Plains in the 1990s

Level	Landscape metrics	Rolling Plains	South Texas Plains	P
Landscape level	PD, #/100ha	26.0 \pm 0.92	31.4 \pm 1.34	0.0012
	MPS, ha	4.2 \pm 0.15	3.8 \pm 0.25	0.1836
	ED, m/ha	119.9 \pm 3.48	124.1 \pm 4.56	0.4658
	MSI	1.31 \pm 0.00	1.31 \pm 0.00	0.5271
	IJI	48.3 \pm 2.01	64.6 \pm 1.34	< 0.0001
Forest class	SEI	0.64 \pm 0.01	0.63 \pm 0.02	0.5484
	PD, #/100ha	2.7 \pm 0.39	7.8 \pm 0.47	< 0.0001
	MPS, ha	0.6 \pm 0.12	1.0 \pm 0.10	0.0080
	ED, m/ha	12.4 \pm 2.32	39.3 \pm 3.25	< 0.0001
	MSI	1.18 \pm 0.01	1.27 \pm 0.01	< 0.0001
Shrubland class	IJI	61.7 \pm 2.63	40.1 \pm 2.58	< 0.0001
	PD, #/100ha	11.1 \pm 0.74	6.1 \pm 0.60	< 0.0001
	MPS, ha	4.0 \pm 0.54	50.1 \pm 18.65	0.0155
	ED, m/ha	91.5 \pm 3.68	97.6 \pm 3.04	0.2057
	MSI	1.35 \pm 0.01	1.35 \pm 0.01	0.5322
Grassland / herbaceous class	IJI	41.4 \pm 2.61	71.9 \pm 1.42	< 0.0001
	PD, #/100ha	8.0 \pm 0.48	10.0 \pm 0.50	0.0042
	MPS, ha	10.8 \pm 2.06	1.3 \pm 0.10	< 0.0001
	ED, m/ha	96.9 \pm 2.95	57.1 \pm 3.49	< 0.0001
	MSI	1.3 \pm 0.01	1.32 \pm 0.01	0.0800
Pasture / cropland class	IJI	46.0 \pm 1.78	52.7 \pm 2.00	0.0133
	PD, #/100ha	1.9 \pm 0.17	4.2 \pm 0.29	< 0.0001
	MPS, ha	26.2 \pm 5.08	5.5 \pm 0.99	0.0001
	ED, m/ha	27.5 \pm 1.76	40.1 \pm 3.52	0.0023
	MSI	1.38 \pm 0.01	1.3 \pm 0.01	< 0.0001
	IJI	56.9 \pm 1.59	71.1 \pm 1.14	< 0.0001

P values are for statistical differences between two ecoregions by t-test.

Landscape change: the 1970s compared with the 1990s

Despite the difference in landscape composition of the Rolling Plains and the South Texas Plains, trends concerning their changing landscapes from the 1970s to the 1990s were similar. Rho (2003) observed that shrubland cover increased in both the Rolling Plains (12.2% to 24.7%) and the South Texas Plains (51.1% to 59.7%). Grassland-herbaceous rangeland decreased in both the Rolling Plains (53.9% to 44.6%) and the South Texas Plains (15.2% to 9.9%). Pasture-cropland and forest patches decreased slightly in both ecoregions (Rho 2003).

At the landscape level, PD increased significantly both in the Rolling Plains (1.56 to 7.07/100 ha) and the South Texas Plains (1.91 to 8.28/100 ha) from the 1970s to the 1990s. ED also increased significantly in the Rolling Plains (11.26 to 25.61 m/ha) and the South Texas Plains (12.43 to 25.46 m/ha). However, MPS decreased in the Rolling Plains (1,404 to 160 ha) and the South Texas Plains (674 to 155ha). MSI increased slightly, but not significantly statistically, in the Rolling Plains (1.57 to 1.65) and the South Texas Plains (1.55 to 1.59). IJI decreased in the Rolling Plains (53.57 to 50.26), but increased in the South Texas Plains (50.93 to 54.67). SEI remained relatively stable for both ecoregions.

Landscape patterns of pasture-cropland patches had dramatically changed during the period from the 1970s to the 1990s when compared to other land cover classes. Pasture-cropland landscape in the South Texas Plains became more fragmented in the 1990s than that in the Rolling Plains. For example, pasture-cropland patches of the South Texas Plains were relatively few (0.40 compared to 0.58/100 ha) and large (981 compared to 901 ha) in the 1970s, but they transformed into many (1.56 compared to 1.13/100 ha) and small (201 compared to 506 ha) patches in the 1990s when compared to the Rolling Plains whereas IJI of pasture-cropland patches remained stable during this period and consistently higher in the South Texas Plains than that of the Rolling Plains.

Correlations between species abundance and landscape characteristics

Scaled quail abundance had significant negative correlations with landscape-level PD, ED, MSI, IJI, and SEI in the Rolling Plains. In the South Texas Plains, scaled quail abundance was negatively correlated with the mean shape index, but had no significant correlation with other landscape metrics (Table 2).

The percentage of cover and most of the spatial attributes of forests were significantly correlated with scaled quail abundance (Table 3). In the Rolling Plains, negative correlations were found between scaled quail abundance and percent cover, PD, ED, MPS, and MSI of forest patches. Negative correlations between species abundance and percentage of forest cover, average size of forest patch were statistically significant in the South Texas Plains, indicating that scaled quail number decreases as forest area and patch size increases in the ecoregion. A highly positive and statistically significant correlation was found between scaled quail abundance and shrubland cover in the South Texas Plains, but not in the Rolling Plains. There was a negative correlation between species abundance and percentage cover of grassland-herbaceous rangeland in the Rolling Plains, but a weak and insignificant correlation in the South Texas Plains.

DISCUSSION

Vegetation cover and scaled quail abundance

We observed different relationships between landscape patterns and their changes on scaled quail abundance in the Rolling Plains and South Texas Plains, which are consistent to results from previous studies. Guthery et al. (2001) noted that scaled quails in the South Texas Plains have different requirements for vegetation cover, compared to other regions. They described shrubland as a key habitat in the South Texas Plains, where scaled quails frequently use woody plants and clusters as escaping cover to avoid harsh weather and predators (Wilson and Crawford 1987), and scaled quail whistle counts positively correlated with shrubland cover (Reid et al. 1979). Our research revealed that the percentage of shrubland cover positively correlated with scaled quail abundance in the South Texas Plains, but not in the Rolling Plains (Table 3).

In the Rolling Plains, grassland-herbaceous rangeland was described as an essential habitat for scaled quails, as it provided loafing (Stormer 1981) and roosting cover (Stormer 1984). Decrease in grassland-herbaceous rangeland and increase of shrublands over the 1970s to the 1990s suggested that increase in woody cover in the Rolling Plains might be related to the decline of scaled quail population, which is contrary to the South Texas Plains, where scaled quail population remained stable even when the shrubland cover increased. Tall and dense areas of woody plants have long been recognized to be of little use to scaled quails, particularly in southeastern

Arizona and northwest Texas (Goodwin and Hungerford 1977, Stormer 1981), because densely vegetated cover impedes the bird's progress on the ground when foraging and escaping.

The reason for different impacts of landscape patterns and their changes on scaled quail abundance of these ecoregions is probably caused by the different scaled quail subspecies in the Rolling Plains and the South Texas Plains, respectively. Wallmo (1957) mentioned that southern Arizona and northwestern Texas had the same subspecies and southern Texas had other subspecies. Interestingly, habitat selection described by Guthery et al. (2001) coincided with the boundary of scaled quail subspecies. But landscape patterns and their changes rarely correlated with population dynamics of the subspecies in the range. Our research showed that the percentage of grassland-herbaceous cover negatively correlated with

the abundance of scaled quails in the Rolling Plains, but not in the South Texas Plains.

Percentage and spatial arrangement of pasture-cropland patches

Pasture-cropland might play a different role with respect to scaled quail habitats in the Rolling Plains compared to the South Texas Plains. It was suggested that croplands was an important habitat in the Rolling Plains ecoregion (Schemnitz 1961, Leyva-Espinosa 2000), but CRP that was initiated in 1985 might be one of the reasons causing population decline in the ecoregion because of expansion of homogeneous cropland patches in the Rolling Plains (Schemnitz 1993). Studies are rarely conducted to evaluate pasture-croplands impact on subspecies abundance in the South Texas Plains. Our results showed

Table 2. Correlations between landscape level metrics and scaled quail abundance for the Rolling Plains and the South Texas Plains in the 1990s

Landscape level metrics	Rolling Plains		South Texas Plains	
	<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
PD	-0.35	0.0121	-0.10	0.6144
MPS	0.26	0.0675	-0.06	0.7750
ED	-0.43	0.0021	-0.18	0.3728
MSI	-0.43	0.0020	-0.45	0.0204
IJI	-0.29	0.0413	-0.25	0.2157
SEI	-0.42	0.0012	-0.28	0.1603

Table 3. Correlation between class level metrics and scaled quail abundance for the Rolling Plains and the South Texas Plains in the 1990s

Cover class	Class level metrics	Rolling Plains		South Texas Plains	
		<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
Forest	% Cover	-0.33	0.0245	-0.44	0.0250
	PD	-0.40	0.0052	-0.15	0.4560
	MPS	-0.31	0.0280	-0.49	0.0117
	ED	-0.38	0.0085	-0.37	0.0642
	MSI	-0.42	0.0030	-0.42	0.0332
Shrubland	% Cover	0.04	0.7599	0.45	0.0209
	PD	-0.28	0.0509	-0.33	0.0947
	MPS	0.34	0.0160	-0.02	0.9393
	ED	-0.35	0.0116	0.01	0.9512
	MSI	-0.09	0.5422	-0.14	0.4850
Grassland / herbaceous	% Cover	-0.25	0.0833	0.01	0.9664
	PD	0.07	0.6084	0.29	0.1528
	MPS	-0.10	0.5084	-0.05	0.8088
	ED	-0.31	0.0280	0.13	0.5421
	MSI	-0.22	0.1324	0.02	0.9233
Pasture / cropland	% Cover	0.25	0.0813	-0.33	0.1050
	PD	-0.42	0.0024	-0.12	0.5749
	MPS	0.26	0.0724	-0.23	0.2599
	ED	-0.13	0.3767	-0.29	0.1514
	MSI	-0.08	0.5694	-0.36	0.0659

there were greater numbers of smaller pasture-cropland patches with more edges and greater interspersions with other cover classes in the Rolling Plains than in the South Texas Plains in the 1970s. These spatial patterns were suitable for scaled quails in the Rolling Plains, since agricultural grains and man-made features (e.g., machinery, fences, brush piles) near or within the cropland patches were frequently used as winter food and loafing-escaping sites by scaled quails (Schemnitz 1961, Silvy et al. 2007). However, from the 1970s to the 1990s, pasture-cropland cover became aggregated (with fewer but larger patches) in the Rolling Plains, and fragmented in the South Texas Plains. These changes in pasture-croplands suggest that declining scaled quail abundance was possibly associated with the decrease and increase in spatial aggregation of pasture-croplands in the Rolling Plains.

Uncertainty of landscape comparison between NLCD and LULC data

Although the NLCD and LULC data was adjusted to make them comparable, the degree of reliability of the landscape comparisons between the 1970s and the 1990s was uncertain. Landscape comparisons based on the assessment of relative differences between the two ecoregions from the 1970s to 1990s, however, should be reliable and can offer insights on differential landscape dynamics. Our study on absolute values of landscape metrics between 1970s and 1990s indicated fragmentation processes in the Rolling Plains and the South Texas Plains landscapes. Changes in landscapes of both the Rolling Plains and the South Texas Plains were different during that period as is shown by pasture-cropland class metrics. For example, pasture-cropland patches in the South Texas Plains were relatively few and large in the 1970s, but they transformed into many small patches in the 1990s, compared to the Rolling Plains. The decrease of pasture-cropland and more aggregated configuration in the Rolling Plains are possibly linked to the long-term population decline of scaled quail in the ecoregion.

The relative differences in landscape pattern metrics from the 1970s to the 1990s remained unchanged in the shrubland and grassland-herbaceous between the Rolling Plains and the South Texas Plains. These results suggest that landscape patterns of vegetation cover remained temporally consistent between the 1970s and 1990s. For example, the number of grassland-herbaceous patches in the Rolling Plains was higher than in South Texas Plains in the 1970s, and continued to grow in the 1990s. Compared to the subspecies in the South Texas Plains, scaled quails

in the Rolling Plains were influenced by the decrease in grassland-herbaceous land cover and the aggregation of cropland-pasture landscape. Habitat management of scaled quail ecology and life cycles is closely related to the CRP and anthropogenic factors affecting the percentage cover of grassland-herbaceous and cropland-pastures.

ACKNOWLEDGMENTS

This work was supported by a grant from the Cross Timbers Chapter of Quail Unlimited, Texas Parks and Wildlife Department, Texas AgriLife Research, and a Tom Slick Graduate Fellowship to Paikho Rho.

LITERATURE CITED

- Bridges AS, Peterson MJ, Silvy NJ, Smeins FE, Wu XB. 2002. Landscape-scale land-cover change and long-term abundance of scaled quail and northern bobwhite in Texas. *Proc Natl Quail Symp* 5: 161-167.
- Elkie PC, Rempel RS, Carr A. 1999. *Patch Analyst User's Manual: A Tool for Quantifying Landscape Structure*. Ontario Ministry of Natural Resources, Boreal Science, Northwest Science & Technology, Thunder Bay.
- Environmental Systems Research Institute. 1998. *Working with ArcView Spatial Analyst*. Environmental Systems Research Institute (ESRI) Inc., Redlands, CA.
- Geissler PH, Sauer JR. 1990. Topics in route-regression analysis. In: *Survey Designs and Statistical Methods for the Estimation of Avian Population Trends* (Sauer JR, S Droege, eds). U.S. Fish and Wildlife Service. Washington, D.C., pp 54-57.
- Goodwin JG Jr, Hungerford CR. 1977. Habitat used by native Gambel's and scaled quail and released masked bobwhite quail in southern Arizona. U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Guthery FS, King NM, Kuvlesky WP Jr, DeStefano S, Gall SA, Silvy NJ. 2001. Comparative habitat use by three quails in desert grassland. *J Wildl Manage* 65: 850-860.
- Leyva-Espinosa IR. 2000. Use of broad-scale data to assess changes of scaled quail population in Texas. PhD Dissertation. Texas Tech University, Lubbock, TX, USA.
- McGarigal K, Marks BJ. 1995. *FRAGSTATS: Spatial Pattern Analysis Program for Quantifying Landscape Structure*. United States Department of Agriculture Forest Service, Pacific Northwest Research Station, Portland, OR.
- Peterson MJ. 2001. Northern bobwhite and scaled quail

- abundance and hunting regulation: a Texas example. *J Wildl Manage* 65: 828-837.
- Reid RR, Grue CE, Silvy NJ. 1979. Competition between bobwhite and scaled quail for habitat in Texas. *Proc Annu Conf Southeastern Fish Wildl Agencies* 33: 146-153.
- Rho P. 2003. GIS-based multiple-scaled study on scaled quail. PhD Dissertation. Texas A&M University, College Station, TX, USA.
- Riitters KH, Wickham JD, Vogelmann JE, Jones KB. 2000. National land-cover pattern data. *Ecology* 81: 604-604.
- Rollins D. 2000. Status, ecology and management of scaled quail in West Texas. In: *Quail IV* (Brennan LA, Palmer WE, Burger LW Jr, Pruden TL, eds). Proceedings of the Fourth National Quail Symposium. Tall Timbers Research Station, Tallahassee, FL, pp165-172.
- Sauer JR, Link WA, Fallon JE, Pardieck KL, Ziolkowski DJ Jr. 2013. The North American breeding bird survey 1966-2011: summary analysis and species accounts. *North American Fauna* 79: 1-32.
- Schemnitz SD. 1961. Ecology of the scaled quail in the Oklahoma Panhandle. *Wildl Monogr* 8: 3-47.
- Schemnitz SD. 1993. Scaled quail habitats revisited – Oklahoma panhandle. In: *Quail III* (Church KE, Dailey TV, eds). Proceedings of the 3rd National Quail Symposium. Kansas Department of Wildlife and Parks, Pratt, KS, pp143-147.
- Silvy NJ, Rollins D, Whisenant SW. 2007. Scaled quail ecology and life history In: *Texas Quails: Ecology and Management* (Brennan LA, ed). Texas A&M University Press, College Station, TX, pp 65-88.
- Stormer FA. 1981. Characteristics of Scaled Quail Loafing Coverts in Northwest Texas. U.S. Department of Agriculture Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Stormer FA. 1984. Night-roosting habitat of scaled quail. *J Wildl Manage* 48: 191-197.
- U.S. Geological Survey (USGS). 1990. Land Use and Land Cover Digital Data from 1:250,000- and 1:100,000-Scale Maps: Data User's Guide 4. United States Department of the Interior, U.S. Geological Survey, Reston, VA.
- Uuemaa E, Antrop M, Roosaare J, Marja R, Mander U. 2009. Landscape metrics and indices: an overview of their use in landscape research. *Living Rev Landscape Res* 3: 1-28.
- Wallmo OC. 1957. Ecology of scaled quail in west Texas. PhD Dissertation. Texas A&M University, College Station, TX, USA.
- Wilson DE. 1992. Small game research and surveys: quail harvest regulations. Texas Parks and Wildlife Department, Performance Report W-126-R-1, Job No. 4.01, Austin, Texas.
- Wilson MH, Crawford JA. 1987. Habitat selection by Texas bobwhites and chestnut-bellied scaled quail in south Texas. *J Wildl Manage* 51: 575-582.
- Wu XB, Smeins FE, Slack RD. 2002. Fundamentals of Ecology Laboratory Manual. 3rd ed. Kendall Hunt Publishing Company, Dubuque, IA.