

# Resistance to Thyroid Hormone Syndrome Mutation in *THRB* and *THRA*: A Review

Jung Eun Moon

Division of Pediatric Endocrinology, Kyungpook National University Children's Hospital, Daegu, Korea

Resistance to thyroid hormone syndrome (RTH) is a genetic disease caused by the mutation of either the thyroid hormone receptor- $\beta$  (*THRB*) gene or the thyroid hormone receptor- $\alpha$  (*THRA*) gene. RTH caused by *THRB* mutations (RTH- $\beta$ ) is characterized by the target tissue's response to thyroid hormone, high levels of triiodothyronine and/or thyroxine, and inappropriate secretion of thyroid-stimulating hormone (TSH). *THRA* mutation is characterized by hypothyroidism that affects gastrointestinal, neurological, skeletal, and myocardial functions. Most patients do not require treatment, and some patients may benefit from medication therapy. These syndromes are characterized by decreased tissue sensitivity to thyroid hormones, generating various clinical manifestations. Thus, clinical changes of resistance to thyroid hormones must be recognized and differentiated, and an approach to the practice of personalized medicine through an interdisciplinary approach is needed.

**Key words:** Thyroid hormone receptor- $\beta$ , Thyroid hormone receptor- $\alpha$ , Thyroid hormone

## REVIEW ARTICLE

**Received:** October 18, 2023

**Revised:** October 30, 2023

**Accepted:** October 30, 2023

**Correspondence to:** Jung Eun Moon  
Division of Pediatric Endocrinology, Kyungpook National University Children's Hospital, 807 Hoguk-ro, Buk-gu, Daegu 41404, Korea  
**Tel:** +82-10-4782-8315  
**Fax:** +82-53-425-6683  
**E-mail:** subuya@daum.net

**ORCID**

<https://orcid.org/0000-0001-9786-7898>



Copyright © 2023, Interdisciplinary Society of Genetic & Genomic Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDeriv License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), provided the original work is properly cited.

## INTRODUCTION

The thyroid gland secretes hormones, namely, thyroxine (T<sub>4</sub>) and triiodothyronine (T<sub>3</sub>). Subsequently, 80% of T<sub>4</sub> is converted to T<sub>3</sub> in the target organ. Various biological actions of thyroid hormones (THs) are mediated by proteins and receptors located in the cell membrane that allows hormones to enter cells. Deiodinases that convert T<sub>4</sub> and T<sub>3</sub> to metabolites with lower iodine content control the cytoplasmic metabolic action of THs. Finally, nuclear receptors bind more closely or exclusively to T<sub>3</sub>; thus, the genomic actions of THs are mediated by T<sub>3</sub> [1]. T<sub>3</sub> binds to nuclear receptors so that gene expression and isoforms of TH receptors (TR $\alpha$ 1, TR $\alpha$ 2, TR $\beta$ 1, and TR $\beta$ 2) are encoded by *THRA* and *THRB*, located in chromosomes 17q21.1 and 3p24.2 [2,3]. Resistance to TH syndromes (RHT) have a genetic origin and are characterized by decreased tissue sensitivity to THs. Most cases are of mutations in *THRB* that encodes for one of the two types of the T<sub>3</sub> nuclear receptor, TR $\beta$ 1 and TR $\beta$ 2 [4].

## RTH CAUSED BY *THRB* MUTATIONS (RTH- $\beta$ )

RTH- $\beta$  is a rare condition caused by *THRB* mutations [5]. RTH was first described in 1967 in a family with deaf congenital disease and pointed epiphysis. Disease prevalence is unclear because routine detection tests for congenital hypothyroidism are based solely on the determination of TSH, which is often normal under this condition [6]. Most RTH- $\beta$  cases are caused by heterozygous variants in *THRB* [4]. *THRB* mutations have been associated with a T<sub>3</sub> receptor binding alteration, retaining homo- and heterodimerization capability, or with an aberrant ability to dissociate from the correlator in the presence of T<sub>3</sub>. The series of

hydrophobic repeats are crucial for homodimer and heterodimer formation in TH receptors, which do not usually exhibit mutations. *THRB* mutations are predominantly grouped into three “hot” regions in the c-terminal region of *THRB*, involved with T3 hormone binding of protein [7]. A retrospective study of 14 patients with RTH and late manifestations of RTH syndrome identified that all had point mutations, located between exons 7–10 within the union domain to T3. Seven patients had widespread resistance to TH, and four showed euthyroidism. However, the other three patients experienced thyroid dysfunction before the age of 10 years, and all patients with early onset had a short stature and different degrees of learning disruption [8]. RTH- $\beta$  is characterized by high levels of T4, T3, and rT3 accompanied by normal or high serum TSH levels [4]. The clinical symptoms of patients with RTH vary from manifestations of hyperthyroidism in organs primarily expressing TR $\alpha$  (heart and bone) to hypothyroidism in tissues expressing TR $\beta$  (hypophysis, brain, and thyroid gland). Hypercholesterolemia, delayed growth, and learning disorders (suggestive of hypothyroidism), can coexist with weight loss, heat intolerance, hyperactivity, and tachycardia [9].

The clinical manifestations of RTH are classified into two subgroups according to the absence or presence of symptoms of thyrotoxicosis, selective pituitary resistance, and generalized resistance of TH (GRTH). Patients with selective pituitary resistance show variable symptoms of hyperthyroidism; however, patients with GRTH exhibit “compensated hypothyroidism,” which controls the effect of high TH levels. In several cohorts, the cardiac phenotypes of patients with different mutations, which included tachycardia and palpitations, were examined, although flutter or atrial fibrillation has been reported up to 20% of patients. No heart failure cases were reported, and no correlation was found between the genotype and the cardiac phenotype [10]. The main differential diagnosis of this condition is thyrotropoma, a TSH-producing pituitary tumor. TSH-producing pituitary tumors are characterized by a biochemical pattern of TH similar to RHT; however, unlike RTH, it presents an increase in  $\alpha$ -subunits and sex hormone binding globulin. The genetic study is definitive for differentiation, and approximately 15% of cases have failed to identify the genetic cause [11].

The treatment mainly aims to maintain a balance between the overstimulation of the tissues expressing TR $\beta$  in patients with RTH [5]. Currently, no definitive therapy is available to correct the molecular defect that causes RTH syndrome, and most patients do not require treatment [5,9]. Patients with tachy-

**Table 1.** Manifestations of resistance to thyroid hormone syndrome

	RTH- $\beta$	RTH- $\alpha$
Mutation receptor	TR- $\beta$ 1, TR- $\beta$ 2	TR- $\alpha$ 1, TR- $\alpha$ 2
Mutation gene	Thyroid hormone receptor- $\beta$ ( <i>THR-<math>\beta</math></i> ) gene	Thyroid hormone receptor- $\alpha$ ( <i>THR-<math>\alpha</math></i> ) gene
Clinical manifestations	Vary from hyperthyroidism to hypothyroidism or simple goiter	Anemia, Constipation, Growth retardation, Developmental delay, Motor and Cognitive development delay
Treatment	Most patients do not require treatment $\beta$ -blocker (with tachycardia) TRIAC or DT4 (with thyrotoxicosis) Dopamine drugs Somatostatin analogs	Levothyroxine

RTH, Resistance to thyroid hormone syndrome; TR, thyroid hormone receptors; TRIAC, triiodothyroacetic acid; DT4, dextrothyroxine.

cardia and palpitations at rest may benefit from  $\beta$ -blocker therapy. If thyroid toxicity symptoms are present, triiodothyroacetic acid (TRIAC) or dextrothyroxine (DT4) can be administered to treat thyrotoxicosis. This compound, through a feedback mechanism, reduces TSH secretion and decreases circulating T4 levels. TRIAC given at a dose of 1.4–2.8 mg/day divided into two or three administrations is effective in both pediatric and adult patients with RTH [6]. The use of dopamine drugs and somatostatin analogs has limited success because TSH secretion quickly escapes the inhibitory effects of both drugs, as T4 reduction strongly stimulates TSH secretion (Table 1) [4,9].

### RTH CAUSED BY *THRA* MUTATIONS (RTH- $\alpha$ )

RTH- $\alpha$  is characterized by tissue-dependent hypothyroidism findings, and >40 *THRA* mutations have been recorded [12]. Most mutations selectively involve the c-terminal domain of TR $\alpha$ , generating severe loss of function and other mutation types associated with a variable-form function of TR $\alpha$ 1 [13]. In these cases, the phenotype is characterized by manifestations of hypothyroidism that affects gastrointestinal, neurological, skeletal, and myocardial functions. Clinical characteristics associated with hypothyroidism (e.g., macrocephaly, hypertelorism, flattened nasal bridge, and a broad face) became more evident with time, together with poor linear growth, resulting in a dysmorphic appearance. Patients with RTH- $\alpha$  have normal hormonal response in the pituitary hypothalamic axis and liver. The severity of the clinical phenotype appears to be associated with the mutation location and type in *THRA*. When hetero-

zygous mutations in this gene are present, abnormalities include anemia, constriction, and delayed growth and development [9]. The A263V mutation of TR $\alpha$ 1 showed partial loss of function, with higher T3 levels. TR $\alpha$ 2 has no mutation-related phenotypes identified; however, some unusual clinical characteristics, such as micrognasia, clavicular agenesis, hyperparathyroidism, and chronic diarrhea, were reported in an adult woman with a different mutation (N359Y), common for both TR $\alpha$ 1 and TR $\alpha$ 2 [14]. In RTH- $\alpha$ , serum T3 levels can be high or normal, T4 and rT3 levels are in the normal range or low, and TSH levels are normal or slightly high [3].

Levothyroxine therapy generates the normalization of T4 levels, with additional increase in T3 and suppressed TSH levels, suggesting conserved negative feedback of TH over TSH secretion (Table 1) [3].

## CONCLUSION

RTH is a group of genetic-related syndromes that result from different mutations, either in the genes they encode for receptors. These syndromes are characterized by decreased tissue sensitivity to THs, generating various clinical manifestations. Thus, clinical changes of resistance to TH must be recognized and differentiated, and an approach to the practice of personalized medicine through an interdisciplinary approach is needed.

## ACKNOWLEDGMENTS

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

## REFERENCES

- Bernal J. Síndromes de resistencia a las hormonas tiroideas. *Endocrinol Nutr* 2011;58:185-96. doi: 10.1016/j.endonu.2011.02.001.
- Bianco AC, Kim BW. Deiodinases: implications of the local control of thyroid hormone action. *J Clin Invest* 2006;116:2571-9. doi: 10.1172/JCI29812.
- Van Gucht AL, Moran C, Meima ME, Visser WE, Chatterjee K, Visser TJ, et al. Resistance to thyroid hormone due to heterozygous mutations in thyroid hormone receptor alpha. *Curr Top Dev Biol* 2017;125:337-55. doi: 10.1016/bs.ctdb.2017.02.001.
- Groeneweg S, Peeters RP, Visser TJ, Visser WE. Therapeutic applications of thyroid hormone analogues in resistance to thyroid hormone (RTH) syndromes. *Mol Cell Endocrinol* 2017;458:82-90. doi: 10.1016/j.mce.2017.02.029.
- Vela A, Pérez-Nanclares G, Ríos I, Rica I, Portillo N, Castaño L. Thyroid hormone resistance from newborns to adults: a Spanish experience. *J Endocrinol Invest* 2019;42:941-9. doi: 10.1007/s40618-019-1007-4.
- Singh BK, Yen PM. A clinician's guide to understanding resistance to thyroid hormone due to receptor mutations in the TR $\alpha$  and TR $\beta$  isoforms. *Clin Diabetes Endocrinol* 2017;15:8. doi: 10.1186/s40842-017-0046-z.
- Censi S, Barollo S, Watutantrige-Fernando S, Manso J, Ferrara AM, Mian C. A Novel thyroid hormone receptor beta mutation (G357R) in a family with resistance to thyroid hormone beta: extending the borders of the "hot" region in the THRB Gene. *Thyroid* 2019;29:449-51. doi: 10.1089/thy.2018.0201.
- Han R, Ye L, Jiang X, Zhou X, Billon C, Guan W, et al. Characteristics of patients with late manifestation of resistance thyroid hormone syndrome: a single-center experience. *Endocrine* 2015;50:689-97. doi: 10.1007/s12020-015-0622-x.
- Persani L, Campi I. Syndromes of resistance to thyroid hormone action. En: genetics of endocrine diseases and syndromes. *Genetics of Endocrine Diseases and Syndromes* 2019;55:84. doi: 10.1007/978-3-030-25905-1\_5.
- Illouz F, Briet C, Mirebeau-Prunier D, Bouhours-Nouet N, Coutant R, Sibilia P, et al. Cardiac complications of thyroid hormone resistance syndromes. *Ann Endocrinol* 2021;82:167-9. doi: 10.1016/j.ando.2020.03.008.
- Sun H, Cao L, Zheng R, Xie S, Liu C. Update on resistance to thyroid hormone syndrome $\beta$ . *Ital J Pediatr* 2020;46:168. doi: 10.1186/s13052-020-00929-x.
- National Library of Medicine (NIH). ClinVar [Internet]. <https://www.ncbi.nlm.nih.gov/clinvar/>
- Moran C, Agostini M, McGowan A, Schoenmakers E, Fairall L, Lyons G, et al. Contrasting phenotypes in resistance to thyroid hormone alpha correlate with divergent properties of thyroid hormone receptor  $\alpha$ 1 mutant proteins. *Thyroid* 2017;27:973-82. doi: 10.1089/thy.2017.0157.
- Moran C, Chatterjee K. Resistance to thyroid hormone  $\alpha$ -emerging definition of a disorder of thyroid hormone action. *J Clin Endocrinol Metab* 2016;101:2636-9. doi: 10.1210/jc.2016-2317.