



Corrigendum: The seven-year cumulative survival rate of Osstem implants

Young-Kyun Kim¹, Bum-Su Kim¹, Pil-Young Yun¹, Sang-Un Mun²,
Yang-Jin Yi³, Su-Gwan Kim⁴, Kyung-In Jeong⁴

¹Department of Oral and Maxillofacial Surgery, ²Department of Orthodontics, ³Department of Prosthodontics,
Section of Dentistry, Seoul National University Bundang Hospital, Seongnam, ⁴Department of Oral and Maxillofacial Surgery,
School of Dentistry, Chosun University, Gwangju, Korea

J Korean Assoc Oral Maxillofac Surg 2014;40:68-75. <http://dx.doi.org/10.5125/jkaoms.2014.40.2.68>

This correction is being published to correct the errors of sentences and spelling in abstract.
The authors apologize for this error.

Before correction

Materials and Methods: A total of 105 patients who had 467 Osstem implants that were placed at the Section of Dentistry, Seoul National University Bundang Hospital (Seongnam, Korea) from June 2003 through December 2005 were analyzed. The life table method and a cross-tabulation analysis, log rank test were used to evaluate the survival curve and the influence that the prognostic factors. The prognostic factors, i.e., age and gender of patients, diameter and length, type of implants, bone graft history and loading time were determined with a Cox proportional hazard model based on logistic regression analysis.

After correction

Materials and Methods: A total of 105 patients who had 467 Osstem implants that were placed at the Section of Dentistry, Seoul National University Bundang Hospital (Seongnam, Korea) from June 2003 through December 2005 were analyzed. The life table method and a cross-tabulation analysis were performed to evaluate the cumulative survival rate. The log rank test was used to evaluate the survival curve. The influence that the prognostic factors had on the survival rate was determined with a Cox proportional hazard model based on logistic regression analysis.