



Common risk factors for postoperative pain following the extraction of wisdom teeth

Vahid Rakhshan^{1,2}

¹*Iranian Tissue Bank and Research Center, Tehran University of Medical Sciences,*

²*Department of Dental Anatomy and Morphology, Dental Branch, Islamic Azad University, Tehran, Iran*

Abstract (J Korean Assoc Oral Maxillofac Surg 2015;41:59-65)

The extraction of third molars is a common task carried out at dental/surgery clinics. Postoperative pain is one of the two most common complications of this surgery, along with dry socket. Knowledge of the frequent risk factors of this complication is useful in determining high-risk patients, planning treatment, and preparing the patients mentally. Since the risk factors for postoperative pain have never been summarized before while the risk factors for dry socket have been highly debated, this report summarizes the literature regarding the common predictors of postextraction pain. Except for surgical difficulty and the surgeon's experience, the influences of other risk factors (age, gender and oral contraceptive use) were rather inconclusive. The case of a female gender or oral contraceptive effect might mainly be associated with estrogen levels (when it comes to dry socket), which can differ considerably from case to case. Improvement in and unification of statistical and diagnostic methods seem necessary. In addition, each risk factor was actually a combination of various independent variables, which should instead be targeted in more comprehensive studies.

Key words: Pain, Risk factors, Third molar, Extraction

[paper submitted 2014. 11. 16 / revised 1st 2014. 11. 25, 2nd 2014. 12. 4 / accepted 2014. 12. 11]

I. Introduction

One of the most common procedures carried out in dental clinics and the most frequent task done at oral and maxillofacial surgery clinics is the extraction of wisdom teeth. This procedure is frequently followed by complications in the mandible¹⁻³, including both iatrogenic (e.g., nerve injury, bone fractures, etc.) and inflammatory ones, such as dry socket, postoperative pain, delayed healing, postoperative infection, hematoma, swelling, trismus, etc.^{2,4-6}. Although the overall complication rate might be generally low, and most complications are minor^{4,7-9}, this surgery is so frequent that the population's morbidity of complications may be noticeable; thus, identifying methods to control or reduce them is

a major concern^{4,10,11}. Besides, not all complications are rare. There are frequent and debilitating complications as well, including postoperative pain.

Pain is also one of the most common postoperative complications of extraction^{6,12-15} and might be caused by the release of pain mediators from the injured tissues^{3,15}. Pain is an important factor in clinical practice^{6,16} and could even discourage patients from seeking dental treatment^{15,17,18}. It begins after the anesthesia subsides and reaches its peak levels during the first postoperative day^{15,19,20}. If dry socket or infection occur, the onset of inflammation will complicate alleviation of postoperative pain^{5,15,20-26}.

In the setting of elective operations, such as third molar removal, patients demand to know the risks, benefits and postoperative quality of life of these procedures^{15,17,27}. The knowledge of the risk factors of postsurgical complications has clinical implications in treatment planning, patient management and prognosis^{15,19,21,24,28-31}. This essay briefly reviews the most common risk factors of pain following third molar removal.

Vahid Rakhshan

#22 Behruz Alley, Karegar St. Tehran 14188-36783, Iran

TEL: +982166929055 FAX: +982166902923

E-mail: vahid.rakhshan@gmail.com

ORCID: <http://orcid.org/0000-0002-9503-3133>

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2015 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.

II. Materials and Methods

The internet was searched to find relevant articles published before July 2014 regarding the risk factors of postextraction pain. The search engines used were Google Scholar, Pubmed/MEDLINE, ISI Web of Science, and Scopus. The keywords were as follows: “third molar”, “wisdom tooth”, “wisdom teeth”, “extraction”, “removal”, “postoperative pain”, “postsurgical pain”, “risk factor”, “prognostic factor”, and “predictor”.

More than 800 unique articles were initially found. All these article “titles” were reviewed to narrow down the scope of the search to more relevant articles, according to the eligibility criteria of the presentation of results of original research or short communications regarding pain perceived after the extraction of the third molars before July 2014; the time span was open to all articles published before this date. The articles of interest were collected and evaluated. The reference lists of the located articles were also consulted to identify additional relevant reports. Each article or abstract was read at least twice, and the proper information was aggregated. Other more general topics were also researched for the sake of discussing the matters.

III. Results

1. Oral hygiene

The effects of hygiene maintenance on postoperative pain have not been widely assessed except in a few English and non-English articles. Sáez Cuesta et al.³² extracted 100 wisdom teeth and found that patients with poor oral hygiene before surgery experienced higher pain levels during the first 6 postoperative hours. Peñarrocha et al.⁵ explored 190 impacted third molars and found that pain increased with increasing lack of care to hygiene. Larrazábal et al.³³ asserted that pain increases were correlated with less brushing before surgery and also during the first postoperative week.

2. Difficulty of the extraction procedure and trauma

An association between different aspects of surgical difficulty (such as the impaction level and angle, extent of bone removal or length of surgery) and pain or paresthesia has been assessed in most previous studies^{6,14,15,28,29,34,37} with a few exceptions^{12,38}. Lago-Méndez et al.¹³, Pedersen¹⁴, Baqain et al.³⁹, and de Santana-Santos et al.⁶ stated that lengthier sur-

geries leave more painful sockets. However, Benediktsdóttir et al.²⁹ found that simply the root morphology and level of impaction were correlated to postoperative pain; they did not identify any impact of the actual surgery time²⁹. Grossi et al.³⁸ also did not find a significant relationship between the elapsed time of surgery and the level of postoperative discomfort. Seymour et al.⁴⁰ stated that the postoperative pain level might not be dependent upon the operator or the extent of surgical trauma as estimated by operating time and radiographic score. On the other hand, Oikarinen³⁵, Garcia Garcia et al.³⁶, Haraji and Rakhshan¹⁵, and Yuasa and Sugiura³⁷ found that more difficult operations were more painful. The release of more inflammatory factors and proximity to the nerve might produce more intense pain in some difficult cases. This pain might shift to paresthesia or anesthesia in cases with direct injury to a nerve³⁴. It should be noted that according to recent research, radiographic indicators alone (i.e., Winter, Pell and Gregory, Pederson)^{14,41-43} cannot totally identify the extent of the difficulty of surgery^{6,13,29,34,44,45}. Additionally, not all the difficulty determinants are necessarily prognostic factors for complications⁴⁶. Therefore, better methods should be used to estimate surgical difficulty, or the latent variables should be assessed independently instead³¹.

3. The operator's expertise

An experienced surgeon might carry out a cleaner, less traumatic and yet faster operation than someone new to the procedure. Additionally, patients might trust experienced clinicians more. These factors (trauma, duration of surgery and anxiety) can play important roles in inducing complications^{8,12,22,28-30,47-53}. Therefore, expert clinicians might obtain better results^{8,12,34,48-51,54-56}. However, the evidence is controversial, as some authors did not denote a link between the surgeon's skill and the patient's postoperative pain²⁹. Some of these “experience levels” were actually different terms of undergraduate study and thus indicated little about the surgeon's expertise⁵⁷. A surgeon's experience might reduce the postoperative pain only within a short period after the surgery but may have no influence on the duration or intensity of longer pains¹². This discrepancy might contribute to the controversial result depending on the time at which the pain is assessed.

4. Tobacco smoking

Smoking might increase pain by reducing blood supply in

the alveolar socket^{33,49}. Meechan et al.⁴⁹ asserted that heavy smokers have a high chance of poor filling of their extraction sockets with blood. They also found a correlation between this phenomenon measured immediately after the extraction and painful sockets⁴⁹. Nevertheless, to the author's knowledge, only certain authors have found a significant link between postoperative pain and smoking³³. The only relationship that was found was between the pains perceived on the first post-surgical day and with postsurgical smoking³³. The same study and many others did not correlate preoperative smoking with postsurgical pain in their total samples^{4,12,15,33,38,39}. In the studies of Grossi et al.³⁸ and Heng et al.², a greater amount of pain perceived by females was associated with smoking^{2,38}. Grossi et al.³⁸ suggested that smoking only affects the perception of pain by females. However, Haraji and Rakhshan¹⁵ adjusted for the role of gender while assessing the effect of smoking on pain. In their analysis, smoking had no significant effects on postoperative pain on the first or third postoperative days or in general. They also assessed the interaction of gender and smoking, and no significant results appeared¹⁵.

5. Gender

Gender is a crucial variable that should be considered when designing and analyzing the findings of studies in all areas and at all levels of biomedical and health-related research³⁸. This issue has been ignored in the past and has gained popularity only in the last few decades⁵⁸. The association between clinical pain and gender is not a simple one, but females have reported more frequent pains compared to men in terms of various anatomic regions, neuropathic conditions, chronic musculoskeletal pains, temporomandibular pains, facial pains, toothaches, etc.⁵⁸⁻⁶⁴. Postoperative pain studies lack standardization and are at some points conflicting; however, in general, it could be inferred that women might experience pain more often and to a greater extent than men^{58,65-68}. Although the research in this regard is rather scarce in terms of postextraction pain by gender, the aforementioned results could imply that females might have a higher sensitivity to pain stimuli perhaps due to psychosocial factors (mood, sex role beliefs, pain coping strategies, and pain-related expectancies), catastrophizing and sex hormones^{38,69,70}. Also the thinner mandible of women might render them more vulnerable to pain and some complications after dental procedures^{6,71}. Some authors have reported more intense postsurgical pains^{28,29,40}, longer symptom recovery times^{17,47} or neurosensory deficit in females³⁴. However, many others found

conflicting results^{4,12,15,37,38,46,57,72,73}. Capuzzi et al.¹² reported a greater extent of pain in males. Yuasa and Sugiura³⁷ declared that postoperative swelling and morbidity but not pain might be greater in females. de Santana Santos et al.⁴¹ observed significantly more pain in females only during the first 4 and 12 postoperative hours, but at the 24th and 48th postoperative hours, the greater pain intensity in women did not reach a level of significance⁴¹. This controversy might be rooted in various missing latent variables (e.g., hormonal, psychological or genetic differences, etc.).

6. Oral contraceptives

Contraceptive consumption might be less likely to affect or confound pain-related results^{4,12,38,74}, although a few studies have reported on its positive role in this regard as well⁷⁵. Regardless, modern contraceptive pills contain considerably lower doses of estrogen and therefore have a reduced role compared with those of the past^{34,76}.

7. Age

The production and process of sensory stimuli might be influenced by aging⁷⁷⁻⁷⁹. The elderly could be at higher risk of complications, such as severe pain and sensory disturbances^{4,12,38,80,81}, possibly because of this group's poorer healing potential, denser bones and completed dental roots^{4,28,34,82}. Some investigators have observed significant deteriorating effects of aging on pain^{9,12,28,81}. Blondeau and Daniel³⁴ reported increased neurosensory problems in patients older than 24 years. However, other studies have not identified such a role^{29,39,83}. Adeyemo et al.⁸⁴ and Bui et al.⁴ found no significant association between age and complications. Yuasa and Sugiura³⁷ reported a significant influence of age on swelling and collective postprocedural morbidity but not pain. Grossi et al.³⁸ observed a significant association between patients older than 23 years and merely severe trismus but not pain either reported subjectively by the patients or implied by the number of painkillers taken. Benediktsdóttir et al.²⁹ found no correlation between age and discomfort, despite their finding indicating that surgery could last significantly longer in older patients. In the study of Capuzzi et al.¹², younger participants reported less pain in the first postextraction day, but the number of painkillers taken was not correlated with age. Haraji and Rakhshan¹⁵ studied younger patients and showed that when the effects of the operation difficulty, smoking and gender were not controlled for, younger people might show

significantly greater pain. However, when these factors were adjusted for, younger patients showed a borderline significantly greater amount of pain only for the first postoperative day but not on the third postsurgical day¹⁵. The narrow range of patient ages could mask such an effect, since third molar extraction is usually indicated in young ages³⁸, and debilitating effects of age might appear in older ages⁷⁹. Some authors have advocated the removal of impacted molars in young adults to avoid severe or permanent sequelae^{9,34,81}. Nonetheless, if the assumption is not confirmed, early prophylactic extraction of wisdom teeth, which is common in Europe and America, might not be justifiable^{38,84,85}.

IV. Conclusion

Based on the number of studies agreeing that a certain po-

tential factor might likely be a real risk factor (Table 1), the trauma of surgery and experience of the surgeon were more likely to be causative or risk factors of pain. High levels of estrogen were not necessarily a risk factor for pain. Evidence suggesting a higher incidence of postextraction pain in females was outnumbered by reports that refuted such an association. The effect of age remained inconclusive. Although only a few studies regarding the effect of oral hygiene on postoperative pain exist, it was shown to be effective in that regard as well. There were at least three obstacles for detecting possible links between pain and risk factors: consumption of painkillers and antibiotics by the patients after surgery, which act as efficient confounders^{15,20,25}, as well as poorer statistical approaches, and confusion of pain caused by a dry socket or infection with pain caused only by the surgery and also with discomfort. Except for a few essays¹⁵, almost

Table 1. A summary of studies supporting or not supporting the role of the searched risk factors (some studies fit both criteria)

Factor	Supporting study	Studies failing to support the risk factor
Oral hygiene	Sáez Cuesta et al. ³² Peñarrocha et al. ⁵ Larrazábal et al. ³³	-
Operation difficulty, duration or trauma as risk indicators of postoperative pain, sensory disruption or discomfort	Lago-Méndez et al. ¹³ Pedersen ¹⁴ Baqain et al. ³⁹ de Santana-Santos et al. ⁶ Oikarinen ³⁵ Garcia Garcia et al. ³⁶ Haraji and Rakhshan ¹⁵ Yuasa and Sugiura ³⁷	Capuzzi et al. ¹² Benediktsdóttir et al. ²⁹ Grossi et al. ³⁸ Seymour et al. ⁴⁰
The expertise of the surgeon	-	Benediktsdóttir et al. ²⁹ Capuzzi et al. ¹²
Smoking	Meechan et al. ⁴⁹ Larrazábal et al. ³³ Grossi et al. ³⁸ Heng et al. ²	Bui et al. ⁴ Haraji and Rakhshan ¹⁵ Larrazábal et al. ³³ Capuzzi et al. ¹² Grossi et al. ³⁸ Baqain et al. ³⁹
Gender	In favor of female gender Phillips et al. ²⁸ de Santana Santos et al. ⁴¹ Benediktsdóttir et al. ²⁹ Seymour et al. ⁴⁰ In favor of male gender Capuzzi et al. ¹²	Eshghpour et al. ⁵⁷ Abu Younis and Abu Hantash ⁷² Barbosa-Rebellato et al. ⁷³ Carvalho and do Egito Vasconcelos ⁴⁶ Haraji and Rakhshan ¹⁵ Yuasa and Sugiura ³⁷ Bui et al. ⁴ Grossi et al. ³⁸
Age	Osborn et al. ⁹ Bruce et al. ⁸¹ Blondeau and Daniel ³⁴ Capuzzi et al. ¹² Phillips et al. ²⁸	Capuzzi et al. ¹² Akadiri et al. ⁸³ Adeyemo et al. ⁸⁴ Bui et al. ⁴ Yuasa and Sugiura ³⁷ Grossi et al. ³⁸ Benediktsdóttir et al. ²⁹ Capuzzi et al. ¹² Haraji and Rakhshan ¹⁵ Baqain et al. ³⁹

Vahid Rakhshan: Common risk factors for postoperative pain following the extraction of wisdom teeth. J Korean Assoc Oral Maxillofac Surg 2015

all previous studies have failed to distinguish dry socket or infection pain from pain caused by the surgery alone when evaluating the risk factors for postoperative pain. Future studies are warranted to account for each type of pain independently. Another issue ignored in almost all studies except a few^{15,28,31} is that the variables that affect pain likely interact with each other. Therefore, analyses not accounting for the interactions are less accurate and less useful than those that consider a broader clinical picture³¹.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

References

- Nordenram A. Postoperative complications in oral surgery. A study of cases treated during 1980. *Swed Dent J* 1983;7:109-14.
- Heng CK, Badner VM, Clemens DL, Mercer LT, Mercer DW. The relationship of cigarette smoking to postoperative complications from dental extractions among female inmates. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;104:757-62.
- El-Soud NA, El Shenawy H. A randomized double blind clinical study on the efficacy of low level laser therapy in reducing pain after simple third molar extraction. *Maced J Med Sci* 2010;3:303-6.
- Bui CH, Seldin EB, Dodson TB. Types, frequencies, and risk factors for complications after third molar extraction. *J Oral Maxillofac Surg* 2003;61:1379-89.
- Peñarrocha M, Sanchis JM, Sáez U, Gay C, Bagán JV. Oral hygiene and postoperative pain after mandibular third molar surgery. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2001;92:260-4.
- de Santana-Santos T, de Souza-Santos JA, Martins-Filho PR, da Silva LC, de Oliveira E Silva ED, Gomes AC. Prediction of postoperative facial swelling, pain and trismus following third molar surgery based on preoperative variables. *Med Oral Patol Oral Cir Bucal* 2013;18:e65-70.
- Calhoun NR. Dry socket and other postoperative complications. *Dent Clin North Am* 1971;15:337-48.
- Muhonen A, Ventä I, Ylipaavalniemi P. Factors predisposing to postoperative complications related to wisdom tooth surgery among university students. *J Am Coll Health* 1997;46:39-42.
- Osborn TP, Frederickson G Jr, Small IA, Torgerson TS. A prospective study of complications related to mandibular third molar surgery. *J Oral Maxillofac Surg* 1985;43:767-9.
- Lopes V, Mumenya R, Feinmann C, Harris M. Third molar surgery: an audit of the indications for surgery, post-operative complaints and patient satisfaction. *Br J Oral Maxillofac Surg* 1995;33:33-5.
- Berge TI, Bøe OE. Predictor evaluation of postoperative morbidity after surgical removal of mandibular third molars. *Acta Odontol Scand* 1994;52:162-9.
- Capuzzi P, Montebugnoli L, Vaccaro MA. Extraction of impacted third molars. A longitudinal prospective study on factors that affect postoperative recovery. *Oral Surg Oral Med Oral Pathol* 1994;77:341-3.
- Lago-Méndez L, Diniz-Freitas M, Senra-Rivera C, Gude-Sampedro F, Gándara Rey JM, García-García A. Relationships between surgical difficulty and postoperative pain in lower third molar extractions. *J Oral Maxillofac Surg* 2007;65:979-83.
- Pedersen A. Interrelation of complaints after removal of impacted mandibular third molars. *Int J Oral Surg* 1985;14:241-4.
- Haraji A, Rakhshan V. Chlorhexidine gel and less difficult surgeries might reduce post-operative pain, controlling for dry socket, infection and analgesic consumption: a split-mouth controlled randomised clinical trial. *J Oral Rehabil* 2015;42:209-19.
- Slade GD, Foy SP, Shugars DA, Phillips C, White RP Jr. The impact of third molar symptoms, pain, and swelling on oral health-related quality of life. *J Oral Maxillofac Surg* 2004;62:1118-24.
- Bienstock DA, Dodson TB, Perrott DH, Chuang SK. Prognostic factors affecting the duration of disability after third molar removal. *J Oral Maxillofac Surg* 2011;69:1272-7.
- Wardle J. Dental pessimism: negative cognitions in fearful dental patients. *Behav Res Ther* 1984;22:553-6.
- Susarla SM, Blaeser BF, Magalnick D. Third molar surgery and associated complications. *Oral Maxillofac Surg Clin North Am* 2003;15:177-86.
- Haraji A, Rakhshan V, Khamverdi N, Alishahi HK. Effects of intra-alveolar placement of 0.2% chlorhexidine bioadhesive gel on dry socket incidence and postsurgical pain: a double-blind split-mouth randomized controlled clinical trial. *J Orofac Pain* 2013;27:256-62.
- Blum IR. Contemporary views on dry socket (alveolar osteitis): a clinical appraisal of standardization, aetiopathogenesis and management: a critical review. *Int J Oral Maxillofac Surg* 2002;31:309-17.
- Bloomer CR. Alveolar osteitis prevention by immediate placement of medicated packing. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000;90:282-4.
- Noroozi AR, Philbert RF. Modern concepts in understanding and management of the "dry socket" syndrome: comprehensive review of the literature. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:30-5.
- Cardoso CL, Rodrigues MT, Ferreira Júnior O, Garlet GP, de Carvalho PS. Clinical concepts of dry socket. *J Oral Maxillofac Surg* 2010;68:1922-32.
- Caso A, Hung LK, Beirne OR. Prevention of alveolar osteitis with chlorhexidine: a meta-analytic review. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2005;99:155-9.
- Kolokythas A, Olech E, Miloro M. Alveolar osteitis: a comprehensive review of concepts and controversies. *Int J Dent* 2010;2010:249073.
- White RP Jr, Shugars DA, Shafer DM, Laskin DM, Buckley MJ, Phillips C. Recovery after third molar surgery: clinical and health-related quality of life outcomes. *J Oral Maxillofac Surg* 2003;61:535-44.
- Phillips C, White RP Jr, Shugars DA, Zhou X. Risk factors associated with prolonged recovery and delayed healing after third molar surgery. *J Oral Maxillofac Surg* 2003;61:1436-48.
- Benediktsdóttir IS, Wenzel A, Petersen JK, Hintze H. Mandibular third molar removal: risk indicators for extended operation time, postoperative pain, and complications. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;97:438-46.
- Halabi D, Escobar J, Muñoz C, Uribe S. Logistic regression analysis of risk factors for the development of alveolar osteitis. *J Oral Maxillofac Surg* 2012;70:1040-4.
- Haraji A, Rakhshan V. Single-dose intra-alveolar chlorhexidine gel application, easier surgeries, and younger ages are associated with reduced dry socket risk. *J Oral Maxillofac Surg* 2014;72:259-65.
- Sáez Cuesta Ú, Peñarrocha Diago M, Sanchis Bielsa JM, Gay Escoda C. Estudio del postoperatorio de 100 terceros molares mandibulares incluidos, en relación a la edad, el sexo, el tabaco y la higiene bucal. *RCOE Revista del Consejo General de Colegios de Odontólogos y Estomatólogos de España* 1999;4:471-5.
- Larrazábal C, García B, Peñarrocha M, Peñarrocha M. Influence of oral hygiene and smoking on pain and swelling after surgical extraction of impacted mandibular third molars. *J Oral Maxillofac*

- Surg 2010;68:43-6.
34. Blondeau F, Daniel NG. Extraction of impacted mandibular third molars: postoperative complications and their risk factors. *J Can Dent Assoc* 2007;73:325.
 35. Oikarinen K. Postoperative pain after mandibular third-molar surgery. *Acta Odontol Scand* 1991;49:7-13.
 36. Garcia Garcia A, Gude Sampedro F, Gandara Rey J, Gallas Torreira M. Trismus and pain after removal of impacted lower third molars. *J Oral Maxillofac Surg* 1997;55:1223-6.
 37. Yuasa H, Sugiura M. Clinical postoperative findings after removal of impacted mandibular third molars: prediction of postoperative facial swelling and pain based on preoperative variables. *Br J Oral Maxillofac Surg* 2004;42:209-14.
 38. Grossi GB, Maiorana C, Garramone RA, Borgonovo A, Creminelli L, Santoro F. Assessing postoperative discomfort after third molar surgery: a prospective study. *J Oral Maxillofac Surg* 2007;65:901-17.
 39. Baqain ZH, Karaky AA, Sawair F, Khraisat A, Duaibis R, Rajab LD. Frequency estimates and risk factors for postoperative morbidity after third molar removal: a prospective cohort study. *J Oral Maxillofac Surg* 2008;66:2276-83.
 40. Seymour RA, Meechan JG, Blair GS. An investigation into postoperative pain after third molar surgery under local analgesia. *Br J Oral Maxillofac Surg* 1985;23:410-8.
 41. de Santana Santos T, Calazans AC, Martins-Filho PR, Silva LC, de Oliveira E Silva ED, Gomes AC. Evaluation of the muscle relaxant cyclobenzaprine after third-molar extraction. *J Am Dent Assoc* 2011;142:1154-62.
 42. Akadir OA, Fasola AO, Arotiba JT. Evaluation of Pederson index as an instrument for predicting difficulty of third molar surgical extraction. *Niger Postgrad Med J* 2009;16:105-8.
 43. Pedersen GW. *Oral surgery*. Philadelphia: Saunders; 1988.
 44. Renton T, Smeeton N, McGurk M. Factors predictive of difficulty of mandibular third molar surgery. *Br Dent J* 2001;190:607-10.
 45. Yuasa H, Kawai T, Sugiura M. Classification of surgical difficulty in extracting impacted third molars. *Br J Oral Maxillofac Surg* 2002;40:26-31.
 46. Carvalho RW, do Egito Vasconcelos BC. Assessment of factors associated with surgical difficulty during removal of impacted lower third molars. *J Oral Maxillofac Surg* 2011;69:2714-21.
 47. Conrad SM, Blakey GH, Shugars DA, Marciani RD, Phillips C, White RP Jr. Patients' perception of recovery after third molar surgery. *J Oral Maxillofac Surg* 1999;57:1288-94.
 48. Larsen PE. Alveolar osteitis after surgical removal of impacted mandibular third molars. Identification of the patient at risk. *Oral Surg Oral Med Oral Pathol* 1992;73:393-7.
 49. Meechan JG, Macgregor ID, Rogers SN, Hobson RS, Bate JP, Dennison M. The effect of smoking on immediate post-extraction socket filling with blood and on the incidence of painful socket. *Br J Oral Maxillofac Surg* 1988;26:402-9.
 50. Johnson WS, Blanton EE. An evaluation of 9-aminoacridine/Gelfoam to reduce dry socket formation. *Oral Surg Oral Med Oral Pathol* 1988;66:167-70.
 51. Yoshii T, Hamamoto Y, Muraoka S, Furudoi S, Komori T. Differences in postoperative morbidity rates, including infection and dry socket, and differences in the healing process after mandibular third molar surgery in patients receiving 1-day or 3-day prophylaxis with lenampicillin. *J Infect Chemother* 2002;8:87-93.
 52. Alexander RE. Dental extraction wound management: a case against medicating postextraction sockets. *J Oral Maxillofac Surg* 2000;58:538-51.
 53. Brekke JH, Bresner M, Reitman MJ. Effect of surgical trauma and polylactate cubes and granules on the incidence of alveolar osteitis in mandibular third molar extraction wounds. *J Can Dent Assoc* 1986;52:315-9.
 54. Adkisson SR, Harris PF. A statistical study of alveolar osteitis. *U S Armed Forces Med J* 1956;7:1749-54.
 55. Berge TI, Gilhuus-Moe OT. Per- and post-operative variables of mandibular third-molar surgery by four general practitioners and one oral surgeon. *Acta Odontol Scand* 1993;51:389-97.
 56. Sisk AL, Hammer WB, Shelton DW, Joy ED Jr. Complications following removal of impacted third molars: the role of the experience of the surgeon. *J Oral Maxillofac Surg* 1986;44:855-9.
 57. Eshghpour M, Moradi A, Nejat AH. Dry socket following tooth extraction in an Iranian Dental Center: incidence and risk factors. *J Dent Mater Tech* 2013;2:86-91.
 58. Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL 3rd. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain* 2009;10:447-85.
 59. Gerdle B, Björk J, Cöster L, Henriksson K, Henriksson C, Bengtsson A. Prevalence of widespread pain and associations with work status: a population study. *BMC Musculoskelet Disord* 2008;9:102.
 60. Wijnhoven HA, de Vet HC, Picavet HS. Prevalence of musculoskeletal disorders is systematically higher in women than in men. *Clin J Pain* 2006;22:717-24.
 61. de Mos M, de Bruijn AG, Huygen FJ, Dieleman JP, Stricker BH, Sturkenboom MC. The incidence of complex regional pain syndrome: a population-based study. *Pain* 2007;129:12-20.
 62. Tsang A, Von Korff M, Lee S, Alonso J, Karam E, Angermeyer MC, et al. Common chronic pain conditions in developed and developing countries: gender and age differences and comorbidity with depression-anxiety disorders. *J Pain* 2008;9:883-91.
 63. LeResche L. Gender considerations in the epidemiology of chronic pain. *Epidemiol Pain* 1999;17:43-52.
 64. Bastos JL, Gigante DP, Peres KG. Toothache prevalence and associated factors: a population based study in southern Brazil. *Oral Dis* 2008;14:320-6.
 65. Chia YY, Chow LH, Hung CC, Liu K, Ger LP, Wang PN. Gender and pain upon movement are associated with the requirements for postoperative patient-controlled iv analgesia: a prospective survey of 2,298 Chinese patients. *Can J Anaesth* 2002;49:249-55.
 66. Mattila K, Toivonen J, Janhunen L, Rosenberg PH, Hynynen M. Postdischarge symptoms after ambulatory surgery: first-week incidence, intensity, and risk factors. *Anesth Analg* 2005;101:1643-50.
 67. Taenzer AH, Clark C, Curry CS. Gender affects report of pain and function after arthroscopic anterior cruciate ligament reconstruction. *Anesthesiology* 2000;93:670-5.
 68. Rosseland LA, Stubhaug A. Gender is a confounding factor in pain trials: women report more pain than men after arthroscopic surgery. *Pain* 2004;112:248-53.
 69. Fillingim RB. Sex, gender, and pain: women and men really are different. *Curr Rev Pain* 2000;4:24-30.
 70. Keefe FJ, Lefebvre JC, Egert JR, Affleck G, Sullivan MJ, Caldwell DS. The relationship of gender to pain, pain behavior, and disability in osteoarthritis patients: the role of catastrophizing. *Pain* 2000;87:325-34.
 71. Nakagawa Y, Ishii H, Nomura Y, Watanabe NY, Hoshiba D, Kobayashi K, et al. Third molar position: reliability of panoramic radiography. *J Oral Maxillofac Surg* 2007;65:1303-8.
 72. Abu Younis MH, Abu Hantash RO. Dry socket: frequency, clinical picture, and risk factors in a palestinian dental teaching center. *Open Dent J* 2011;5:7-12.
 73. Barbosa-Rebellato NL, Thomé AC, Costa-Maciel C, Oliveira J, Scariot R. Factors associated with complications of removal of third molars: a transversal study. *Med Oral Patol Oral Cir Bucal* 2011;16:e376-80.
 74. Heasman PA, Jacobs DJ. A clinical investigation into the incidence of dry socket. *Br J Oral Maxillofac Surg* 1984;22:115-22.
 75. Garcia AG, Grana PM, Sampedro FG, Diago MP, Rey JM. Does oral contraceptive use affect the incidence of complications after extraction of a mandibular third molar? *Br Dent J* 2003;194:453-5.
 76. Catellani JE, Harvey S, Erickson SH, Cherkin D. Effect of oral contraceptive cycle on dry socket (localized alveolar osteitis). *J Am Dent Assoc* 1980;101:777-80.

77. Enkling N, Nicolay C, Bayer S, Mericske-Stern R, Utz KH. Investigating interocclusal perception in tactile teeth sensibility using symmetric and asymmetric analysis. *Clin Oral Investig* 2010;14:683-90.
78. Jacobs R, Van Steenberghe D. From osseoperception to implant-mediated sensory-motor interactions and related clinical implications. *J Oral Rehabil* 2006;33:282-92.
79. Kazemi M, Geramipناه F, Negahdari R, Rakhshan V. Active tactile sensibility of single-tooth implants versus natural dentition: a split-mouth double-blind randomized clinical trial. *Clin Implant Dent Relat Res* 2014;16:947-55.
80. Goldberg MH, Nemarich AN, Marco WP 2nd. Complications after mandibular third molar surgery: a statistical analysis of 500 consecutive procedures in private practice. *J Am Dent Assoc* 1985;111:277-9.
81. Bruce RA, Frederickson GC, Small GS. Age of patients and morbidity associated with mandibular third molar surgery. *J Am Dent Assoc* 1980;101:240-5.
82. Precious DS, Mercier P, Payette F. Risks and benefits of extraction of impacted third molars: a critical review of the literature. 2. *J Can Dent Assoc* 1992;58:845-52.
83. Akadiri OA, Okoje VN, Arotiba JT. Identification of risk factors for short-term morbidity in third molar surgery. *Odontostomatol Trop* 2008;31:5-10.
84. Adeyemo WL, Ogunlewe MO, Ladeinde AL, Hassan OO, Taiwo OA. A comparative study of surgical morbidity associated with mandibular third-molar surgery in young and aging populations. *J Contemp Dent Pract* 2010;11:E001-8.
85. Adeyemo WL, Ogunlewe MO, Ladeinde AL, Abib GT, Gbotolorun OM, Olojede OC, et al. Prevalence and surgical morbidity of impacted mandibular third molar removal in the aging population: a retrospective study at the Lagos University Teaching Hospital. *Afr J Med Med Sci* 2006;35:479-83.