



A retrospective study of mandibular fractures in children

Santanu Mukhopadhyay

Department of Pedodontics and Preventive Dentistry, Dr. R. Ahmed Dental College & Hospital, Kolkata, India

Abstract (J Korean Assoc Oral Maxillofac Surg 2018;44:269-274)

Objectives: The incidence and patterns of mandibular fractures vary by country and population age. This retrospective study evaluated the etiologies and patterns of mandibular fractures in children.

Materials and Methods: The clinical records of 89 children (45 males and 44 females) aged 0 to 12 years who presented with mandibular fractures from July 2012 to June 2016 were retrospectively reviewed. The sex, patient age, site of fracture, etiology of trauma, and monthly variations of the fractures were recorded. Descriptive statistics, the z-test and chi-square test were used for statistical analysis and the *P*-value was set at <0.05.

Results: Eighty-nine children (male-to-female ratio 1.02:1) sustained 131 mandibular fractures. Within the study sample, the 6 to 9 year age group most frequently experienced fractures (47.3%). Falls and road traffic accidents (RTA) were the two most common etiological factors that accounted for 44.9% and 24.7% of cases. The condylar fracture was the most common anatomical location (38.9%) followed by the angle (20.6%), parasymphysis (18.3%), body (15.3%), and symphysis (5.3%). A single fracture (51.7%) was more common than multiple fractures (48.3%). The month-wise distribution of mandibular fractures was fairly constant.

Conclusion: The condylar region was the most common anatomic site for fractures; in addition, a fall and RTA were the major etiological factors for mandibular fractures. A single fracture was observed in 51.7% of patients while multiple fracture lines accounted for 48.3% of cases.

Key words: Mandibular fractures, Child, Epidemiology

[paper submitted 2017. 12. 5 / revised 2018. 1. 17 / accepted 2018. 1. 17]

I. Introduction

Compared to adults, facial fractures are uncommon injuries in the pediatric population. The incidence is even rarer in children younger than five years of age. It is estimated that only 1% of facial fractures occur in toddlers and preschool-aged children. Pediatric and teenage maxillofacial trauma account for about 15% of all maxillofacial trauma¹⁻⁴.

Research investigation has shown that the mandible is the most commonly fractured facial bone³⁻⁹. Mandibular fractures in children may lead to serious complications like asymmet-

ric mandibular growth, temporomandibular joint ankylosis, and malocclusion. In general, soft tissue injuries are more common in children than skeletal fractures. Worldwide, falls from height, road traffic accidents (RTA), sports-related injuries, and bicycle accidents are the leading causes of mandibular fractures in the pediatric population. The incidence and patterns of mandibular fractures vary with geographical location, socioeconomic conditions, and the age of the population¹⁰. The majority of previous studies have focused on maxillofacial trauma in the adult population. These reports have shown that RTA is the most common mode of maxillofacial fractures in developing countries, while interpersonal violence accounts for the majority of such cases in developed countries^{6,7,9,11-13}. The purpose of this retrospective study was to evaluate the etiologies and patterns of mandibular fractures in children who reported to a tertiary care center in Kolkata, India between 2012 and 2016.

II. Materials and Methods

A retrospective study of mandibular fracture was performed in patients who presented with maxillofacial trauma

Santanu Mukhopadhyay

Department of Pedodontics and Preventive Dentistry, Dr. R. Ahmed Dental College & Hospital, 114 Acharya Jagadish Chandra Bose Road, Sealdah, Kolkata 700014, India

TEL: +91-33226511032 FAX: +91-33222268509

E-mail: msantanu25@gmail.com

ORCID: <https://orcid.org/0000-0002-7810-987X>

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2018 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.

to the Department of Pedodontics and Preventive Dentistry of the Dr. R. Ahmed Dental College & Hospital, Kolkata, from July 2012 to June 2016. This institute is affiliated with the West Bengal University of Health Sciences, Kolkata in India. The study was reviewed and approved by the ethical committee of the institute of Dr. R. Ahmed Dental College & Hospital. In this study, information was obtained from the injury report register and entered in a data collection sheet. We reviewed all patients aged 12 years and younger at the time of mandibular fractures. The subjects were divided into three groups: Group A (0-5 years), Group B (6-9 years), and Group C (10-12 years). Patients were categorized based on age, sex, place of residence, fracture mechanism, anatomic location of fracture and date of trauma. The trauma etiology was classified as fall from height, RTA, play- or sports-related injury, bicycle accident, assault, or other. The anatomical sites of the mandibular fractures were determined by panoramic and/or computed tomographic examination of the patients. Furthermore, mandibular fracture sites were classified according to Killey¹⁴: symphyseal, parasymphyseal, body, angle, ramus and condyle. Patient charts with incomplete information, extensive head injury and the presence of pathology were excluded from the study.

The results were tabulated and analyzed using the Epi Info (ver. 7; CDC-INFO, Atlanta, GA, USA). Descriptive analyses including frequency, percentage, and proportions were performed. Where appropriate, the significance of the findings was evaluated using the z-test and chi-square test. The level of significance was set at $P < 0.05$.

III. Results

1. Age and sex distribution

A total of 202 cases of maxillofacial trauma records were

extracted for review. Among them, 89 pediatric patients (44.1%) sustained mandibular fractures. Of the 89 patients, 45 patients (50.6%) were male and 44 patients (49.4%) were female with a male to female ratio of 1.02:1. The mean patient age (7.73 ± 3.02 years; 95% confidence interval [CI], 7.1-8.36) was slightly higher in females (8.20 ± 3.02 years; 95% CI, 7.31-9.1) than in males (7.26 ± 2.89 years; 95% CI, 6.39-8.14). However, this difference was not statistically significant (z for 95% CI=1.96). Most patients belonged to Group B (39 patients, 43.8%), followed by Group C (32 patients, 36.0%) and Group A (18 patients, 20.2%).(Table 1)

2. Place of residence

There were significantly ($P=0.0103$, Fisher exact test) more fracture cases (50 patients, 56.2%) reported from rural areas than from urban areas (39 patients, 43.8%) of residence.

3. Mandible fracture etiology

The causes of mandibular fractures are shown in Table 2. Most of the mandibular fractures were caused by falls (40 patients, 44.9%), and other causes included RTA (22 patients, 24.7%), play (16 patients, 18.0%), bicycle accident (8 patients, 9.0%), and violence (2 patients, 2.2%). Only one male

Table 1. Patient demographics

Age group	Male	Female	Total
Group A	11 (24.4)	7 (15.9)	18 (20.2)
Group B	20 (44.4)	19 (43.2)	39 (43.8)
Group C	14 (31.1)	18 (40.9)	32 (36.0)
Total	45 (50.6)	44 (49.4)	89 (100)

Values are presented as number (%).

Group A: 0-5 years, Group B: 6-9 years, Group C: 10-12 years.

Santanu Mukhopadhyay: A retrospective study of mandibular fractures in children. J Korean Assoc Oral Maxillofac Surg 2018

Table 2. Mandibular fracture etiologies by age group

Etiology	Group A		Group B		Group C		Total
	Male	Female	Male	Female	Male	Female	
Fall	5 (45.5)	5 (71.4)	11 (55.0)	10 (52.6)	3 (21.4)	6 (33.3)	40 (44.9)
Road traffic accidents	3 (27.3)	1 (14.3)	5 (25.0)	6 (31.6)	2 (14.3)	5 (27.8)	22 (24.7)
Play	2 (18.2)	1 (14.3)	3 (15.0)	1 (5.3)	6 (42.9)	3 (16.7)	16 (18.0)
Bicycle	1 (9.1)	0	1 (5.0)	2 (10.5)	1 (7.1)	3 (16.7)	8 (9.0)
Assault	0	0	0	0	1 (7.1)	1 (5.6)	2 (2.2)
Other	0	0	0	0	1 (7.1)	0	1 (1.1)
Total	11 (100)	7 (100)	20 (100)	19 (100)	14 (100)	18 (100)	89 (100)

Values are presented as number (%).

Group A: 0-5 years, Group B: 6-9 years, Group C: 10-12 years.

Santanu Mukhopadhyay: A retrospective study of mandibular fractures in children. J Korean Assoc Oral Maxillofac Surg 2018

Table 3. Fracture distribution by age group (n=89)

Site	Group A		Group B		Group C	
	Male	Female	Male	Female	Male	Female
Symphysis alone	0	0	0	1	0	1
Parasymphysis alone	2	0	0	0	0	0
Body alone	0	0	3	4	1	1
Angle alone	1	0	0	2	3	3
Condyle alone	4	4	4	2	2	7
Ramus alone	0	0	0	0	0	1
Symphysis+condyle	1	1	1	0	0	2
Parasymphysis+condyle	2	1	7	3	1	0
Body+angle	1	1	3	1	0	0
Body+condyle	0	1	2	1	1	0
Parasymphysis+angle	0	0	0	3	3	1
Angle+condyle	0	0	0	1	1	2
Symphysis+angle	0	0	0	0	1	0
Parasymphysis+ramus	0	0	0	0	1	0

Group A: 0-5 years, Group B: 6-9 years, Group C: 10-12 years.

Santanu Mukhopadhyay: A retrospective study of mandibular fractures in children. *J Korean Assoc Oral Maxillofac Surg* 2018

patient (1.1%) in Group C sustained a mandibular fracture due to trauma from a tube well handle.

4. Fracture distribution

The distribution of fractures is shown in Table 3. A total of 131 fracture locations were observed in 89 patients. The majority of fractures (70 patients, 53.4%) occurred in males. Approximately half of these fractures, (62 patients, 47.3%) were seen in Group B, followed by Group C (44 patients, 33.6%) and Group A (25 patients, 19.1%). There were 20 boys and 26 girls, a total of 46 patients (51.7%) that reported a single fracture location and 43 patients (48.3%), 25 boys and 18 girls presented with multiple fractures. However, no significant sex difference ($P=0.20$, Fisher exact test) was observed in the distribution of fractures. The most common site was the condylar region (38.9%) followed by angle (20.6%), parasymphysis (18.3%), body (15.3%), and symphysis (5.3%). Fracture of the ramus (1.5%) was observed only in Group C. There were no coronoid fractures observed in our study. Multiple fractures were most commonly observed in the condylar and parasymphyseal regions.

5. Month-wise distribution

With regard to the month-wise distribution of mandibular fractures, the maximum number of cases was reported in Au-

Table 4. Monthly distribution of mandibular fractures

Month	No. (%) of fractures
January	7 (7.9)
February	7 (7.9)
March	6 (6.7)
April	6 (6.7)
May	7 (7.9)
June	7 (7.9)
July	8 (9.0)
August	12 (13.5)
September	11 (12.4)
October	4 (4.5)
November	6 (6.7)
December	8 (9.0)
Total	89 (100)

Santanu Mukhopadhyay: A retrospective study of mandibular fractures in children. *J Korean Assoc Oral Maxillofac Surg* 2018

gust (12 patients, 13.5%) and the minimum number of cases was documented in October (4 patients, 4.5%).(Table 4)

IV. Discussion

Maxillofacial trauma is one of the principal causes of morbidity and mortality in children. Mandibular fractures at young ages may lead to functional impairment and disfigurement. Patterns and etiologies of mandibular fractures vary by geographical location, cultural characteristics, and socioeconomic status. The majority of investigations have shown that males are more prone to maxillofacial trauma than females in all age groups^{3-5,7,8,15,16}. Generally, the male to female ratio of maxillofacial trauma and mandibular fractures is 2:1. In contrast, the results of our study showed no significant sex bias. Cole et al.¹⁵ concluded that at a young age, the etiology of mandibular fractures is similar in both sexes. Therefore, sex-related fracture differences are less significant at a young age.

In this study, only 20.2% of fractures were reported in children below five years of age. This is consistent with the results of other investigations that also reported that maxillofacial trauma is uncommon in toddlers and preschool age children^{2,3,7,8,15}. This could be attributable to the fact that the younger age groups experience more parental supervision and less independence than older children. In addition, the retruded position of the face in relation to the skull, a greater cranium to face ratio in infants and toddlers, lack of pneumatization of nasal bones and greater flexibility of facial bones may contribute to a lower incidence of maxillofacial fractures in this age group. In our study, most injuries (39 patients, 43.8%) occurred in the 6 to 9 years age group. Several factors, like facial skeleton growth, participation in school ac-

tivities, and an increase in social interaction may increase the risk of maxillofacial trauma in this age group.

The results of this study support the finding of a study by Namdev et al.¹⁶, who also observed the majority of mandibular fracture cases in rural areas. Poor road conditions in some parts of rural India, especially during the rainy season, may have contributed to an increased incidence of maxillofacial trauma.

Falls from height (44.9%) were the major cause of mandibular fracture in this study. Our results are in agreement with the observations of Collao-González et al.⁵, Joshi et al.⁸, Namdev et al.¹⁶, Owusu et al.¹⁷, Kumaraswamy et al.¹⁸, and Atilgan et al.¹⁹, who also reported that falls from height were the most common mechanism of maxillofacial trauma in children. In this study, more than half of the population in Group A and Group B sustained mandibular fractures due to falls from height, representing low velocity/energy trauma. Kumaraswamy et al.¹⁸ observed that in children up to 6 years of age, falls in the home were the most common mode of maxillofacial trauma, and with increasing age and more outdoor exposure, falls tended to occur outside the home.

This study demonstrated that RTA was the second most common cause of mandibular fracture. Some reports have shown that RTA is the main etiology of trauma^{3,7,20}. A recent Indian study by Singhal et al.⁷ also reported that RTA was the most common mode of maxillofacial trauma (57.27%) in children and adolescents. In this report, motor vehicle as well as motorbike accidents were categorized as RTA, and accounted for approximately 25% of fracture cases. In addition, 9.0% of mandibular fractures occurred secondary to bicycle accidents. One reason for mandibular fractures due to RTA might be the lack of helmet use by children while traveling in motorbikes. Furthermore, the use of seat belts and airbags as safety measures is limited, especially in semi-urban and rural parts of India where traffic rules are not strictly enforced.

In the present study, play- or sports-related injuries increased with age and the highest incidence was observed in the 10 to 12 years age group. Muraoka et al.²¹ also reported similar trends among Japanese children. A rise in fracture incidence during puberty and adolescence may be due to increased involvement in sports.

Schlievert²² and Knoche et al.²³ reported child abuse to be one cause of mandibular fractures in children. In this study, we did not identify child abuse as a mechanism for mandibular fractures, which was similar to the pattern observed by Namdev et al.¹⁶ and Fasola et al.²⁴. In comparison, Almahdi and Higzi³, Gassner et al.¹⁰, Atilgan et al.¹⁹, and Tanaka et al.²⁵

found that interpersonal violence is an important etiology for pediatric facial fractures. Additionally, Bamjee²⁶ reported assaults and gunshot wounds to be the cause of mandibular fracture in 48% of patients. In contrast, only two cases of mandibular fractures due to assaults or interpersonal violence were observed in this study. The higher incidence of mandibular fractures caused by assaults in previous studies may be attributed to differences in socioeconomic factors, behavioral habits, and the inclusion of adolescents in the study population.

In this study, the condylar region was the most common site of mandibular fracture. Our results are in agreement with the observations of other studies reporting the condyle as the most common fracture site of the mandible. In preschool children, condylar fractures accounted for the majority of cases (52%) and the incidence decreases with increasing age. The condylar process in children has high bone marrow content and the cortex is relatively thin. Therefore, this causes low resistance to low velocity trauma during fall. Overall, condylar fractures in the present study accounted for 39% of all mandibular fractures. This is comparable with the results of Almahdi and Higzi³ (29.8%), Joshi et al.⁸ (40.9%), Namdev et al.¹⁶ (40.3%), Owusu et al.¹⁷ (27.9%), and Shi et al.²⁷ (55.7%). In addition, approximately half of patients (48.3%) presented with fractures at more than one site. Multiple fractures most commonly involve the parasymphiseal and condylar regions. Patients with RTA as the main etiology for mandible fractures have reported the parasymphysis and condyle to be common fracture sites⁹.

Haug and Foss² concluded that mandibular angle fractures are rare in children and adolescents. Shi et al.²⁷ reported mandibular angle fractures in only 3.5% of cases, which primarily occurred in patients with permanent dentition. In our study, the mandibular angle was the second most common fracture site and the incidence of fracture to this area increased with age. This finding is supported by the observations of Almahdi and Higzi³, Owusu et al.¹⁷, and Thorén et al.²⁸, who also reported that mandibular angle fractures consistently increased with age. In addition, the overall incidence of fractures of the parasymphysis (18.3%) and body (15.3%) were comparable with the results of Muñante-Cárdenas et al.⁴ who observed similar incidences of fractures of the parasymphysis (18.7%) and body of the mandible (15.1%).

In this study, the incidence of mandibular fractures was highest in August, although it was relatively constant with seasonal variations. This is in agreement with the results of Joshi et al.⁸, Haug and Foss², and Tanaka et al.²⁵ who also ob-

served that month-wise distribution of mandibular fractures remains fairly constant with seasonal variations. Jung et al.²⁹ reported that the highest incidences of fractures occurred during the months of autumn and that the January was the month with the lowest reported incidence. In contrast to our findings, some studies have reported that the monthly distribution of maxillofacial fractures peaked in summer^{30,31}.

The limitations of the present study include the retrospective nature and small sample size. In spite of these limitations, our results show the epidemiology and patterns of mandibular fractures in the eastern parts of India. We hope that the results of our study will provide insight for healthcare providers to formulate preventive strategies.

V. Conclusion

The results of this study indicated that there was no significant difference in the incidence of mandibular fractures according to sex. Another distinguishing feature of this study was a high incidence of fractures in rural populations. Our finding that condylar fractures were the most common type of fractures and that falls are the primary causes of mandibular fractures was supported by the majority of published data. In addition, RTA and bicycle accidents account for approximately one-third of cases, indicating the need for strict implementation of traffic rules.

Author's Contributions

S.M. has contributed to the design of the work, acquisition, and analysis of data, drafting of the work and manuscript preparation.

Ethics Approval and Consent to Participate

The study was reviewed and approved by the Ethical Committee of the institute of Dr. R. Ahmed Dental College & Hospital, but the Ethical Committee does not provide an approval number.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

References

1. Marano R, de Oliveira Neto P, Sakugawa KO, Zanetti LSS, de Moraes Z. Mandibular fractures in children less than 3 years: a rare case report. *Rev Port Estomatol Med Dent Cir Maxilofac* 2013;54:115-76.
2. Haug RH, Foss J. Maxillofacial injuries in pediatric patient. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000;90:126-34.
3. Almahdi HM, Higzi MA. Maxillofacial fractures among Sudanese children at Khartoum Dental Teaching Hospital. *BMC Res Notes* 2016;9:120.
4. Muñante-Cárdenas JL, Olate S, Asprino L, de Albergaria Barbosa JR, de Moraes M, Moreira RW. Pattern and treatment of facial trauma in pediatric and adolescent patients. *J Craniofac Surg* 2011;22:1251-5.
5. Collao-González C, Carrasco-Labra A, Sung-Hsieh HH, Cortés-Araya J. Epidemiology of pediatric facial trauma in Chile: a retrospective study of 7,617 cases in 3 years. *Med Oral Patol Oral Cir Bucal* 2014;19:e99-105.
6. Pandey S, Roychoudhury A, Bhutia O, Singhal M, Sagar S, Pandey RM. Study of the pattern of maxillofacial fractures seen at a tertiary care hospital in North India. *J Maxillofac Oral Surg* 2015;14:32-9.
7. Singhal R, Singh V, Bhagol A, Agrawal A, Kumar P. Pediatric maxillofacial injuries - if a new look is required? *Int J Pediatr Otorhinolaryngol* 2013;77:1333-6.
8. Joshi SR, Saluja H, Pendyala GS, Chaudhari S, Mahindra U, Kini Y. Pattern and prevalence of maxillofacial fractures in rural children of central Maharashtra, India. A retrospective study. *J Maxillofac Oral Surg* 2013;12:307-11.
9. Kumar GB, Dhupar V, Akkara F, Kumar SP. Patterns of maxillofacial fractures in goa. *J Maxillofac Oral Surg* 2015;14:138-41.
10. Gassner R, Tuli T, Hächl O, Rudisch A, Ulmer H. Cranio-maxillofacial trauma: a 10 year review of 9,543 cases with 21,067 injuries. *J Craniomaxillofac Surg* 2003;31:51-61.
11. Abdullah WA, Al-Mutairi K, Al-Ali Y, Al-Soghier A, Al-Shnwni A. Patterns and etiology of maxillofacial fractures in Riyadh city, Saudi Arabia. *Saudi Dent J* 2013;25:33-8.
12. Ferreira P, Marques M, Pinho C, Rodrigues J, Reis J, Amarante J. Midfacial fractures in children and adolescents: a review of 492 cases. *Br J Oral Maxillofac Surg* 2004;42:501-5.
13. Verma S, Chambers I. Update on patterns of mandibular fracture in Tasmania, Australia. *Br J Oral Maxillofac Surg* 2015;53:74-7.
14. Killey HC. Killey's fractures. In: Banks P, ed. *Killey's fractures of the mandible*. 3rd ed. London: Wright; 1988:15-28.
15. Cole P, Kaufman Y, Hollier LH Jr. Managing the pediatric facial fracture. *Cranio-maxillofac Trauma Reconstr* 2009;2:77-83.
16. Namdev R, Jindal A, Bhargava S, Dutta S, Singhal P, Grewal P. Patterns of mandible fracture in children under 12 years in a district trauma center in India. *Dent Traumatol* 2016;32:32-6.
17. Owusu JA, Bellile E, Moyer JS, Sidman JD. Patterns of pediatric mandible fractures in the United States. *JAMA Facial Plast Surg* 2016;18:37-41.
18. Kumaraswamy SV, Madan N, Keerthi R, Singh DS. Pediatric injuries in maxillofacial trauma: a 5 year study. *J Maxillofac Oral Surg* 2009;8:150-3.
19. Atilgan S, Erol B, Yaman F, Yilmaz N, Ucan MC. Mandibular fractures: a comparative analysis between young and adult patients in the southeast region of Turkey. *J Appl Oral Sci* 2010;18:17-22.
20. Holland AJ, Broome C, Steinberg A, Cass DT. Facial fractures in children. *Pediatr Emerg Care* 2001;17:157-60.
21. Muraoka M, Nakai Y, Nakagawa K, Yoshioka N, Nakaki Y, Yabe T, et al. Fifteen-year statistics and observation of facial bone fracture. *Osaka City Med J* 1995;41:49-61.
22. Schlievert R. Infant mandibular fractures: are you considering child abuse? *Pediatr Emerg Care* 2006;22:181-3.
23. Knoche JW, LeBlanc KK, King TW, Knox BL. An infant with a

- unilateral mandibular fracture: when to consider nonaccidental trauma. *Clin Pediatr (Phila)* 2012;51:404-7.
24. Fasola AO, Obiechina AE, Arotiba JT. Fractures of the mandible in children. *East Afr Med J* 2001;78:616-8.
 25. Tanaka N, Uchida N, Suzuki K, Tashiro T, Tomitsuka K, Kimijima Y, et al. Maxillofacial fractures in children. *J Craniomaxillofac Surg* 1993;21:289-93.
 26. Bamjee Y. Paediatric maxillofacial trauma. *J Dent Assoc S Afr* 1996;51:750-3.
 27. Shi J, Chen Z, Xu B. Causes and treatment of mandibular and condylar fractures in children and adolescents: a review of 104 cases. *JAMA Otolaryngol Head Neck Surg* 2014;140:203-7.
 28. Thorén H, Iso-Kungas P, Iizuka T, Lindqvist C, Törnwall J. Changing trends in causes and patterns of facial fractures in children. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:318-24.
 29. Jung HW, Lee BS, Kwon YD, Choi BJ, Lee JW, Lee HW, et al. Retrospective clinical study of mandible fractures. *J Korean Assoc Oral Maxillofac Surg* 2014;40:21-6.
 30. Posnick JC, Wells M, Pron GE. Pediatric facial fractures: evolving patterns of treatment. *J Oral Maxillofac Surg* 1993;51:836-44; discussion 844-5.
 31. Burlini D, Conti G, Amadori F, Bardellini E, De Giuli C. Management of paediatric maxillofacial fractures: conventional methods and resorbable materials. *Eur J Paediatr Dent* 2015;16:24-8.