

한국심리학회지

발달

33권 1호 (2020년 3월)



THE KOREAN JOURNAL OF DEVELOPMENTAL PSYCHOLOGY

목 차

교사-아동 관계가 학업수행적응에 미치는 영향: 아동의 자아존중감 및 친사회적 행동을 매개변수로	김진성
청소년이 닮고 싶어 하는 또래는 누구인가? 또래 지위 지표로서의 선망도 탐색	신희영
학교 밖 청소년의 심리·사회적 발달궤적과 성인초기 적응 여부에 따른 차이	최지은, 이지원, 정다빈, 이해나, 김현경
한국 아동의 선택적 신뢰 발달: 5세 아동이 4세와 달리 정보제공자의 연령을 정확도보다 중시할 때	범지민, 최영은
대학생의 분배 및 절차공정성에 대한 개인적 믿음, 미래지향 시간관과 정신적 안녕감의 관계	손연경, 이승연
자폐스펙트럼장애에서 발달 기능을 위한 가상현실-기반 중재: 메타분석	홍승주, 김미성
그릿(Grit), 학업적 실패내성, 성취목표지향성에 대한 잠재프로파일분석	안태영, 양수진

발행처 : 한국발달심리학회
발행인 : 송하나(성균관대학교)
주 소 : 서울특별시 마포구 월드컵북로2길 93 진빌딩 2층
전 화 : 02-336-6672

인쇄일 : 2020년 3월 15일
발행일 : 2020년 3월 15일
제작처 : 책과공간
(02-725-9371)

편집위원장 : 이승연(이화여자대학교)

편집위원 : 강연욱(한림대학교) 김기연(중앙대학교) 김혜리(충북대학교) 노수림(충남대학교)
서미정(제주대학교) 송주현(California State University) 송현주(연세대학교) 신희영(전북대학교)
유연옥(계명대학교) 최유정(Harvard University) 한세영(이화여자대학교)
Daniel Choe(UC Irvine Human Development and Family Studies)

심사위원 : 김근영(서강대학교) 김기연(중앙대학교) 김민희(한국상담대학원대학교) 김수안(서울대학교)
김수영(이화여자대학교) 김혜리(충북대학교) 맹세호(성신여자대학교) 박다은(충북대학교)
서미정(제주대학교) 설경옥(이화여자대학교) 성은현(호서대학교) 손정락(전북대학교)
송경희(University of Toronto, Mississauga) 송주현(California State University)
심희옥(군산대학교) 양수진(이화여자대학교) 오인수(이화여자대학교) 유연옥(계명대학교)
윤 황(경기대학교) 이승연(이화여자대학교) 이승진(건국대학교) 이주영(동덕여자대학교)
정은정(명지대학교) 최미경(덕성여자대학교) 최유정(Harvard University) 한세영(이화여자대학교)

한국심리학회지: 발달은 한국발달심리학회의 기관지로서 연 4회 간행되며, 발달심리학 분야의 연구논문, 자료 및 논평을 게재한다. 한국심리학회지: 발달은 일정한 구독료를 받고 배부하며, 한국심리학회에 연회비를 납부한 회원은 한국심리학회 홈페이지에서 온라인 구독이 가능하다. 비회원의 구독에 관해서는 편집위원회 사무실로 문의하기 바란다.

Vol. 33, No. 1

March 15, 2020

THE KOREAN JOURNAL OF DEVELOPMENTAL PSYCHOLOGY
published quarterly-annually
by THE KOREAN PSYCHOLOGICAL ASSOCIATION

This journal is issued quarterly-annually and carries research articles based on empirical data & theoretical review. Subscription inquiries and manuscript submission should be directed to: Editor, The Korean Journal of Developmental Psychology, Department of Psychology, Ewha Womans University, Seoul, Korea.

Editor : Seung-yeon Lee(Ewha Womans University)

Associate Editor : Yeonwook Kang(Hallym Univeristy) Giyeon Kim(Chung-Ang University)
Hei-Rhee Ghim(Chungbuk National University) Soo Rim Noh(Chungnam National University)
Mijung Seo(Jeju National University) Ju-Hyun Song(California State University)
Hyun-joo Song(Yonsei University) Huiyoung Shin(Chonbuk National University)
Younoak Yu(Keimyung University) Youjung Choi(Harvard University)
Sae-Young Han(Ewha Womans University)
Daniel Choe(UC Irvine Human Development and Family Studies)

한국심리학회지

발 달

제 33 권 제 1 호 / 2020. 3.

교사-아동 관계가 학업수행적응에 미치는 영향: 아동의 자아존중감 및 친사회적 행동을 매개변수로	김진성 / 1
청소년이 닮고 싶어 하는 또래는 누구인가? 또래 지위 지표로서의 선망도 탐색	신희영 / 19
학교 밖 청소년의 심리·사회적 발달궤적과 성인초기 적응 여부에 따른 차이	최지은, 이지원, 정다빈, 이해나, 김현경 / 37
한국 아동의 선택적 신뢰 발달: 5세 아동이 4세와 달리 정보제공자의 연령을 정확도보다 중시할 때	범지민, 최영은 / 53
대학생의 분배 및 절차공정성에 대한 개인적 믿음, 미래지향 시간관과 정신적 안녕감의 관계	손연경, 이승연 / 65
Virtual Reality-Based Intervention for Developmental Functioning in ASD: A Meta-Analysis	Seungju Hong, Hillary Mi-Sung Kim / 85
그릿(Grit), 학업적 실패내성, 성취목표지향성에 대한 잠재프로파일분석	안태영, 양수진 / 103

한국발달심리학회

Virtual Reality-Based Intervention for Developmental Functioning in ASD: A Meta-Analysis

Seungju Hong¹, Hillary Mi-Sung Kim²

Department of Child Psychology and Education, Sungkyunkwan University/ Master's student¹
Department of Child Psychology and Education, Sungkyunkwan University/ Lecturer²

Received: January 16, 2020
Revised: February 06, 2020
Accepted: February 27, 2020

교신저자: 김미성
성균관대학교 아동청소년학과
서울시 종로구 성균관로 25-2

E-MAIL:
hmisungk1023@gmail.com

자폐스펙트럼장애에서 발달 기능을 위한 가상현실-기반 중재: 메타분석

홍승주¹, 김미성²
성균관대학교 아동청소년학과/ 석사과정¹
성균관대학교 아동청소년학과/ 강사²

© Copyright 2019. The
Korean Journal of
Developmental Psychology.
All Rights Reserved.

*이 논문은 2019년 자폐학회
추계학술대회에서 포스터 발표한
내용을 수정 보완한 것임

ABSTRACT

This meta-analysis aggregated and analyzed the research in studying the pretest-posttest outcomes of VR-based interventions for children and young adults with ASD. Seven studies published between 2009 and 2019 with a total of 150 participants were included. Outcome measures of social competences were classified into 5 categories. The random-effects model of the current meta-analysis showed that the average effect of VR-based intervention as follows: social skills($g = .59, SE = .12, 95\% CI: .35-.82$), emotion recognition($g = .50, SE = .16, 95\% CI: 0.19-.82$), attention($g = .35, SE = .14, 95\% CI: .07-.63$), executive function($g = .52, SE = .12, 95\% CI: .28-.77$) and conversational skills($g = .46, SE = .21, 95\% CI: .05-.88$). The implication for VR-based interventions for ASD was discussed with respect to environmental factor such as age, ASD functioning, number of sessions, country where the study conducted and additional treatment combined.

Keywords: ASD, virtual reality, social skills, attention, emotion recognition, executive function, conversational skills

According to the Korean Ministry of Health and Welfare's 「Fact-finding Survey of Persons with Disabilities」 in 2017, the prevalence rate of autism spectrum disorder is gradually increasing to 0.01 percent in 2011, 0.02 percent in 2014 and 0.03 percent in 2017. According to the Health and Welfare Ministry's 「Status of Registration as Persons with Disabilities」 in 2018, the number of children with disabilities (ages below 17) is 72,260 and among them, about 17.4 percent, or 12,559 children present with autism spectrum disorder(ASD). With the increasing number of children with ASD, the development and implementation of various interventions are being studied at home and abroad (Lee, 2018). To implement effective interventions for children with ASD, it is crucial to evaluate newly introduced interventions.

The fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5; American Psychiatric Association, 2013) made changes on autism diagnosis. The diagnosis is called ASD, which no longer distinguishes subdiagnoses (Autistic Disorder, Asperger Syndrome, Pervasive Developmental Disorder Not Otherwise Specified, Disintegrative Disorder) that were in the DSM-IV. Children with ASD exhibit delay or impairments in social behavior and communication including social orienting, joint attention, emotion recognition and

expression, verbal and non-verbal communication (Wenar & Kerig, 2000). Associated features including abnormal responses to sensory stimuli, deficits in theory of mind, central coherence, and executive functions are often seen as well (Wicks-Nelson, Israel, 2013).

The interest in the utility of virtual reality(VR) as a treatment tool has been relatively recently increased, and VR-based interventions have been applied for a wide range of disorders such as PTSD, OCD, phobias, as well as ASD (Mesa-Gresa et al., 2018). Especially, VR-based interventions showed advantages for people with ASD because of the ecological validity of the intervention, in which the environment could be manipulated to suit the characteristics of the individuals (Jarrold et al. 2013). For example, VR interventions were developed to involve the participants with ASD in life-like contexts such as classrooms, peer groups, and meeting new people. The intervention was provided with a computer set up, a half-CAVE design, or a 3D VR technology. By setting up multiple social environment and partners, the possibility of generalization of the treatment into a real life context can be increased. Meanwhile, VR technology to improve for VR based interventions was noted by researchers. This, for example, includes difficulties in displaying immediate facial emotions and dizziness caused by a

lack of reality. It may also be a limitation of VR-based interventions that clinicians and caregivers need to be trained in order to use the technology appropriately.

Previous researches on virtual reality in technology, rehabilitation and mental health (Mesa-Gresa et al., 2018; Miller & Bugnariu, 2016; Wass & Porayska-Pomsta, 2014) contributed to reveal the usefulness of VR as a diagnostic tool. However, to our knowledge, since VR-based interventions have been introduced, their effectiveness as an intervention tool for autism spectrum disorders has not been analyzed. The current study aimed to synthesize previous studies of VR as an intervention tool for ASD.

Method

Study selection

Studies of VR based interventions administered to children and young adults with ASD were selected for the analysis. For the data, following databases are searched: ERIC, PubMed, Cochrane, psycINFO and EBSCOhost from 2009 to 2019. The searches included the terms “virtual reality” or “virtual” in combination with “autism”, “asd”, or “aspergers”. Review papers were searched in order to find any missing data as well.

Subjects were limited to the children and young adults. Studies meeting the following criteria were included: 1) at least one VR-based intervention, 2) pre and post treatment tests and 3) subjects with ASD. A total of 7 studies with a total sample size of 150 participants (mean age = 14.29 year) met the final inclusion criteria and were included.

The following studies were excluded according to the following reasons: studies that did not use VR as an intervention tool (Bystrova et al., 2019; Cai et al., 2017; Chia & Kee, 2014; Lahiri et al., 2011) studies with post-test only (Jarrold et al., 2013; Ke & Moon, 2018; Kuriakose & Lahiri, 2015; Wallace et al., 2017; Wallace et al., 2010; Wang & Reid, 2013; Wang et al., 2016); studies that did not evaluate disorder related skills or symptoms (Zhao et al., 2018); studies that used a single subject (Cheng et al., 2015; De Luca et al., 2019; Stendal & Balandin, 2015) and studies in which insufficient statistical data were available (Beach & Wendt, 2015; Lahiri et al., 2012; Lorenzo et al., 2016; Millen et al., 2014; Moon & Ke, 2019; Politis et al., 2017). See Fig. 1 for a diagram of the study selection process.

All analyses were completed with the Comprehensive Meta-Analysis software version 3.




Figure. 1 Study Selection Process

Procedure

In addition to outcome data, the following

data were collected: characteristics of the study participants (age, diagnosis), attrition information, treatment condition assignment

Table 1. Outcome measures

Outcome	Measure
Social Skills	Triangles, PEP-3 social reciprocity
Emotion recognition	NEPSY-II Affect recognition, Faces test, Eyes test, DANVA
Attention	NEPSY-II Auditory Attention, WebNeuro software-attention, Conners' CPT
Executive Function	Analogical reasoning task, WebNeuro software-executive function, D-KEFS
Conversational Skills	ATEC(speech, language, communication), ABAS Communication, SSPA, WebNeuro software-verbal

ABAS: Adaptive Behavior Assessment System; ATEC: Autism Treatment Evaluation Checklist; CPT: Continuous Performance Test; DANVA: Diagnostic Analysis of Non-Verbal Accuracy; D-KEFS: Dells-Kaplan Executive Functioning system; NEPSY: a developmental NEUROPSYchological Assessment; PEP: Psychoeducational Profile; SSPA: Social Skills Performance Assessment

methods (random/matched), amount of treatment (number and length of sessions), number of participants per condition, and year of publication, qualities of outcome measures (validity, reliability and rater). Outcome measures were classified into such categories as social skills, emotion recognition, attention, executive function and conversational skills (see Table 1). Analyses were conducted using a combined score if more than one measure is used for a category.

Effect Size Calculation

The effect sizes were computed using Hedge's *g* for each study. Hedge's *g* is used in order to fix biased effect size when Cohen's *d* is used for small sample size. Hedge's *g* may be interpreted similarly to

Cohen's *d* (Cohen, 1988), with a cutoff for small (0.2), medium (0.5) and large (0.8) effects. Also, inverse variance weight is used which provides relatively stable result compare to other weighted mean effect size. The effect sizes were calculated using this

$$\text{formula: Hedge's } g = J \times \frac{(\bar{x}_1 - \bar{x}_2)}{s_{pooled}} = J \times d,$$

where \bar{x}_1 is the mean of the posttest score, \bar{x}_2 is the mean of pretest score, and S_p is the pooled standard deviation. *J* is the correction constant, $J = 1 - \frac{3}{4(n_1 + n_2 - 2) - 1}$. A

combined effect size was computed for studies using multiple outcome measures. The overall mean effect size for studies combined was computed using a weighted

$$\text{formula, } \overline{ES^*} = \frac{\sum_{i=1}^n ES_i w_i^*}{\sum_{i=1}^n w_i^*}, \text{ where } w_i^* \text{ is the}$$

Table 2. Study Characteristics

Study	Location	Mean age of sample (range)	n	Sample characteristics	VR type	Additional treatment	Design	Session Design (# of people)	No. of sessions (total time)	Outcome measured time	Primary outcome measures	Primary outcome Hedge's g	Additional outcome measures
Amaral et al. (2018)	Portugal	22 yr (16-38 yr)	15	High functioning ASD	Brain Computer Interface (VR)	None	One-group pretest-posttest	Group (2)	7 (N/R)	Last session	ATEC- speech, Language, communication	0.925	JAAT, ATEC; VABS; POMS; HADS; BDI
Didehbani et al. (2016)	US	11.4 yr (7-16 yr)	30	ASD	VR-SCT	None	One-group pretest-posttest	Group (2)	10 (10hr)	Within 2 weeks of the last session	NEPSY-II Affect Recognition Triangles NEPSY-II auditory attention Ekman 60 Analogical Reasoning Task	0.725 0.389 0.210 0.558 0.455	NEPSY-II response set
Ip et al. (2018)	Hong Kong	9.03 yr (7-10 yr)	36	ASD	Cave Automatic Virtual Environment	None	Experimental group pretest-posttest (waitlist control)	Group (3-4)	28 (N/R)	Within 2 weeks of the last session	Face test Eyes test	0.204 0.227	PEP-3; ABAS-II
Kandalaft et al. (2012)	US	21.25 yr (8-26 yr)	8	Aspergers or PDD-NOS	VR-SCT	None	One-group pretest-posttest	Individual	10 (10hr)	Within 2 weeks of the last session	Elkman 60 Eyes test Triangles SSPA	1.266 0.319 1.078 0.731	ACS-SP
Lamash et al. (2017)	Israel	14 yr (11-19 yr)	33	ASD	VAP-S Virtual Environment	Meta-cognition strategies	One-group pretest-posttest	N/R	8 (N/R)	N/R	WebNeuro software-attention WebNeuro software-executive function WebNeuro software-verbal	0.576 0.585 0.377	Webneuro software; BRIEF-SR; TOGSS
Stichter et al. (2013)	US	12 yr (11-14 yr)	11	ASD	iSocial-3D Virtual Learning Environment	None	Experimental group pretest-posttest (alternating programs)	Individual	5 (31-45 min each)	2 weeks after the last session	DANVA D-KEFS Conners CPT-II commission Conners CPT-II omission	0.440 0.407 0.153 0.089	Eyes test: The Faux Pas Stories; The Strange Stories
Yang et al. (2018)	US	22.5 yr (18-31 yr)	17	High functioning ASD	VR-SCT	None	One-group pretest-posttest	Individual	N/R (10hr)	N/R	Triangles	0.460	ACS-CP

weight for each study, $w_i^* = \frac{1}{v_i + r^2}$. The random effects analysis was used, which assumes that the studies included are only a sample of the entire population of studies (Hwang, 2016, Chin, 2015).

Result

Seven studies published between 2009 and 2019 met criteria for inclusion with a total of 150 participants. Social skills were measured in 4 studies, and emotion recognition was measured in 4 studies, attention was measured in 3 studies, executive function was measured in 3 studies and conversational skills were measured in 4 studies. Both pre-test and post-test was demonstrated in all 7 studies, while only one study had both experimental and comparison groups. Interventions ranged in number from 8 to 28 sessions ($n = 89$, $Mdn = 14.8$ sessions). Hours of intervention ranged from 0.5 to 28 hours ($n = 155$, $Mdn = 51.6$ hours). Length of intervention ranged from 20 to 120 days ($n = 337$, $Mdn = 56.1$ days). Study characteristics are presented in Table 2.

The pooled effect size of VR-based interventions for 4 studies in social skills yielded a medium effect size ($g = .59$, $SE = .12$, 95% $CI: .35 - .82$, $p < .001$) (Fig. 2-1).

The pooled effect size VR-based

interventions for 4 studies in emotion recognition yielded a medium effect size ($g = .50$, $SE = .16$, 95% $CI: .19 - .82$, $p = .002$) (Fig. 2-2).

The pooled effect size VR-based interventions for 3 studies in attention yielded a small effect size ($g = .35$, $SE = .14$, 95% $CI: .07 - .63$, $p = .015$) (Fig. 2-3).

The pooled effect size VR-based interventions for 3 studies in executive function yielded a medium effect size ($g = .52$, $SE = .12$, 95% $CI: .28 - .77$, $p < .001$) (Fig. 2-4).

The pooled effect size of VR-based interventions for 4 studies in conversational skills yielded a medium effect size ($g = .46$, $SE = .21$, 95% $CI: .05 - .88$, $p = .027$) (Fig. 2-5).

Publication bias

Publication bias in meta-analysis refers that published studies do not represent all the results. There is possibility that studies with positive and significant results are tend to be published more often than those with negative, non-significant results (Dickerson, 2005). It may lead a meta-analysis to overestimate the effect size of the studies. In order to detect publication bias, we used a funnel plot asymmetry test for each outcome. A funnel plot of standard error and effect size revealed some funnel plot




Figure 2-1. Forest plot of the VR-based intervention effect in social skills for ASD




Figure 2-2. Forest plot of the VR-based intervention effect in emotion recognition for ASD




Figure 2-3. Forest plot of the VR-based intervention effect in attention for ASD




Figure 2-4. Forest plot of the VR-based intervention effect in executive function for ASD




Figure 2-5. Forest plot of the VR-based intervention effect in conversational skills for ASD




Figure 3-1. Funnel plot for estimates of VR-based intervention effect in social skills




Figure 3-3. Funnel plot for estimates of VR-based intervention effect in attention




Figure 3-2. Funnel plot for estimates of VR-based intervention effect in emotion recognition




Figure 3-4. Funnel plot for estimates of VR-based intervention effect in executive function




Figure 3-5. Funnel plot for estimates of VR-based intervention effect in conversational skills

asymmetry for emotion recognition and conversation skills (Figure. 3-1, 3-2, 3-3, 3-4, 3-5). The Duval and Tweedie (2000) trim and fill procedure was conducted. The filled effect sizes for emotion recognition and conversational skills were not significant: for emotion recognition ($g = .29$, 95% $CI = -.02-.60$), for conversational skills ($g = .20$, 95% $CI = -.21-.62$).

Discussion

The current meta-analysis of VR-based intervention trials (7 studies, $n = 150$) revealed that, for the developmental functions related to ASD including social skills, emotion recognition, attention executive function and conversational skills, VR-based interventions have medium effect size.

Considering publication bias, the effects were not significant for emotion recognition and conversational skills. From this sensitivity analysis, we can conservatively infer that the effects of the VR-based interventions for emotion recognition and conversational skills were not significant. However, the improvement in social skills, attention and executive function remained significant.

When there were no significant effects for emotion recognition and conversational skills, we may consider these functions are relatively inherent characteristics of ASD, which are hard to change and require a sophisticated work. According to a literature review (Danaka & Sung, 2016), emotion recognition deficits in ASD are plausibly accounted for by their avoiding the eye region. Studies showed that direct eye contact elicits a heightened physiological response. As for conversation skills, they require both social cognition and perspective taking (Dixon et al. 2009) that can not be adopted in a short period. Lack of these underlying skills may have not made enough improvement in the conversational skills despite the VR intervention.

Further, when we looked at the characteristics of subgroups that were grouped based on mean effect sizes, the effects of the VR-based interventions were different by some characteristics such as

age, ASD functioning, number of sessions, country where the study conducted and additional treatment. These findings suggested factors to be considered when VR-based intervention is planned for children and young adults with ASD.

For social skills, the study by Kandaloft, et al. (2012), in which samples were diagnosed with Aspergers or PDD-NOS, showed larger effect size compared to the other three studies with the samples diagnosed with ASD or high functioning ASD. This suggests that a VR-based intervention is more effective for those with relatively normal language and intelligence than those below normal in them.

For emotion recognition, there were differences in the country where the study was conducted and the number of sessions operated. The study by Ip et al. (2018) that showed the smaller effect size was constructed with 28 sessions in Hong Kong whereas the other three studies were constructed with 5 or 10 sessions in US. This suggests that a VR-based intervention may have different effect depending on the participants' cultural and language background and can be effective with shorter sessions.

For attention and executive function, the study by Lamash et al. (2017) combined the VR-based intervention with meta-cognition strategies. The participants were introduced

to meta-cognition strategies prior to the intervention. Its larger effect size suggests that combining VR intervention with teaching meta-cognitions strategies may increase the effect size than a single intervention. This finding is supported by Diamond (2013) showing that executive functioning involves attention. Also, a study showed that cortical network underlies both executive functioning and attention (Makris et al., 2007).

For conversational skills, the effect sizes were different by the participants' age. Studies by Amaral et al. (2018) and Kandaloft et al. (2012), in which the participants had the higher mean age, showed larger effect sizes than studies with younger the participants. Older participants may have experienced various situations which require conversational interactions. Our result implies that people with more experiences are better benefited by a VR-based intervention for conversational skills.

In conclusion, VR-based interventions can be considered as an acceptable treatment tool for the developmental functions of children and young adults with ASD, especially social skills, attention and executive function.

This current meta-analysis has limitations to be considered. First, this meta-analysis is limited by the limited number of studies aggregated. More studies on VR-based interventions for ASD in the future can allow

calibrating the effect size. Second, this meta-analysis focused on the virtual reality. The future study can compare it with non-VR based interventions in their effectiveness. Lastly, the future study can test a moderator such as age, ASD functioning, number sessions, country where the study conducted and combined treatment, when VR-based intervention studies are sufficiently accumulated.

Finally, future studies about VRs' negative side effects for children are needed in order to advance VR-based interventions. Several studies of VR-based interventions for adults reported that VR caused no or low side effects including sickness, oculomotor effects, disorientation, and nausea (Freeman, 2010; McMamara et al., 2018).

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Reference

- Amaral, C., Mouga, S., Simões, M., Pereira, H. C., Bernardino, I., Quental, H., ... & Castelo-Branco, M. (2018). A feasibility clinical trial to improve social attention in autistic spectrum disorder (ASD) using a brain computer interface. *Frontiers in neuroscience, 12*, 1-13.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). <https://doi.org/10.1176/appi.books.9780890425596>
- Beach, J., & Wendt, J. (2015). Using virtual reality to help students with social interaction skills. *Journal of the International Association of Special Education, 16*(1), 26-33.
- Bystrova, Y., T., Tokarskaya, V., L., Vuković, B. D. (2019). Optimum virtual environment for solving cognitive tasks by individuals with autism spectrum disorders: the questions and methods of design. *International Journal of Cognitive Research in Science, Engineering and Education (IJCRSEE), 7*(1), 63-72
- Cai, Y., Chiew, R., Nay, Z. T., Indhumathi, C., Huang, L. (2017). Design and development of VR learning environments for children with ASD. *Interactive Learning Environments, 25*(8), 1098-1109.
- Cheng, Y., Huang, C. L., & Yang, C. S. (2015). Using a 3D immersive virtual environment system to enhance social understanding and social skills for children with autism spectrum disorders. *Focus on Autism and Other Developmental Disabilities, 30*(4), 222-236.
- Chia, N., Kee, N. (2014). Application of universal design for learning (Ud11) and

- living (Udl2) in virtual dolphin-assisted intervention (VR) for children with autism. *The Journal of the International Association of Special Education*, 15(1), 75-82.
- Chin, Y. A. (2015). **Stata를 이용한 메타분석**. 서울: Korea University Press.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum Associates.
- De Luca, R., Leonardi, s., Portaro, S., Cause, M., Domenico, C., Colucci, P., ... & Calabrò, R., (2019). Innovative use of virtual reality in autism spectrum disorder: A case-study. *Applied Neuropsychology: Child*. DOI: 10.1080/21622965.2019.1610964.
- Diamond A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135-168.
- Didehbani, N., Allen, T., Kandalaft, M., Krawczyk, D., & Chapman, S. (2016). Virtual reality social cognition training for children with high functioning autism. *Computers in Human Behavior*, 62, 703-711.
- Dixon, D. R., Tarbox, J., & Najdowski, A. (2009). Social skills in autism spectrum disorders. In J. L. Matson (Ed.), *Social behavior and skills in children* (pp. 117-140). New York: Springer Science + Business Media.
- Freeman, D., Pugh, K., Vorontsova, N., Antley, A., & Slater, M. (2010). Testing the continuum of delusional beliefs: An experimental study using virtual reality. *Journal of Abnormal Psychology*, 119(1), 83-92.
- Hwang, S. D. (2016). *Meta-Analysis using OpenMeta, R, or RevMan*. Kyungpook National University Press.
- Ip, H. H., Wong, S. W., Chan, D. F., Byrne, J., Li, C., Yuan, V. S., ... & Wong, J. Y. (2018). Enhance emotional and social adaptation skills for children with autism spectrum disorder: A virtual reality enabled approach. *Computers & Education*, 117, 1-15.
- Jarrold, W., Mundy, P., Gwaltney, M., Bailenson, J., Hatt, N., McIntyre, N., ... & Swain, L. (2013). Social attention in a virtual public speaking task in higher functioning children with autism. *Autism Research*, 6(5), 393-410.
- Kandalaft, M. R., Didehbani, N., Krawczyk, D. C., Allen, T. T., & Chapman, S. B. (2013). Virtual reality social cognition training for young adults with high-functioning autism. *Journal of Autism and Developmental Disorders*, 43(1), 34-44.
- Ke, F., Moon, J. (2018). Virtual collaborative gaming as social skills training for high-functioning autistic children. *British Journal of Educational Technology*, 29(4), 728-741.

- Korean Ministry of Health and Welfare (2017). *National survey on persons with disabilities*.
- Korean Ministry of Health and Welfare (2018). *Status of Registration as Persons with Disabilities*.
- Kuriakose, S., Lahari U. (2015). Understanding the psycho-physiological implications of interaction with a virtual reality-based system in adolescents with autism: A feasibility study. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 23(4), 665-675.
- Lahiri, U., Bekele, E., Dohrmann, E., Warren, Z., & Sarkar, N. (2012). Design of a virtual reality based adaptive response technology for children with autism. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 21(1), 55-64.
- Lahiri, U., Trewyn, A., Warren, Z., & Sarkar, N. (2011). Dynamic eye gaze and its potential in virtual reality based applications for children with autism spectrum disorders. *Autism-Open Access*, 1(1). DOI: 10.4172/2165-7890.1000101.
- Lamash, L., Klinger, E., & Josman, N. (2017). Using a virtual supermarket to promote independent functioning among adolescents with Autism Spectrum Disorder. *International Conference on Virtual Rehabilitation (ICVR)* (pp. 1-7). *IEEE*.
- Lee, S. (2018). A review on focused intervention practices and comprehensive treatment models in autism spectrum disorder. *Learning Disability*, 15(3), 275-313.
- Lorenzo, G., Lledó, A., Pomares, J., & Roig, R. (2016). Design and application of an immersive virtual reality system to enhance emotional skills for children with autism spectrum disorders. *Computers & Education*, 98, 192-205.
- Makris, N., Biederman, J., Valera, E. M., Bush, G., Kaiser, J., Kennedy, D. N., ... Seidman, L. J. (2007). Cortical thinning of the attention and executive function networks in adults with attention-deficit/hyperactivity disorder. *Cerebral Cortex*, 17(6), 1364-1375.
- McNamara, P., Moore, K. H., Papeis, Y., Diallo, S., & Wildman, W. J. (2018). Virtual reality-enabled treatment of nightmares. *Dreaming*, 28(3), 205-224.
- Mesa-Gresa, P., Gil-Gómez, H., Lozano-Quilis, J. A., & Gil-Gómez, J. A. (2018). Effectiveness of virtual reality for children and adolescents with autism spectrum disorder: An evidence-based systematic review. *Sensors*, 18(8), 2486.
- Millen, L., Cobb, S., Patel, H., & Glover, T. (2014). A collaborative virtual environment for conducting design sessions with students with autism spectrum disorder. *International Journal of Child Health and Human Development*,

- 74), 367.
- Miller, H. L., & Bugnariu, N. L. (2016). Level of immersion in virtual environments impacts the ability to assess and teach social skills in autism spectrum disorder. *Cyberpsychology, Behavior, and Social Networking, 19*(4), 246-256.
- Moon, J., & Ke, F. (2019). Exploring the treatment integrity of virtual reality-based social skills training for children with high-functioning autism. *Interactive Learning Environments, 1-15*.
- Politis, Y. (2017). Involving people with autism in development of virtual world for provision of skills training. *International Journal of E-Learning & Distance Education 32*(2), Retrieved February 29, 2020, from <http://www.ijede.ca/index.php/jde/article/view/1032>
- Stendal, K., & Balandin, S. (2015). Virtual worlds for people with autism spectrum disorder: A case study in second life. *Disability and rehabilitation, 37*(17), 1591-1598.
- Stichter, J. P., Laffey, J., Galyen, K., & Herzog, M. (2014). iSocial: Delivering the social competence intervention for adolescents (SCI-A) in a 3D virtual learning environment for youth with high functioning autism. *Journal of Autism and Developmental Disorders, 44*(2), 417-430.
- Tanaka, J. W. & Sung, A. (2016). The "eye avoidance" hypothesis of autism face processing. *Journal of Autism and Developmental Disorder, 46*(5), 1538-1552.
- Wallace, S., Parsons, S., & Bailey, A. (2017). Self-reported sense of presence and responses to social stimuli by adolescents with ASD in a collaborative virtual reality environment. *Journal of Intellectual & Developmental Disability, 42*(2), 131-141.
- Wallace, S., Parsons, S., Westbury, A., White, K., White, K., & Bailey, A. (2010). Sense of presence and atypical social judgments in immersive virtual environments: Responses of adolescents with Autism Spectrum Disorders. *Autism, 14*(3), 199-213.
- Wang, M., & Reid, D. (2013). Using the virtual reality-cognitive rehabilitation approach to improve contextual processing in children with autism. *The Scientific World Journal*. <https://doi.org/10.1155/2013/716890>.
- Wang, X., Laffey, J., Xing, W., Ma, Y., & Stichter, J. (2016). Exploring embodied social presence of youth with Autism in 3D collaborative virtual learning environment: A case study. *Computers in Human Behavior, 55*, 310-321.
- Wass, S. V., & Porayska-Pomsta, K. (2014). The uses of cognitive training technologies in the treatment of autism spectrum disorders. *Autism, 18*(8), 851-

871.

- Wenar, C., & Kerig, P. (2000). *Developmental psychopathology: From infancy through adolescence*. McGraw-Hill.
- Wicks-Nelson, R., Israel, A. (2013). *Abnormal child and adolescent psychology*. New York: Routledge
- Yang, Y. D., Allen, T., Abdullahi, S. M., Pelphrey, K. A., Volkmar, F. R., & Chapman, S. B. (2018). Neural mechanisms of behavioral change in young adults with high functioning autism receiving virtual reality social cognition training: A pilot study. *Autism Research, 11*(5), 713-725.
- Zhao, H., Swanson, A. R., Weitlauf, A. S., Warren, Z. E., & Sarkar, N. (2018). Hand - in - hand: A communication - enhancement collaborative virtual reality system for promoting social interaction in children with autism spectrum disorders. *IEEE Transactions on Human-Machine Systems, 48*(2), 136-148.

자폐스펙트럼장애에서 발달 기능을 위한 가상현실-기반 중재: 메타분석

홍승주¹

김미성²

성균관대학교 아동청소년학과/ 석사과정¹

성균관대학교 아동청소년학과/ 강사²

본 메타분석은 ASD를 가진 아동과 청년들을 위한 VR-기반 중재의 효과에 대한 연구를 집약하고 분석하였다. 2009년부터 2019년 사이에 발표된 연구 중, 사전사후 결과를 포함하고 있는 7개의 VR-기반 중재 연구가 메타분석에 포함되었다. 메타분석에 포함된 연구의 연구대상자는 총 150명이었다. 측정 결과는 ASD의 특성으로 나타나는 발달 기능인 사회적 기술, 정서인식, 주의집중, 실행기능, 의사소통기술로 분류하여 분석하였다. 무선효과모형에 따른 효과크기는 다음과 같이 나타났다: 사회적 기술($g = .59$, $SE = .12$, 95% $CI: .35-.82$), 정서인식($g = .50$, $SE = .16$, 95% $CI: .19-.82$), 주의집중($g = .35$, $SE = .14$, 95% $CI: .07-.63$), 실행기능($g = .52$, $SE = .12$, 95% $CI: .28-.77$), 의사소통기술($g = .46$, $SE = .21$, 95% $CI: .05-.88$). VR-기반 중재의 효과가 대체로 중간이며, 주의집중에 대한 효과크기는 상대적으로 작았다. ASD에 대한 VR-기반 중재의 효과와 관련하여 연령, ASD 기능 수준, 연구가 이루어진 국가, 중재 회기 수 및 VR-기반 중재와 함께 제공된 추가 치료개입 등 중재의 효과에 영향을 미칠 수 있는 여러 환경 요인을 논의하였다.

주요어 : 자폐스펙트럼장애, 가상현실, 사회적기술, 정서인식, 주의집중, 실행기능, 의사소통기술