

## Asymbiotic seed germination and *in vitro* propagation condition in *Calanthe discolor* Lindl.

Huyk Joon Kwon<sup>1</sup>, So Lim Shin<sup>1</sup>, Kee Hwa Bae<sup>1,2</sup> and Soo-Young Kim<sup>1,\*</sup>

<sup>1</sup>National Institute of Biological Resources, 42 Hwangyeong-ro, Seo-gu, Incheon 22689, Republic of Korea

<sup>2</sup>Nakdonggang National Institute of Biological Resources, 137 Donam 2-gil, Sangju-si, Gyeongsangbuk-do 37242, Republic of Korea

\*Correspondent: sy7540@korea.kr

We investigated the effects of sodium hypochlorite (NaOCl) and culture medium on embryo swelling and germination of *Calanthe discolor* Lindl., and established a method for determining the swelling and protocorm formation of *C. discolor* seeds via *in vitro* examination of immature seeds. Treatment of immature seeds with NaOCl greatly enhanced the extent of embryo swelling and protocorm formation of immature zygote embryos compared to seeds without NaOCl treatment. The effects of the culture media were also evaluated with regard to embryo swelling and protocorm formation of *in vitro* cultured seeds with and without NaOCl treatment. Additionally, the effects of white fluorescent light and red and blue LEDs lights on seedling growth in *in vitro* culture were examined. The most suitable condition for seedling growth after 12 weeks of culture was the red LEDs light with POM medium. These results show effective asymbiotic germination and growth of *C. discolor*.

Keywords: chlorophyll, germination, growth, LEDs, terrestrial orchid

© 2017 National Institute of Biological Resources  
DOI:10.12651/JSR.2017.6.1.068

### INTRODUCTION

Orchidaceae comprise a fascinating group of ornamental plants, and numerous novel cultivars have been produced by interspecific and intergeneric hybridization of plants with exotic and elegant flowers. However, many orchid species are endangered as a consequence of environmental disruption, succession of natural habitats, and overexploitation for horticultural purposes. *In situ* conservation by preservation and enhancement of dwindling populations of endangered orchid species is difficult because of the relatively slow growth and low germination rate, and because these processes require symbiotic relationships with mycorrhizal fungi in natural habitats (Bernard, 1903; Warcup, 1973; Rasmussen, 1995).

The genus *Calanthe* is widely distributed in tropical and subtropical regions, including Africa, South America, Asia, and the Pacific Islands (Gale and Drinkell, 2007; Lee *et al.*, 2008). It contains approximately 171 species worldwide, of which approximately 29 inhabit Asian regions, such as Korea, Japan, and China (Lee *et al.*, 2008). However, only four species (*Calanthe discolor*

Lindl., *C. sieboldii* Ohwi, *C. reflexa* Maximowicz, and *C. aristulifera* Hayata) and two natural hybrids (*Calanthe* × *bicolor* and *Calanthe* × *kibanakirishima*) (Hyun *et al.*, 1999a; 1999b; Lee and Choi, 2006; Lee *et al.*, 2008) inhabit Korea.

*Calanthe discolor* has beautiful flowers that consist of a green perianth and red lips in raceme inflorescences, and is highly valued. Therefore, establishment of protocols for mature seed germination in this species is a prerequisite for *ex situ* conservation. Seed germination is the limiting step for the establishment of plantlets from the germination of terrestrial orchids because germination is more difficult than in epiphytic species (Arditti and Ernst, 1984). Kano (1972) reported that *Calanthe* germination in temperate regions is the most difficult among all terrestrial orchid species studied. Several studies have reported propagation of *Calanthe* species by asymbiotic germination (Miyoshi and Mii, 1988; 1995a; 1995b; Park *et al.*, 2000; Lee *et al.*, 2007). Various strategies, such as presoaking of seeds in water (Mii and Kako, 1974), organic solvent treatment (Mii, 1978), phytohormones treatment (Miyoshi and Mii, 1995a), pre-

treatment with ultrasonication (Miyoshi and Mii, 1988), and the addition of a polyphenol absorbent to the medium (Miyoshi and Mii, 1995b), have been used in efforts to improve the germination and/or protocorm formation of *Calanthe* species. Recently, Lee *et al.* (2007) collected seeds of *C. tricarinata* at various stages of seed formation and maturation, and the frequency of germination of immature seeds at 150 days postanthesis reached approximately 30%. The germination frequency of fully matured seeds was low and only a few seeds germinated. Lee *et al.* (2007) also noted that scarification, for example, by prolonged treatment with chemical solutions and ultrasound, stimulated the germination of fully matured seeds and resulted in frequencies of seed germination (32% to 39%) similar to that of immature seeds.

The effectiveness of seed disinfection reagents such as sodium hypochlorite (NaOCl) and calcium hypochlorite  $\text{Ca}(\text{OCl})_2$  for stimulation of orchid germination has been reported in *Dendrobium* species (Malmgren, 1996). Previously, we reported that NaOCl treatment effectively stimulated the *in vitro* germination of seeds in the genus *Cypripedium* (Bae *et al.*, 2009; Bae *et al.*, 2010; Bae and Choi, 2013). The mechanism underlying this stimulating effect on germination of orchids is not well understood (Harvais and Hadley, 1967). Germination of epiphytic orchids and some terrestrial orchids has been observed under both dark and light conditions (Werckmeister, 1971; Ueda, 1972; Arditti and Ernst, 1984), whereas stimulation of seed germination in darkness has been described for many terrestrial species (Harvais, 1973; Stoutamire, 1974; Arditti and Ernst, 1984; Kauth *et al.*, 2006). Additionally, there is a need for an efficient light source to improve production efficiency, reduce costs, and improve the quality of micropropagated plants. The utilization of a more versatile, efficient light source for *in vitro* plant regeneration and growth could offer new and important possibilities to achieve success in commercial micropropagation. Light-emitting diodes (LEDs) have been proposed as a primary light source for space-based plant research chambers and bioregenerative life-support systems, and as a potential alternative light source for *in vitro* plant growth and development (Bula *et al.*, 1991; Yeh and Chung, 2009; Nhut and Nam, 2010). However, most LEDs studies examined vegetables (pepper, tomato, and onion); only a few studies have evaluated orchid (Bae *et al.*, 2014) and *Gerbera* spp. (Kwon *et al.*, 2012).

In the present study, we evaluated the effects of NaOCl treatment and LEDs illumination on germination and seedling growth, respectively, for *C. discolor*. We also discuss the importance of a reproducible protocol for establishment of germination and seedling growth from seeds using asymbiotic culture.

## MATERIALS AND METHODS

### Plant materials and culture conditions

Capsules of *C. discolor* were collected from a single population near Youngarioreum in Jeju, Korea, in July 2013 (voucher specimen, NIBRVP0000439107 and seed number, KWPSGR0000141268). Following collection, the immature seeds were immersed in deionized, sterilized water and agitated for 30 min.

### Morphological observation on seed coat by Scanning Electron Microscopy (SEM)

For scanning electron microscopy (SEM) selected seeds were mounted directly on stubs using double-sided adhesive tape and coated with gold in a sputter coater. The changes of seed coat by 1% NaOCl treatment was observed. The SEM micrographs were taken in a SEM (Jeol, Japan) at an accelerating voltage of 20 kV.

### Effects of NaOCl and culture medium on embryo swelling and seed germination

Mature seed capsules were then treated with 30 mL of 1% NaOCl in deionized water (v/v) for 30 min, followed by three 30-s rinses in deionized, sterilized water. The capsules were then cleaved with a scalpel blade, and the seeds were scraped off. The seeds were soaking treated with 1% NaOCl for 30 min. Subsequently, the seeds were left in the final rinse water until transfer to POM (Phytomax Orchid Maintenance, P6668; Sigma, USA) medium, SGM (Seed Germination Maintenance, P6543; Sigma, USA) medium, and MS (Murashige and Skoog, 1962) medium after 10 weeks of culture on medium supplemented with  $20 \text{ g} \cdot \text{L}^{-1}$  sucrose and  $3.0 \text{ g} \cdot \text{L}^{-1}$  gelrite.

All media were supplemented with  $20 \text{ g} \cdot \text{L}^{-1}$  sucrose and the pH was adjusted to 5.5 with 0.1 M KOH before addition of  $3.0 \text{ g} \cdot \text{L}^{-1}$  gelrite. Media were autoclaved at 117.7 kPa and 121-°C for 15 min. The cultures were maintained in a growth room at  $20 \pm 2^\circ\text{C}$ . Two basal media were used in this study: POM and SGM media without plant growth regulators. Embryo swelling was defined as at least a doubling in size and formation of a protocorm. The number of swollen embryos and embryo diameter were recorded every 2 weeks by examination was replicated three times with a microscope.

### Effects of LEDs on seedling growth and analysis of chlorophyll contents

Protocorms were transferred to POM and MS medium (pH 5.2, sucrose  $20 \text{ g} \cdot \text{L}^{-1}$ , gelrite  $3.0 \text{ g} \cdot \text{L}^{-1}$ ) for seedling. Cultures were incubated in LEDs growth chambers (Se-

jong Scientific, Korea) at  $23 \pm 2^\circ\text{C}$  and 60% relative humidity. Seedlings were subjected to three different culture treatments with the following sources of illumination: (1) fluorescent light (FL,  $40\text{--}70 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ), (2) red LEDs (RL, peak wavelength: 660 nm,  $3.5\text{--}4.0 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ), and (3) blue LEDs (BL, peak wavelength: 450 nm,  $15\text{--}20 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ ). The duration of lighting for all treatments was 16 h per day. Plantlet height was evaluated by measuring the average length of the shoots and roots and the weight per plantlet after 12 weeks of culture. The content of chlorophyll a and b and the total chlorophyll content were determined using methods developed by Hiscox and Israelsham (1979) with some modifications. The color intensity of the resulting extract was measured using a spectrophotometer (Shimadzu, UV-120-01) at 645 nm, 663 nm and the chlorophyll content (mg/g fresh weight) was calculated using the following formulae. Chl a (mg/g FW of leaf) =  $(\text{OD}_{663}0.127 - \text{OD}_{645}0.00269) \times 100$ , Chl b (mg/g FW of leaf) =  $(\text{OD}_{645}0.0229 - \text{OD}_{663}0.00468) \times 100$ , total chlorophyll content =  $(\text{OD}_{645}0.0202 - \text{OD}_{663}0.00802) \times 100$  and ratio of Chl a to Chl b = Chl a/Chl b.

### Statistical analysis

All data were expressed as the mean  $\pm$  standard error (SE) and analyzed using an analysis of variance (ANOVA). All experiment was replicated three times with at least 200 seeds per replication. Significant differences among the treatments were determined by performing multiple comparison tests using Duncan's multiple range test with  $p < 0.05$  defined as significant (SAS, 2003).

## RESULTS AND DISCUSSION

### Effect of NaOCl treatment and culture medium on seed germination


The effect of NaOCl treatment and media on seed germination was examined. After 30 min of 1% NaOCl treatment, the seed coat was bleached completely and the zygotic embryo inside the seed coat could be clearly seen (Fig. 1C). Embryo swelling was first scored 2 weeks after sowing (Fig. 1D). After NaOCl treatment of the seeds, embryo swelling was frequent after 8 weeks of culture. The highest embryo swelling in zygotic embryos was noticed when seeds were treated for 30 min with 1% NaOCl in POM medium (Table 1). POM medium (88.2%) induced significantly greater embryo swelling than SGM (55.4%) and MS (66.7%) media. When immature seeds of *C. discolor* were cultured on POM, SGM, and MS media without 1% NaOCl treatment 11.4%, 8.2%, and 4.1% swelling was observed 2 weeks of cul-

ture (Table 1).

After 8 weeks in POM medium without NaOCl treatment, the protocorm formation was very low compared to that in seeds treated with NaOCl treatment (Table 1). In contrast, immature seeds treated with NaOCl began to germinate within 6 weeks (Fig. 1E). Immature seeds with NaOCl treatment cultured in POM medium showed a greatly enhanced frequency of protocorm formation (Table 1). The maximum protocorm formation was recorded for seeds treated with 1% NaOCl for 30 min and cultured on POM medium (82.3%), followed by SGM (44.1%) and MS (14.8%) medium, whereas the three media without NaOCl treatment showed the least protocorm formation (Table 1). Morphological development of *C. discolor* from seed to protocorms was observed (Fig. 1E-G). The maximum embryo diameter was recorded for seeds treated with 1% NaOCl and cultured on POM medium (Fig. 2A), followed by SGM (Fig. 2B) medium, and MS medium (Fig. 2C). Within 10 weeks of culture on POM medium, the plantlets attained a height of 2-2.5 cm with converting shoot primordia and 1-2 roots (Fig. 1H). The well-rooted plantlets (approximately 2-3 cm long with 3-4 roots) were removed from the POM medium (Fig. 1I).

SEM analysis revealed that NaOCl treatment resulted in the small perforation of seed coats (Fig. 1B) but intact in NaOCl non-treatment (Fig. 1A), indicating that NaOCl treatment strongly affect the rigidity of seed coat by decomposition of cell wall materials. Seed germination of some orchid species is typically very low or nonexistent in *ex vitro* and *in vitro* conditions (Ault and Blackmon, 1987; Anderson, 1996). Terrestrial orchids have more stringent requirements for germination, but little information is available regarding the specific requirements for terrestrial orchids (Fast, 1982). NaOCl is a disinfecting agent that is widely used for seed surface sterilization and it is also known to favor seed germination or to overcome seed dormancy in *Cypripedium reginae*, *C. parviflorum* and *Platanthera grandiflora* (Vujanovic *et al.*, 2000). Yildiz and Celal (2002) reported that treatment with NaOCl for 20 minutes enhanced the germination of *Linum usitatissimum* seeds. The promotion of germination by NaOCl is thought to be due to scarification of the seed coat, which allows more water and oxygen absorption, or from the enhancement of oxidative respiration by the extra supply of oxygen arising from decomposition of NaOCl (Vujanovic *et al.*, 2000). In addition, the effectiveness of disinfection agents such as NaOCl and  $\text{Ca}(\text{OCl})_2$  for stimulation of germination of orchid seeds has been reported in *Cypripedium acaule* and *Dendrobium* species (St-Arnaud *et al.*, 1992; Malmgren, 1996), but the underlying mechanism has not yet been reported. The possible mechanisms of action underlying the in-






**Fig. 1.** Developmental stages of asymbiotically cultured *Calanthe discolor* seeds. (A) surface of seed coat without NaOCl treatment; (B) surface of seed coats after 1% NaOCl treatment for 30 min by SEM (arrows: perforation); (C) hyaline embryo, seed coat intact; (D) embryos swollen after 2 weeks of culture; (E) swelled embryos present rhizoids after 6 weeks of culture; (F) appearance of protomeristem and rhizoid elongation after 7 weeks culture; (G) appearance of chlorophyllous protomeristem after 10 weeks of culture; (H) shoot formation *via* the shoot axis; (I) The well-developed plantlets were removed from POM medium; Scale bars: A, B = 100  $\mu$ m, C-G = 1.0 mm, H, I = 25.0 mm.

**Table 1.** Effect of 1% NaOCl treatment and culture media on embryo swelling and protocorm formation from *Calanthe discolor* seeds.

NaOCl treatment time (min)	Swelled embryo formation (2 week after sowing, %)			Protocorm formation (8 week after sowing, %)		
	POM <sup>z</sup>	SGM	MS	POM	SGM	MS
0	11.4 d <sup>y</sup>	8.2 e	4.1 f	1.3 d	0.6 d	0.2 d
30	88.2 a	55.4 c	66.7 b	82.3 a	44.1 b	14.8 c


<sup>z</sup>POM, SGM and MS indicate phytomax orchid maintenance medium, seed germination medium and Murashige and Skoog medium, respectively.

<sup>y</sup>Different alphabetical letters are significantly different according to Duncan's multiple range test at  $P < 0.05$ .



**Fig. 2.** Effects of culture medium and 1% NaOCl treatment on embryo diameter of *Calanthe discolor* after 0, 2, 4, 6 and 8 weeks *in vitro* culture. (A) Effect of treatment with or without 1% NaOCl on seeds in POM medium; (B) SGM medium; (C) MS medium. Values are represented as the mean  $\pm$  SD.

duction of seed germination or disruption of dormancy by NaOCl have been considered to be partial degradation of the seed coat and/or the solubilization and oxida-



**Fig. 3.** Effect of light source (fluorescent light (FL) and red (RL) or blue LEDs light (BL)) on seedling growth (shoot (SLT) and root length (RLT), fresh weight (FWT) measurement) of *Calanthe discolor* after 12 weeks of culture on POM medium supplemented with sucrose ( $20 \text{ g} \cdot \text{L}^{-1}$ ) and gelrite ( $3.0 \text{ g} \cdot \text{L}^{-1}$ ). Values are represented as the mean  $\pm$  SD. <sup>a-c</sup>Different alphabetical letters are significantly different according to Duncan's multiple range test with significant defined as  $P < 0.05$ .

tion of growth inhibitors. Harvais (1982) considered the stimulating effect of surface sterilization with NaOCl to be a physiological effect of the endogenous inhibitor abscisic acid (ABA) being washed away from the seed.

### Effect of LEDs light on seedling growth and chlorophyll measurement

Shoot length at 12 weeks after culture was 2.74 cm, 3.11 cm, and 1.39 cm under fluorescent light (FL), red LEDs light (RL), and blue LEDs light (BL), respectively, in the POM medium (Fig. 3). The root length measurements were all significantly greater for seedlings cultured under BL with a wavelength of 625 nm (Fig. 3). The highest total chlorophyll content was observed in seedlings cultured under RL ( $4.55 \text{ mg} \cdot \text{g}^{-1}$ ), followed by FL ( $3.37 \text{ mg} \cdot \text{g}^{-1}$ ), and BL ( $0.11 \text{ mg} \cdot \text{g}^{-1}$ ) (Table 2). The induction of protocorm growth using LEDs has also been previously reported in *Calanthe satsuma* (Fukai *et al.*, 1997). The inhibition of seed germination by lighting has been reported for many terrestrial orchid species, such as *Calanthe tricarinata* (Godo *et al.*, 2010) and

**Table 2.** Effect of light source (LEDs) on chlorophyll content of *Calanthe discolor* after 12 weeks of culture on POM medium.

Light sources	Chlorophyll contents			
	Chl a	Chl b	Total Chlorophyll contents	Chl a ratio b
FL <sup>z</sup>	3.21 b <sup>y</sup>	0.17 b	3.37 b	18.52 a
RL	4.32 a	0.23 a	4.55 a	18.87 a
BL	0.08 c	0.02 c	0.11 c	5.23 b

<sup>z</sup>FL, RL and BL indicate fluorescent light, red LEDs (660 nm) and blue LEDs (450 nm), respectively.

<sup>y</sup>Different alphabetical letters are significantly different according to Duncan's multiple range test at  $P < 0.05$ .

*Habenaria macroceratitis* (Stewart and Kane, 2006). LEDs lighting systems also regulate *in vitro* seedling growth. In *Bletilla ochracea*, a maximum seedling growth of 55% was obtained using various LEDs lights (Godo *et al.*, 2011).

This study demonstrated an efficient micropropagation system for *C. discolor* using seeds and axenic seedling-derived rhizome for explants culture. A symbiotic seed germination and cloning through rhizomes provided a large number of vigorous seedlings and rhizome-derived clones for ornamental exploitation and eco-restoration of the ensuing rhizomes as the storage organ for propagation. Association with a symbiotic fungus, at least during transplantation from *in vitro* culture, may be an option to enhance growth and to establish species in the field. However, the protocol described in the present study could be useful for commercial nurseries that conduct mass propagation and *ex situ* conservation of *C. discolor*. It is thought that seed propagation and protocorm formation of *C. discolor* can be facilitated by 30 min of 1% NaOCl treatment.

In conclusion, red LEDs light and 1% NaOCl treatment conditions enhance the germination and protocorm formation of *C. discolor*. Therefore, the optimal *C. discolor in vitro* propagation methods of submerging the seeds in a 1% NaOCl for 30 min then sowing them in POM medium at red LEDs light.

## ACKNOWLEDGEMENTS

This work was carried out with the support of NIBR project (NIBR201619101).

## REFERENCES

- Anderson, A.B. 1996. The re-introduction of *Platanthera ciliaris* in Canada. In: C. Allen (ed.), Proceedings of the North American Native Terrestrial Orchids: Propagation and Production. North American Native Terrestrial Orchid Conference, Maryland. pp. 73-76.
- Arditti, J. and R. Ernst. 1984. Physiology of germinating orchid seeds. In: J. Arditti (ed.), Orchid biology: Reviews and perspectives 3. Cornell University Press, Ithaca.
- Ault, J.R. and W.J. Blackmon. 1987. *In vitro* propagation of *Ferocactus acanthodes* (Cactaceae). Hort. Sci. 22:126-127.
- Bae, K.H., C.H. Kim, B.Y. Sun and Y.E. Choi. 2010. Structural changes of seed coats and stimulation of *in vitro* germination of fully mature seeds of *Cypripedium macranthos* Swartz by NaOCl pretreatment. Prog. Ornament. Plant 10:107-113.
- Bae, K.H., H.K. Kwon and Y.E. Choi. 2009. *In vitro* germination and plantlet conversion from the culture of fully mature seeds of *Cypripedium guttatum* Swartz. Prog. Ornament. Plant 9:160-165.
- Bae, K.H. and Y.E. Choi. 2013. Factors affecting fruit bearing in natural habitat and *in vitro* culture of zygotic embryos of *Cypripedium japonicum*. Prog. Ornament. Plant 11: 146-152.
- Bae, K.H., K.H. Oh and S.Y. Kim. 2014. Sodium hypochlorite treatment and light-emitting diode (LED) irradiation effect on *in vitro* germination of *Oreorchis patens* (Lindl.) Lindl. J. Plant Biotechnol. 41:44-49.
- Bernard, N. 1903. La germination des orchidées. Revue Générale de Botanique 16:458-475.
- Bula, R.J., R.C. Morrow, T.W. Tibbitts, D.J. Barta, R.W. Ingnatius and T.S. Martin. 1991. Light-emitting diodes as a radiation source for plants. Hort. Sci. 26:203-205.
- Fast, G. 1982. European terrestrial orchids. Symbiotic and asymbiotic methods. In: J. Arditti (ed.), Orchid biology: reviews and perspectives II. Ithaca, NY: Comstock Publishing Associates. pp. 326-329.
- Fukai, S., K. Fujikawa, K. Okamoto, A. Hasegawa and M. Goi. 1997. Effects of red and blue light on germination and protocorm growth of *Calanthe satsuma*. Lindleyana 12:169-171.
- Gale, S. and C. Drinkell. 2007. *Calanthe arisanensis* Orchidaceae. Curtis's Bot. Mag. 24:206-210.
- Godo, T., K. Fujiwara, K. Guan and K. Miyoshi. 2011. Effect of wavelength of LED-light on *in vitro* asymbiotic germination and seedling growth of *Bletilla ochracea* (Orchidaceae), Plant Biotechnol. 28:397-400.
- Godo, T., M. Komori, E. Nakaoki, T. Yukawa and K. Miyoshi.



2010. Germination of mature seeds of *Calanthe tricarinata* Lindl., an endangered terrestrial orchid, by asymbiotic culture *in vitro*. *In Vitro Cell Dev. Biol. Plant* 46: 323-332.
- Harvais, G. 1973. Growth requirements and development of *Cypripedium reginae* in axenic culture. *Can. J. Bot.* 51:327-332.
- Harvais, G. 1982. An improved culture medium for growing the orchid *Cypripedium reginae* axenically. *Can. J. Bot.* 51:327-332.
- Harvais, G. and G. Hadley. 1967. The relation between host and endophyte in orchid mycorrhiza. *New Phytol.* 66: 205-215.
- Hiscox, J.D. and G.F. Israelsham. 1979. A method for the extraction of chlorophyll from leaf tissue without maceration, *Can. J. Bot.* 57:1332-1334.
- Hyun, M.R., J.Y. Choi, J.N. Suh, I.S. So and J.S. Lee. 1999a. Isozyme and Randomly Amplified Polymorphic DNA (RAPD) analysis for genetic relationship among *Calanthe discolor*, *C. sieboldii*, and *C. bicolor* native to Cheju Island. *Korean J. Hort. Sci. Technol.* 17:141-143.
- Hyun, M.R., J.Y. Choi, J.N. Suh, I.S. So and J.S. Lee. 1999b. Studies on distributions and morphological characteristics of *Calanthe discolor*, *C. sieboldii*, and *C. bicolor* native to Cheju Province. *Kor. J. Hort. Sci. Technol.* 17: 497-499.
- Kano, K. 1972. *The orchids*. Seibundo-shinkosha, Tokyo.
- Kauth, P.J., W.A. Vendrame and M.E. Kane. 2006. *In vitro* culture and seedling development of *Calopogon tuberosus*. *Plant Cell Tiss. Org. Cult.* 85:91-102.
- Korea Forest Service. 2008. Rare and Endangered plants. pp. 332.
- Kwon, M.K., H.H. Nam, J.J. Sung and J.H. Lim. 2012. Effect of Light-emitting Diodes on growth of *in vitro* propagules in *Gerbera hybrida*. *Flower Res. J.* 20(4):233-237.
- Lee, G.J., K.S. Kim, J.S. Kim, J.H. Park, J.B. Kim, D.S. Kim and S.Y. Kang. 2008. A wide array of phenotypic components and their contributions to variations in *Calanthe* orchid landraces. *Hort. Environ. Biotechnol.* 49:418-426.
- Lee, J.S. and B.H. Choi. 2006. Distribution and red data of wild orchids in the Korean Peninsula. *Kor. J. Plant Taxon* 36:335-360.
- Lee, Y.I., F.L. Lu, E.C. Yeung and M.C. Chung. 2007. Developmental change in endogenous abscisic acid concentrations and asymbiotic seed germination of a terrestrial orchid, *Calanthe tricarinata* Lindl. *J. Amer. Soc. Hort. Sci.* 132:246-252.
- Malmgren, S. 1996. Orchid propagation: theory and practice. In: C. Allen (ed.), *Proceedings of the North American Native Terrestrial Orchids: Propagation and Production*. North American Native Terrestrial Orchid Conference, Maryland. pp. 63-71.
- Mii, M. 1978. Effect of organic solvent pretreatment on germination of *Calanthe discolor*. *Proc. Japan Soc. Hort. Sci. Conf.* pp. 330-331.
- Mii, M. and S. Kako. 1974. Studies on germination of *Calanthe izu-insularis*. I. Effect of water treatment and light condition on germination. *Proc. Japan Soc. Hort. Sci. Conf.* pp. 326-327.
- Miyoshi, K. and M. Mii. 1988. Ultrasonic treatment for enhancing seed germination of terrestrial orchid *Calanthe discolor* in asymbiotic culture. *Sci. Hortic.* 35:127-130.
- Miyoshi, K. and M. Mii. 1995a. Phytohormone pre-treatment for the enhancement of seed germination and protocorm formation by terrestrial orchid, *Calanthe discolor* (Orchidaceae), in asymbiotic culture. *Sci. Hortic.* 63:263-267.
- Miyoshi, K. and M. Mii. 1995b. Enhancement of seed germination and protocorm formation in *Calanthe discolor* (Orchidaceae) by NaOCl and polyphenol absorbent treatments. *Plant. Tiss. Cult. Lett.* 12:267-272.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant* 15:473-497.
- Nhut, D.T. and N.B. Nam. 2010. Light emitting diodes (LEDs): an artificial lighting source for biological studies. In: *Proceedings of the 3rd international conference of the development of BME in Vietnam*, 11-14 January 2010.
- Park, S.Y., H.N. Murthy and K.Y. Paek. 2000. *In vitro* seed germination of *Calanthe sieboldii*, an endangered orchid species. *J. Plant. Biol.* 43:158-161.
- Rasmussen, H. 1995. *Terrestrial orchids from seed to mycotrophic plant*. Cambridge University Press, Cambridge.
- SAS, Institute Inc. 2003. SAS version 9.0. SAS Institute, North Carolina.
- St-Arnaud, M.D., D. Lauzer and D. Barabe. 1992. *In vitro* germination and early growth of seedlings of *Cypripedium acaule* (Orchidaceae). *Lindleyana* 7:22-27.
- Stewart, S.L. and M.E. Kane. 2006. Asymbiotic germination and *in vitro* seedling development of *Habenaria macroceratitis* (Orchidaceae), a rare Florida terrestrial orchid. *Plant Cell Tiss. Org. Cult.* 86:147-158.
- Stoutamire, W. 1974. Terrestrial orchid seedling. In: C.L. Withner (ed.), *The orchids: Scientific studies*. Wiley-Interscience, New York. pp. 101-128.
- Ueda, H. 1972. Effects of light and culture medium on adventitious root formation by *Cymbidium*s in aseptic culture. *Amer. Orchid Soc. Bull.* 41:322-327.
- Vujanovic, V., M. St-Arnaud, D. Barabe and G. Thibeault. 2000. Viability testing of orchid seed and the promotion of colouration and germination. *Ann. Bot.* 86:79-86.
- Warcup, J.H. 1973. Symbiotic germination of some Australian terrestrial orchids. *New Phytol.* 72:387-392.
- Werckmeister, P. 1971. Light induction of geotropism and the control of proliferation and growth of *Cymbidium* in tissue culture. *Bot. Gaz.* 132:346-350.
- Yeh, H. and J.P. Chung. 2009. High-brightness LEDs-energy efficient lighting sources and their potential in indoor plant cultivation. *Renew. Sust. Energy Rev.* 13:2175-

2180.

Yildiz, M. and E.R. Celal. 2002. The effect of sodium hypochlorite solution on *in vitro* seedling growth and shoot regeneration of flax (*Linum usitatissimum*). Natul. wissen. Schffen. 89:259-261.

*Submitted: June 27, 2016*

*Revised: August 31, 2016*

*Accepted: February 7, 2017*