

# First report and morphological description of two *Acrobeloides* species (Nematoda: Rhabditida: Cephalobidae) in South Korea

Taeho Kim<sup>1</sup>, Yucheol Lee<sup>2</sup> and Joong-Ki Park<sup>2,\*</sup>

<sup>1</sup>Animal Resources Division, National Institute of Biological Resources, Incheon 22689, Republic of Korea

<sup>2</sup>Division of EcoScience, Ewha Womans University, 52 Ewhayeodae-gil, Seodaemun-gu, Seoul 03760, Republic of Korea

\*Correspondent: jkpark@ewha.ac.kr

The genus *Acrobeloides* (Cobb, 1924) Thorne, 1937 are bacterial feeders and are one of the most abundant and widely distributed nematode groups in various terrestrial environments. Based on morphological and morphometric analyses, we found two *Acrobeloides* species reported in Korea for the first time: *A. bodenheimeri* (Steiner, 1936) Thorne, 1937 and *A. tricornis* (Throne, 1925) Thorne, 1937. These species exhibit morphological characters concordant with typical features of the genus *Acrobeloides*, such as a fusiform pharyngeal corpus with swollen metacarpus and lateral incisures extending to the tail terminus. However, *A. bodenheimeri* is distinguished from other acrobeloids by having its low and rounded labial probolae, distinct post-uterine sac and five lateral incisures. *Acrobeloides tricornis* is distinguished from its congeners by the following characteristics: its high labial probolae with acute termini, inconspicuous post-uterine sac and five lateral incisures. Morphological characters and their measurements, and illustrations of *A. bodenheimeri* and *A. tricornis* are described in this study.

Keywords: *Acrobeloides bodenheimeri*, *Acrobeloides tricornis*, Cephalobidae, Nematoda, South Korea

© 2021 National Institute of Biological Resources  
DOI:10.12651/JSR.2021.10.4.405

## INTRODUCTION

Members of the genus *Acrobeloides* (Cobb, 1924) Thorne, 1937, belonging to the family Cephalobidae, are bacterial feeders and are one of the most abundant and widely distributed nematode groups in various terrestrial environments, such as agricultural land (Pervez, 2011), forest (Háněl, 1999), highlands (Boström, 1993) and sand dunes (Yeates, 1967). Morphological and morphometric variations (such as body size, nerve ring and excretory position and tail shape) in this group have been reported from previous studies (Anderson, 1965; 1968; De Ley *et al.*, 1999; Abolafia and Peña-Santiago, 2003). These morphological variations often hinder species delimitation and identification that mislead species-level taxonomy. To date, 40 nominal species have been reported in this genus; however, only 29 species may be accepted as valid (Andrássy, 2005), including two species previously reported in Korea (Kim *et al.*, 2016; 2017): *A. nanus* (de Man, 1880) Anderson, 1968 and *A. varius* Kim, Kim and Park, 2017.

Following a survey of plots of farmland and overgrown fields, *A. bodenheimeri* (Steiner, 1936) Thorne, 1937 and *A. tricornis* (Thorne, 1925) Thorne, 1937 were isolated

from soil samples. In this paper, we provide a detailed description of the morphological characters and morphometrics, and illustrations of these two species.

## MATERIALS AND METHODS

### Nematode isolation

Soil samples were collected from a potato farm (Bukmyeon, Uichang-gu, Changwon-si, Gyeongsangnam-do, South Korea [GPS coordinates: 35°22'22.5"N, 128°36'47.3"E]) and overgrown fields (Mechuri Island, Daenam-ro, Danwon-gu, Ansan-si, Gyeonggi-do, South Korea [GPS coordinates: 37°12'02.9"N, 126°32'24.6"E]). Nematode specimens were extracted from soil samples using the Baermann funnel method (Baermann, 1917).

### Sample processing and morphological observations

For fixation, the nematode specimens were transferred to a 15 mL tube containing 2 mL water, to which was added 4 mL of 80°C TAF (2% triethanolamine and 7% formaldehyde) solution. The fixed nematodes were dehydra-

**Table 1.** Morphometric measurements of *Acrobelloides bodenheimeri* and *A. tricornis*.

Character	<i>Acrobelloides bodenheimeri</i>	<i>Acrobelloides tricornis</i>	Character	<i>Acrobelloides bodenheimeri</i>	<i>Acrobelloides tricornis</i>
	♀, n = 1	♀, n = 1		♀, n = 1	♀, n = 1
L	529.5	332.0	Excretory pore position (% pharynx)	63.8	79.7
a	27.2	17.5	Deirid position (% pharynx)	71.5	86.9
b	3.4	3.3	Vulva from anterior end	369.6	220.2
c	19.3	15.9	Vulva to anus	133.6	87.8
c'	2.0	1.9	Vulva to anus/tail length	4.9	4.2
V	69.8	66.3	Reproductive tract length	207.0	95.2
G	39.1	28.7	Vagina	12.2	5.1
Body diameter	19.5	19.0	Post-uterine sac	50.3	–
Pharynx length	156.3	99.8	Uterus	62.1	34.4
Tail length	27.4	20.8	Spermatheca	36.9	17.6
Anal body diameter	13.7	10.8	Oviduct	11.0	13.8
Tail annuli <sup>a</sup>	14	15	Ovary	155.4	85.9
Lips region diameter	7.5	6.1	Vagina/body diameter	0.6	0.3
Stoma	12.2	10.7	Post-uterine sac/body diameter	2.6	–
Stoma diameter	4.3	4.2	Uterus/body diameter	3.2	1.8
Stoma/lips region diameter	1.6	1.8	Spermatheca/body diameter	1.9	0.9
Stoma/stoma diameter	2.8	2.6	Oviduct/body diameter	0.6	0.7
Corpus	91.3	52.0	Ovary/body diameter	8.0	4.5
Isthmus	29.3	19.5	Rectum	19.9	12.9
Basal bulb	18.7	12.8	Rectum/anal body diameter	1.5	1.2
Basal bulb diameter	13.5	10.6	Anus to phasmid	13.2	9.7
Basal bulb length/diameter	1.4	1.2	Phasmid position (% tail)	48.1	46.3
Corpus:isthmus ratio	3.1	2.7	Lateral field width	4.3	4.5
Nerve ring to anterior end	99.6	70.7	Lateral field width/body diameter (%)	22.2	23.9
Excretory pore to anterior end	99.7	79.5	Cuticle thickness	1.0	0.7
Deirid to anterior end	111.8	86.7	Annuli width	2.0	1.4
Nerve ring position (% pharynx)	63.7	70.9			

All measurements are in  $\mu\text{m}$ .

<sup>a</sup>Number of annuli from anus to the tail end.

L, body length; a, body length/body diameter; b, body length/Pharynx length; c, body length/tail length; c', tail length/diameter at anus region; V, % distance of vulva from anterior end/body length; G, % reproductive tract length/body length.

ted using Seinhorst's (1959) method and mounted in pure glycerin on HS slides (Shirayama *et al.*, 1993). Under an optical microscope (BX-51; Olympus, Tokyo, Japan) equipped with differential interference contrast (DIC), morphological and morphometric characters of the nematode specimens were observed and measured using a Cool Snap Photometrics color CCD digital camera (MP5.0-RTV-R-CLR-10; Photometrics, Tucson, AZ, USA) and the program QCapture Pro 5 (QImaging, Surrey, Canada).

## SYSTEMATIC ACCOUNTS

Order Rhabditida Chitwood, 1933  
Suborder Tylenchina Thorne, 1949

Infraorder Cephalobomorpha De Ley and Blaxter, 2002  
Family Cephalobidae Filipjev, 1934  
Genus *Acrobelloides* (Cobb, 1924) Thorne, 1937

### *Acrobelloides bodenheimeri* (Steiner, 1936)

Thorne, 1937 (Table 1, Fig. 1)

몽뚝꼬리뿔족입술선충 (신칭)

*Acrobeles bodenheimeri*: Steiner, 1936: 77, fig. 24.


*Cephalobus bodenheimeri*: Andr assy, 1984: 170.

*Rafiqius bodenheimeri*: Khan and Hussain, 1997: 140.

*Acrobelloides rotundifolius*: Bussau, 1991: 122, abb. 7.

*Acrobelloides bodenheimeri*: Thorn, 1937: 11.

**Material examined.** 1♀, Buk-myeon, Uichang-gu, Changwon-si, Gyeongsangnam-do, South Korea (GPS coordi-



**Fig. 1.** *Acrobeloides bodenheimeri* (Steiner, 1936) Thorne, 1937. A, Entire female; B, Female neck region; C, Female posterior region; D, Female head region; E, Female reproductive system. am, amphid; an, anus; bb, basal bulb; ca, cardia; co, corpus; cpa, cephalic papilla; de, deirid; ep, excretory pore; in, intestine; is, isthmus; lf, lateral field; lpa, labial papilla; lpr, labial probolae; nr, nerve ring; ovi, oviduct; ova, ovary; pa, primary axil; ph, phasmid; pus, post-uterine sac; re, rectum; spe, spermatheca; st, stoma; ut, uterus; va, vagina; vu, vulva.

nates: 35°22'22.5"N, 128°36'47.3"E), extracted by sieving and the Baermann funnel method from potato farm soil. The specimen (slide No. NIBRIV0000862892) is deposited at the National Institute of Biological Resources, South Korea.

**Measurements.** See Table 1.

**Description.** Female: Body cylindrical, length 529.5  $\mu\text{m}$  long, ventrally curved after fixation (Fig. 1A). Cuticle annulated; annuli 2.0  $\mu\text{m}$  wide and 1.0  $\mu\text{m}$  thick at mid-body. Lateral field occupying 22.2% of width of body at

mid-body. Lateral incisures varying in number along body length: three incisures between procorpus region and deirid, branching off from deirid into five incisures (Fig. 1B); two incisures fading out between anus and phasmid region; three incisures extending to tail terminus (Fig. 1C). Head region continuous with neck. Lip region 7.5  $\mu\text{m}$  in diameter, with triradiate symmetry with 6 + 4 papillae (Fig. 1D). Three pairs of asymmetrical lips; pairs of lips separated by U-shaped primary axils. Guarding process absent. Cephalic probolae absent. Three low and rounded

labial probolae present. Stoma cephaloboid; length 1.6 times the lip region diameter (Fig. 1B). Cheilorhabdions oval-shaped. Small dorsal denticle on metastom. Pharyngeal corpus fusiform with swollen metacarpus, 3.1 times isthmus length. Isthmus narrower than corpus, distinctly demarcated from metacarpus. Basal bulb oval-shaped with well-developed valves; 1.4 times as long as its width. Cardia inconspicuous, surrounded by intestinal tissue. Nerve ring located in posterior corpus, at 63.7% of pharynx length. Excretory pore position at posterior corpus, at 63.8% of pharynx length. Position of deirids in lateral field at anterior isthmus, at 71.5% of total neck length. Reproductive system monodelphic-prodelphic (Fig. 1E). Vulva lips not protruding. Vagina length 0.6 times body diameter. Post-uterine sac 2.6 times body width. Uterus length 3.2 times body diameter. Spermatheca 1.9 times body width. Oviduct short. Ovary straight, with a single row of oocytes. Rectum length 1.5 times anal body diameter (Fig. 1C). Tail conoid, with truncated terminus. Phasmids located middle of tail, at 48.1% of tail length.

Male: Unknown.

**Distribution.** Denmark (Bussau, 1991), Israel (Steiner, 1936), Mongolia (Andrássy, 1967), Malawi (Siddiqi *et al.*, 1992), South Korea (this study), Spain (Abolafia and Peña-Santiago, 2003), and USA (De Ley *et al.*, 1999).

**Habitat.** Soil sample from a potato farm.

**Remarks.** Morphological characters of the specimen described in this study generally agree with previous studies (Steiner, 1936; Siddiqi *et al.*, 1992; De Ley *et al.*, 1999; Abolafia and Peña-Santiago, 2003), except for total body length (529 vs 613–1,530  $\mu\text{m}$ ), the ratio of body length to body width ( $a = 27.2$  vs 15.0–24.3), the ratio of total body length to pharynx length ( $b = 3.4$  vs 4.0–7.8) and corpus to isthmus ratio (3.1 vs 3.3–7.2). Although these morphometric measurements of the specimen examined from a single individual differ from *A. bodenheimeri*, they are considered intraspecific variations as previously reported from many other *Acrobelloides* species (Anderson, 1965; 1968; Boström and Gydemo, 1983; De Ley *et al.*, 1999; Abolafia and Peña-Santiago, 2003). Morphometric analysis using multiple nematode individuals is further required to delimitate morphological variation of the species.

***Acrobelloides tricornis* (Thorne, 1925) Thorne, 1937 (Table 1, Fig. 2)**

삼각뿔족입술선충 (신칭)

*Acrobeles tricornis*: Thorne, 1925: 189, figs. 13.

*Acrobelloides setosus*: Brzeski, 1962: 337, figs. 7–9.

*Acrobelloides uberrinus*: Anderson, 1965: 232, fig. 1.

*Acrobelloides syrtisus*: Yeates, 1967: 530, fig. 2.

*Acrobelloides tricornis*: Thorne, 1937: 11.

**Material examined.** 1♀, Mechuri Island, Daenam-ro, Danwon-gu, Ansan-si, Gyeonggi-do, South Korea (GPS

coordinates: 37°12'02.9"N, 126°32'24.6"E), extracted by sieving and the Baermann funnel method from overgrown field soil. The specimen (slide No. ZCIVIV0000003207) is deposited at the National Institute of Biological Resources, South Korea.

**Measurements.** See Table 1.


**Description.** Female: Body cylindrical, length 332.0  $\mu\text{m}$  long, ventrally curved after fixation (Fig. 2A). Cuticle annulated; annuli 1.4  $\mu\text{m}$  wide and 0.7  $\mu\text{m}$  thick at mid-body. Lateral field occupying 23.9% of width of body at mid-body. Lateral incisures varying in number along body length: three incisures before deirid, branching off from deirid into five incisures (Fig. 2B); two incisures fading out at anus level; three incisures extending to tail terminus (Fig. 2C). Head region continuous with neck. Lip region 6.1  $\mu\text{m}$  in diameter, with triradiate symmetry with 6 + 4 papillae (Fig. 2D). Three pairs of asymmetrical lips; pairs of lips separated by U-shaped primary axils. Secondary axils shallow. Guarding process absent. Cephalic probolae absent. Three labial probolae high, with conical basal part and acute distal part. Stoma cephaloboid, length 1.8 times lip region diameter (Fig. 2B). Cheilorhabdions oval-shaped, strongly cuticularized. Pharyngeal corpus fusiform with swollen metacarpus, 2.7 times isthmus length. Isthmus narrower than corpus, distinctly demarcated from metacarpus. Basal bulb oval-shaped with well-developed valves; 1.2 times as long as its width. Cardia surrounded by intestinal tissue. Nerve ring located at anterior isthmus, at 70.9% of pharynx length. Excretory pore position at posterior isthmus, at 79.7% of pharynx length. Position of deirids in lateral field at basal bulb level, at 86.9% of total neck length. Reproductive system monodelphic-prodelphic (Fig. 2E). Vulva lips not protruding. Vagina length 0.3 times body diameter. Post-uterine sac inconspicuous. Uterus length 1.8 times body diameter. Spermatheca 0.9 times body width. Oviduct short. Rectum length 1.2 times anal body diameter (Fig. 2C). Tail conoid, with truncated terminus. Phasmids located middle of tail, at 46.3% of tail length.

Male: Unknown.

**Distribution.** Austria, Bulgaria, Canada, France, Georgia, Germany (Andrássy, 1984), Hungary (Andrássy, 2005), Kazakhstan, Kyrgyzstan, Lithuania, The Netherlands (Andrássy, 1984), New Zealand (Yeates, 1967; Boström and Holovachov, 2010), Norway (Spitzbergen) (Loof, 1971; Boström, 1987), Poland (Brzeski, 1962), Russia (Andrássy, 1984), Senegal (De Ley *et al.*, 1990; Boström and Holovachov, 2010), South Korea (this study), Spain (Abolafia and Peña-Santiago, 2003), USA (Thorne, 1925; Anderson, 1965), Tajikistan (Andrássy, 1984), Turkey (Boström, 1993), Turkmenistan, and Uzbekistan (Andrássy, 1984).

**Habitat.** Soil from an overgrown field.

**Remarks.** The specimen here described generally matches



**Fig. 2.** *Acrobeloides tricornis* (Thorne, 1925) Thorne, 1937. A, Entire female; B, Female neck region; C, Female posterior region; D, Female head region; E, Female reproductive system. am, amphid; an, anus; bb, basal bulb; ca, cardia; co, corpus; cpa, cephalic papilla; de, deirid; ep, excretory pore; in, intestine; is, isthmus; lf, lateral field; lpa, labial papilla; lpr, labial probolae; nr, nerve ring; ovi, oviduct; ova, ovary; pa, primary axil; ph, phasmid; re, rectum; sa, secondary axil; spe, spermatheca; st, stoma; ut, uterus; va, vagina; vu, vulva.

with morphological features that previously studied for *A. tricornis* (Thorne, 1925; Loof, 1971; Boström, 1993). *Acrobeloides setosus* Brezski, 1962, *A. syrtisus* Yeates, 1967 and *A. uberrinus* Anderson, 1965 were treated as synonyms of *A. tricornis* by Boström and Holovachov (2010). The examined specimen in this study is morphologically very similar to *A. uberrinus*, a junior synonym of *A. tricornis*, in the longitudinal incisures in the lateral field with three incisures extending to tail terminus (Anderson, 1965; De Ley *et al.*, 1990) (Fig. 2C). Further examination of multiple nematode individuals is needed to pro-

vide a species delimitation of their morphological variations.

The two newly reported *A. bodenheimeri* and *A. tricornis* species described herein are distinguished by shape of labial probolae (low and rounded vs high with acute distal part), post-uterine sac (distinct vs inconspicuous) and some morphometric characters (such as body length [529.5 vs 332.0] and the ratio of body length to body width [ $a=27.2$  vs 17.5]). In this study, we were not able to find male specimens of both species and species identification of nematodes was conducted based purely on morphologi-

cal characters examined from a single female individual, which is very common in soil nematode taxonomy. Moreover, morphological characters from male representatives in many *Acrobeloides* species (such as *A. apiculatus*, *A. nanus* and *A. varius*) have not yet been documented. Although a single female specimen was examined from each of the two species, their morphological characters described herein generally agree with those of *A. bodenheimeri* and *A. tricornis*.

### ACKNOWLEDGEMENTS

This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR201902204).

### REFERENCES

- Abolafia, J. and R. Peña-Santiago. 2003. Nematodes of the order Rhabditida from Andalucía Oriental, Spain. The genus *Acrobeloides* (Cobb, 1924) Thorne, 1937, with description of *A. arenicola* sp. n. and a key to species. *Journal of Nematode Morphology and Systematics* 5(2):107-130.
- Anderson, R.V. 1965. *Acrobeloides uberrinus* n. sp., with a note on morphologic variation within soil and bacteria-reared populations. *Proceedings of the Helminthological Society of Washington* 32(2):232-235.
- Anderson, R.V. 1968. Variation in taxonomic characters of a species of *Acrobeloides* (Cobb, 1924) Steiner and Bührer, 1933. *Canadian Journal of Zoology* 46(3):309-320.
- Andrássy, I. 1967. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 92. Weitere Bodennematoden aus den Jahren 1964 und 1965. *Opuscula Zoologica Budapest* 6(2):203-233.
- Andrássy, I. 1984. Klasse Nematoda (Ordnungen Monhysterida, Desmoscolecida, Araeolaimida, Chromadorida, Rhabditida). *Gustav Fischer Verlag, Stuttgart*.
- Andrássy, I. 2005. Free-living nematodes of Hungary: Nematoda errantia. Volume I. *Hungarian Natural History Museum, Budapest*.
- Baermann, G. 1917. Eine einfache Methode zur Auffindung von *Ankylostomum* (Nematoden) Larven in Erdproben. *Geneeskundig Tijdschrift voor Nederlandsch-Indië* 57:131-137.
- Boström, S. and R. Gydemo. 1983. Intraspecific variability in *Acrobeloides nanus* (de Man) Anderson (Nematoda, Cephalobidae) and a note on external morphology. *Zoologica Scripta* 12(4):245-255.
- Boström, S. 1987. A scanning electron microscope study of some species of terrestrial nematodes from Spitzbergen. *Nematologica* 33(4):366-374.
- Boström, S. 1993. Some cephalobids from Turkey (Nematoda: Rhabditida). *Nematologia Mediterranea* 21(2):295-300.
- Boström, S. and O. Holovachov. 2010. Description of *Acrobeloides uberrinus* Anderson, 1965 and *A. syrtisus* Yeates, 1967 (Rhabditida: Cephalobidae) with a discussion of the taxonomic status of *A. tricornis* (Thorne, 1925) Thorne, 1937 and *A. setosus* Brzeski, 1962. *Journal of Nematode Morphology and Systematics* 13(2):97-106.
- Brzeski, M. 1962. Three new species of the genus *Acrobeloides* Cobb. (Nematoda, Cephalobidae). *Bulletin de l'Académie Polonaise des Sciences. Classe II. Série des Sciences Biologiques* 10(8):335-339.
- Bussau, C. 1991. Freilebende Nematoden aus Küstendünen und angrenzenden Biotopen der deutschen und dänischen Küsten IV. Rhabditida und Tylenchida (Nematoda). *Zoologischer Anzeiger* 226(3-4):114-148.
- De Ley, P., E. Geraert and A. Coomans. 1990. Seven cephalobids from Senegal (Nematoda: Rhabditida). *Journal of African Zoology* 104(4):287-304.
- De Ley, P., M.-A. Félix, L.M. Frisse, S.A. Nadler, P.W. Sternberg and W.K. Thomas. 1999. Molecular and morphological characterisation of two reproductively isolated species with mirror-image anatomy (Nematoda: Cephalobidae). *Nematology* 1(6):591-612.
- Háněl, L. 1999. Fauna of soil nematodes (Nematoda) in Trojmezná hora Reserve. *Silva Gabreta* 3:89-94.
- Khan, H.A. and S.S. Hussain. 1997. Biosystematics of *Rafiqius saeedi* (Siddiqi, Deley and Khan, 1992) Gen. N. Comb. (Nematoda: Cephalobidae) with observation on its life cycle. *Pakistan Journal of Zoology* 29(2):139-143.
- Kim, T., J. Kim, Y.J. Bae and J.-K. Park. 2016. First Record of *Acrobeloides nanus* (Cephalobidae: Rhabditida: Nematoda) from Korea. *Animal Systematics, Evolution and Diversity* 32(4):258-265.
- Kim, T., J. Kim and J.-K. Park. 2017. *Acrobeloides varius* sp. n. (Rhabditida: Cephalobidae) from South Korea. *Nematology* 19(4):489-496.
- Loof, P.A.A. 1971. Freelifving and plant parasitic nematodes from Spitzbergen, collected by Mr. H. van Rossen. *Mededelingen Landbouwhogeschool Wageningen* 71(7):1-86.
- Pervez, R. 2011. *Acrobeloides ishraqi* sp. n. and *Acrobeloides mushtaqi* sp. n. (Nematoda: Rhabditida) from chickpea rhizosphere, Uttar Pradesh, India. *Archives of Phytopathology and Plant Protection* 44(15):1438-1446.
- Seinhorst, J.W. 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica* 4(1):67-69.
- Shirayama, Y., T. Kaku and R.P. Higgins. 1993. Double-sided microscopic observation of meiofauna using an HS-slide. *Benthos Research* 44:41-44.
- Siddiqi, M.R., P. De Ley and H.A. Khan. 1992. *Acrobeloides saeedi* sp. n. from Pakistan and redescription of *A. bodenheimeri* (Steiner) and *Placodira lobata* Thorne (Nematoda: Cephalobidae). *Afro-Asian Journal of Nematology* 2(1-2): 5-16.

Steiner, G. 1936. Opuscula miscellanea nematologica, IV. Proceedings of the Helminthological Society of Washington 3(2):74-80.

Thorne, G. 1925. The genus *Acrobeles* von Linstow, 1887. Transactions of the American Microscopical Society 44(4): 171-210.

Yeates, G.W. 1967. Studies on nematodes from dune sands. 5.

Acrobelinae. New Zealand Journal of Science 10(2):527-547.

*Submitted: October 6, 2021*

*Revised: October 19, 2021*

*Accepted: October 19, 2021*