

The first record of *Ulva adhaerens* (Ulvaceae, Chlorophyta) from Jeju Island, Korea

Hyung Woo Lee¹, Eun Hee Bae² and Myung Sook Kim^{1,3,*}

¹Research Institute for Basic Sciences, Jeju National University, Jeju 63243, Republic of Korea

²Microorganism Resources Division, National Institute of Biological Resources, Incheon 22689, Republic of Korea

³Department of Biology, Jeju National University, Jeju 63243, Republic of Korea

*Correspondent: myungskim@jejunu.ac.kr

The current surveys of *Ulva* in the subtidal area around Jeju Island give a chance to discover unrecorded green algal species of the Korean macroalgal flora. As a result of this investigation, we found *Ulva adhaerens* Matusmoto & Shimada, inhabiting the subtidal regions, up to 15 m deep, and conducted the DNA barcoding on plastid *rbcL*-3P and *tufA* regions with describing the morphological characteristics. Our specimens of *U. adhaerens* forms a monophyletic clade with the Japanese type specimen and *U. piritoka* Ngāti Kuri, Heesch & W.A. Nelson from New Zealand exhibiting each 0.3% sequence divergences, respectively, in the plastid *rbcL*-3P. The genetic variation of *U. adhaerens* clade is 1.0–3.9% in *rbcL*-3P and 4.8–9.8% in *tufA* to each *Ulva* species, including the generic type, *U. lactuca* Linnaeus. The morphology of Korean *U. adhaerens* specimens is identical to the type specimens of *U. adhaerens* from Japan having the development of rhizoidal filaments from both of the cell layers of the distromatic blade and the extension of rhizoidal clumps with adhesive trait between blades by extended rhizoidal clumps at the basal blades. The thallus attachment to substrate is by numerous minute discoidal plates made up of rhizoids originating from the inner part of distromatic blades in basal. Although there are still some problems to resolve the relationship between *U. adhaerens* and *U. piritoka* in the *rbcL* dataset and the phylogenetic pattern of the Group II intron of *rbcL*, we propose the new record of *U. adhaerens* in Korean macroalgal flora based on the morphological characteristics of Korean specimens. Continued study of the genus *Ulva* by morphological and molecular assessment will delimit the species of *Ulva*, elucidate the relationships between them, and uncover the species diversity.

Keywords: biodiversity, DNA barcoding, morphology, *rbcL*-3P, *tufA*, *Ulva adhaerens*

© 2022 National Institute of Biological Resources
DOI:10.12651/JSR.2022.11.4.266

INTRODUCTION

The genus *Ulva* Linnaeus includes 89 species as one of the most commonly distributed green macroalgae from intertidal and subtidal environments worldwide (Guiry and Guiry, 2022). Although the species identification of *Ulva* has traditionally relied on the morphological characteristics such as shape and thickness of blade, presence/absence of microscopic marginal denticulations, and the type of rhizoidal filaments (Hiraoka *et al.*, 2004), there are still experimental limitation to recognize each *Ulva* species correctly showing morphological plasticity (Spalding *et al.*, 2016). Since the molecular approach was applied to the genus *Ulva* to redefine the phylogenetic border between monostromatic and distromatic species

(Hayden *et al.*, 2003), the necessity of genetic information has been enlarged to reveal the species diversity (Heesch *et al.*, 2009), to investigate the systematics of *Ulva* (Kraft *et al.*, 2010), and to monitor the occurrence of invasive species and green tides in coastal areas (Krupnik *et al.*, 2018; Kang *et al.*, 2019).

The feasibility of DNA barcoding on *Ulva* was assessed using plastid markers, *rbcL*-3P, 3' end of the large subunit of the ribulose-bisphosphate carboxylase, and *tufA*, the elongation factor Tu (Saunders and Kucera, 2010). These markers have given the proof to delimit species boundaries in *Ulva* with understanding the local diversity (Heesch *et al.*, 2009; Kraft *et al.*, 2010; Kirkendale *et al.*, 2013; Miladi *et al.*, 2018) and revealing *Ulva* cryptic taxa (Kirkendale *et al.*, 2013; Lee *et al.*, 2019). In addition, the phy-

logenetic affinities within *Ulva* have been reassessed by the molecular analyses, giving us a more precise understanding of the evolutionary relationships. The advance of genetic assessment leads us to discover new species diversity of *Ulva* from extreme environment (Spalding *et al.*, 2016) and improve the ongoing taxonomic misapplication of species names by analyzing the lectotype specimens (Hughey *et al.*, 2019; 2021; 2022).

In Korea, 20 *Ulva* species have been recorded from the high intertidal shore through to the subtidal habitats (NIBR, 2019; MABIK, 2022). We discovered an unrecorded species during the surveys of subtidal zone and conducted a DNA barcoding assessment of the genus *Ulva* from Jeju Island of Korea. The aim of this study is to determine the taxonomic position of this undiscovered *Ulva* in Korea based on DNA barcoding of the plastid *rbcL*-3P and *tufA* regions with detailed morphological observation.

MATERIALS AND METHODS

Three specimens of *Ulva*, showing the blossom-like habit, were collected from the subtidal regions, up to 15 m depth on Munseom, an islet of Jeju Island, Korea (Table 1) using SCUBA. These fresh samples were photographed using an Olympus TG-4 camera (Olympus, Tokyo, Japan). Thallus fragments of each specimen were dried for molecular analyses using silica gel. In order to describe morphological characteristics, the samples were preserved in 5% formalin in seawater and sectioned using a bench-top freezing microtome (NK-101-II; Nippon Optical Works Co., Ltd., Tokyo, Japan). Sectioned materials were stained with 1% aniline blue acidified with 1% HCl after bleaching under sunlight. Sections were mounted in 35% custom-made corn syrup. Sectioned materials were photographed under a microscope (BX43; Olympus) using an EOS 600D digital camera (Canon, Tokyo, Japan). Digitized images were edited for clarity using Adobe Photoshop software (v. 6.1; Adobe Systems Inc., San Jose, CA, USA). Pressed herbarium specimens were deposited as voucher specimens in the herbarium of Jeju National University (JNUB) and the National Institute of Biological Resources (KB), Incheon, Korea.

The genomic DNA of *Ulva* specimens (Table 1) was extracted following the protocol of the LaboPass™ Tissue Genomic DNA Isolation Kit Mini (COSMO GENTECH, Seoul, Korea). AccuPower PCR Premix (Bioneer, Daejeon, Korea) was used according to the manufacturer's protocol for all PCR reactions. Two plastid DNA barcoding marker, *rbcL*-3P and *tufA*, were amplified and sequenced. The primer combinations of *rbcL*-3P and *tufA* in this study were *GrbcL*Fi (Saunders and Kucera, 2010)/1385R (Manhart, 1994), and *tufGF*4/*tufGR* (Saunders and Kucera, 2010), respectively. The PCR amplifi-

cation procedure followed that of Saunders and Kucera (2010). All purification of successfully amplified PCR products were conducted by AccuPrep PCR Purification Kit (Bioneer) and sequenced by Macrogen (Seoul, Korea) using forward and reverse primers. The *rbcL*-3P and *tufA* sequences for the phylogenetic analysis within *Ulva* were selected from GenBank (Table 1). The taxa selected for outgroups are as follows: *Umbraulva japonica* (Holmes) Bae & I.K. Lee, LC507134, *Ryuguphycus kuaweuweu* (H.L. Spalding & A.R. Sherwood) H. Kawai, T. Hanyuda & T. Kitayama, KT932987, and *Ulvaria obscura* (Kützinger) Gayral ex Bliding, HQ603651 for *rbcL*-3P; *Um. japonica*, LC507141, *R. kuaweuweu*, KT932969, and *Ul. obscura*, HQ610415 for *tufA*. The *rbcL*-3P and *tufA* datasets were aligned visually using BioEdit (Hall, 1999) after editing the *Ulva* sequences obtained in this study using Chromas ver. 1.45 software (Technelysium Pty Ltd., South Brisbane, Australia). Additionally, we obtained three sequences of 18S rRNA-5P from the same specimens without performing phylogenetic reconstruction (Table 1).

We assessed the levels of variation in the *rbcL*-3P and *tufA* sequences based on uncorrected pair-wise genetic distances (*p*-distance) using MEGA 5.1 (Tamura *et al.*, 2011), and the neighbor-joining (NJ) algorithm based on the Kimura-2-parameter (K2P) distance method. To determine the taxonomic position of *Ulva* specimens (Table 1), each *rbcL*-3P and *tufA* dataset was analyzed using maximum-likelihood (ML) in the RAxML software (Stamatakis, 2006). We ran the analysis with the GTR + Γ + I model of evolution and 1000 bootstrap replicates. Bayesian phylogenetic inference (BI) was generated by MrBayes ver. 3.1.2 (Ronquist and Huelsenbeck, 2003). Each phylogenetic tree was viewed FigTree v1.4.0 (Rambaut, 2012).

RESULTS

In the molecular comparison of the 767 bp dataset of *rbcL*-3P (Fig. 1) including 22 *Ulva* taxa and three outgroups (Table 1) sequences, 115 sites (15.0%) were variable and 77 sites (10.0%) were parsimoniously informative. The three blossom-like specimens from Jeju of Korea (MT978111, MT978113, MT978112) formed a clade with the Japanese type specimens of *Ulva adhaerens* Matsu-moto & Shimada (AB897327, AB894328) with 0.3% intra-specific variation and the type specimen from New Zealand of *U. piritoka* Ngāti Kuri, Heesch & W.A. Nelson (MW389665) with 0.3%. The topology of *rbcL*-3P ML analysis showed that the three sequences from Korea are the sister group to a sequence of *U. piritoka* from New Zealand (Fig. 1) with poor support (43% bootstrap and no posterior probability value). On the other hand, the *U. adhaerens* clade including *U. piritoka* in *rbcL*-3P phylo-

Table 1. The information of sequence dataset of *Ulva adhaerens* in this study and the representative ulvacean species from GenBank including outgroup taxa.

Species	Information	GenBank accession No.		Reference
		<i>rbcL</i>	<i>tufA</i>	
<i>Ulva adhaerens</i> Matsumoto & Shimada	MSK-GA00073; Korea	MT978111	MT978120	This study
	MSK-GA00074; Korea	MT978112	MT978121	This study
	MSK-GA00075; Korea	MT978113	MT978122	This study
	Japan	AB894327	–	Matsumoto and Shimada, 2015
	Japan	AB894328	–	Matsumoto and Shimada, 2015
<i>Ulva aragoënsis</i> (Bliding) Maggs	Israel	MG704815	MG976875	Krupnik <i>et al.</i> , 2018
<i>Ulva arasakii</i> Chihara	Japan	AB097621	–	Shimada <i>et al.</i> , 2003
	Japan	–	AB561079	Matsumoto <i>et al.</i> , 2011
<i>Ulva australis</i> Areschoug	Australia	MT815849		Hughey <i>et al.</i> , 2021
		MT624966	MT625139	Lee <i>et al.</i> , 2019
<i>Ulva brisbanensis</i> L.G. Kraft, Kraft & R.F. Waller	Australia	EU933945	–	Kraft <i>et al.</i> , 2010
<i>Ulva californica</i> Wille	Canada	HQ603514	HQ610279	Saunders and Kucera, 2010
<i>Ulva chaugulii</i> Kavale & Kazi	Israel	MG704805	MG976863	Krupnik <i>et al.</i> , 2018
<i>Ulva compressa</i> Linnaeus	Canada	HQ603521	HQ610285	Saunders and Kucera, 2010
<i>Ulva conglobata</i> Kjellman	Japan	AB894326		Matsumoto and Shimada, 2015
<i>Ulva expansa</i> (Setchell) Setchell & Gardner	USA	MH731009	MH731007	Hughey <i>et al.</i> , 2019
<i>Ulva fenestrata</i> Postels & Ruprecht	Russia	MK456393	MK456404	Hughey <i>et al.</i> , 2019
<i>Ulva gigantea</i> (Kützting) Bliding	Ireland	MT160606	MT160716	Fort <i>et al.</i> , 2021
<i>Ulva howensis</i> (A.H.S. Lucas) Kraft	Australia	JN082214	JN029310	Kirkendale <i>et al.</i> , 2013
<i>Ulva iliohaha</i> Spalding & Sherwood	USA	KT932995	KT932976	Spalding <i>et al.</i> , 2016
<i>Ulva intestinalis</i> Linnaeus	Canada	HQ603542	HQ610306	Saunders and Kucera, 2010
<i>Ulva lactuca</i> Linnaeus	Chile	MH730972	MH730972	Hughey <i>et al.</i> , 2019
<i>Ulva lacunculata</i> (Kützting) Wittrock	Croatia	MW543061		Hughey <i>et al.</i> , 2022
	Ireland	MT160587	MT160697	Fort <i>et al.</i> , 2021
<i>Ulva linza</i> Linnaeus	Canada	HQ603603	HQ610368	Saunders and Kucera, 2010
<i>Ulva ohiohilulu</i> Spalding & Sherwood	USA	KT932996	KT932978	Spalding <i>et al.</i> , 2016
<i>Ulva ohnoi</i> M. Hiraoka & S. Shimada	Japan	AB116037		Hiraoka <i>et al.</i> , 2004
	Korea	MT624835	MT625009	Lee <i>et al.</i> , 2019
<i>Ulva piritoka</i> Ngāti Kuri, Heesch & W.A. Nelson	New Zealand	MW389665		Heesch <i>et al.</i> , 2021
<i>Ulva procera</i> Hayden <i>et al.</i>	Canada	HQ603628	HQ610392	Saunders and Kucera, 2010
<i>Ulva prolifera</i> O.F. Müller	China	JQ867403	–	Shao <i>et al.</i> , 2015
<i>Ulva pseudo-ohnoi</i> H.W. Lee, J.C. Kang & M.S. Kim	Korea	MT624855	MT625027	Lee <i>et al.</i> , 2019
<i>Ulva rigida</i> C. Agardh	Ireland	MT160612	MT160722	Fort <i>et al.</i> , 2021
	Ireland	EU484401		Longhnae <i>et al.</i> , 2008
	Spain	MW543060		Hughey <i>et al.</i> , 2022
<i>Ulva sublittoralis</i> Segawa	Japan	AB741535		Ichihara <i>et al.</i> , 2013
<i>Ulva tepida</i> Masakiyo & Shimada	Israel	MG704820	MG976864	Krupnik <i>et al.</i> , 2018
<i>Umbraulva japonica</i>	Japan	LC507134	LC507141	Kawai <i>et al.</i> , 2020
<i>Ryugophycus kuaweuweu</i> (Spalding & Sherwood) H. Kawai, T. Hanyuda & T. Kitayama	Japan	KT932987	KT932969	Spalding <i>et al.</i> , 2016
<i>Ulvaria obscura</i> (Kützting) Gayral ex Bliding	Canada	HQ603651	HQ610415	Saunders and Kucera, 2010

geny was supported moderately by 86% BS and 1.0 PP value in ML and BI analysis, respectively. The *U. adhaerens* clade exhibited 2.9–3.1% interspecific divergence to the *U. lactuca* sequence of Lectotypic specimen (MH730972). The *U. adhaerens* clade formed a large clade with

the four foliose species, the type specimen of *U. rigida* C. Agardh from Spain (MW543060) with *U. rigida* from Ireland (EU484401, MT160612), *U. arasakii* M. Chihara from Japan (AB097621), *U. fenestrata* Postels & Ruprecht from Russia (MK456393), *U. expansa* (Setchell) Set-




Fig. 1. Phylogenetic tree of Ulvacean species inferred from the plastid *rbcL*-3P including *Ulva adhaerens* Matsumoto & Shimada collected from Jeju, Korea. Support values on each branch are ML bootstrap (>60%, left) and Bayesian posterior probability (>0.60, right). Sequences produced in this study are marked in bold font. Branch lengths are proportional to substitution rate.

chell & N.L. Gardner from USA (MH731009), and *U. australis* Areschoug from Korea (MT624966) exhibiting 1.0–1.4%, 1.6–2.1%, 2.3–2.6%, 2.5–2.9% and 2.2–2.4% interspecific divergence in *rbcL*-3P, respectively including the four tubular species, *U. compressa* Linnaeus (HQ603521) *U. intestinalis* Linnaeus (HQ603542), *U. howensis* (A.H.S. Lucas) Kraft (JN082214) and *U. prolifera* O.F. Müller (JQ867403) with 2.4–2.6%, 2.8–3.1%, 1.9–2.3% and 2.4–2.7%, respectively (Fig. 1).

The *tufA* sequence dataset (859 bp) composed of our

three specimens including 18 *Ulva* species and three outgroups (Table 1) had 225 variable sites (26.2%) and 115 parsimony-informative sites (13.4%). There are no *tufA* data for the type specimen of *U. adhaerens* from Japan and *U. piritoka* from New Zealand. The three specimens of *U. adhaerens* (MT978110, MT978111, MT978112) formed a distinct clade (Fig. 2). The topology of the *tufA* phylogeny (Fig. 2) was similar to that of the *rbcL* phylogeny (Fig. 1), inferring a large monophyly containing *U. rigida* from Ireland (MT160722), *U. australis* from Korea




Fig. 2. Phylogenetic tree of Ulvacean species inferred from the plastid *tufA* including *Ulva adhaerens* Matsumoto & Shimada collected from Jeju, Korea. Support values on each branch are ML bootstrap (>60%, left) and Bayesian posterior probability (>0.60, right). Sequences produced in this study are marked in bold font. Branch lengths are proportional to substitution rate.

(MT625139), *U. arasaki* from Japan (AB561079), *U. fenestrata* from Russia (MK456404), and *U. expansa* from USA (MH731007) with *U. adhaerens* from Korea. The interspecific divergence was 8.1–8.3% between *U. adhaerens* and *U. lactuca*, the type species of *Ulva* (MH730972). *Ulva adhaerens* was sister to the clade composed of *U. australis*, *U. arasaki*, *U. fenestrata* and *U. expansa* (Fig. 2), with 8.2%, 5.5–5.9%, 6.6–6.8% and 6.5–6.7% interspecific divergence, respectively. And it showed the biggest interspecific divergence as 9.0–9.3% to *U. pseudo-ohnoi* H.W. Lee, J.C. Kang & M.S. Kim (MT625027) (Fig. 2).

***Ulva adhaerens* Matsumoto & Shimada (Fig. 3A–N)**

Holotype. TNS-AL183435; TNS (Matsumoto and Shimada, 2015: 107, Fig. 60).

Type locality. Tenjin-jima, Sajima, Yokosuka, Kanagawa prefecture, Japan.

Korean name. 연접갈파래 (국명신칭): The Korean spe-

cific epithet, yeonjeop, is means “connected each other” in reference to the adherence between blades by the haptera of bundles of rhizoidal filaments arising from the cells of the ventral layer of the upper blade.

Habitat. Epilithic or epiphytic from the intertidal area (Matsumoto and Shimada, 2015) to the subtidal zone, up to 15 m deep (this study).

Distribution. Jeju Island, Korea (this study); Japan (Matsumoto and Shimada, 2015).

Specimens examined. MSK-GA00073 (Fig. 3B), MSK-GA00074, MSK-GA00075, Munseom, Jeju Island, Korea, Jan. 30, 2013 (deposited in JNUB); NIBR AL0000117920, Munseom, Jeju Island, Korea, Feb. 11, 2009, NIBR AL0000117921, Feb. 11, 2009, NIBR AL0000118004, Nov. 13, 2009, NIBR AL0000118012, May 13, 2009 (deposited in KB).

Other examined specimens. MSKL160414-17-1, MSKL-160414-17-2, MSKL-160414-17-3, MSKL-160414-17-4, and MSKL160414-17-5, Apr. 14, 2016 (deposi-




Fig. 3. *Ulva adhaerens* Matsumoto & Shimada. (A) Natural habit of *U. adhaerens* from Munseom, Jeju, growing blossom-like attached on benthic invertebrates, seaweeds or sands. (B) The voucher specimen (MSK-GA00073) of *U. adhaerens* from Munseom, Jeju, Korea, Jan 30, 2013. Thallus habit composed blossom-like and dorsiventral lobes having rounded and slightly ruffled blade. (C) Marginal part slightly uneven without marginal denticulations or microscopic protuberances. (D) Sub-roundish to polygonal cells at the surface view of upper part with irregular arrangement. (E) More angular cells at the surface view of basal part with irregular arrangement. Pyrenoids are one or two per cell (white arrows). (F) Transsection of blade at upper part composed of cuboidal cells with solid cell profiles. The height of both dorsal and ventral cells is similar. (G) Transsection at middle part composed of cuboidal cells with solid cell profiles. Dorsal cells are slightly higher than ventral ones. (H) Basal transsection composed of cuboidal cells with solid cell profiles. Dorsal cells are much higher than ventral ones. (I–K) Blade adherence by numerous and interwoven secondary rhizoids (black arrows). Blade adherence (black arrowheads) occurs at the basal part between dorsal plane of main blade and ventral plane of subsidiary one. Both secondary rhizoids originated from distromatic cells are fused to each other. (L–N) The development of minute rhizoidal holdfasts. Rhizoids are bud off from distromatic cells (black arrow) (L). Rhizoids are expanding through ventral cell plane downwardly (black arrows) (M). Fully expanded and interwoven rhizoids (black arrow) produce minute and disciform holdfast (black arrowhead) at the tips (N). Scale bars: B = 1 cm; C–H, L = 30 μ m; I, J, N = 150 μ m; K = 60 μ m; M = 50 μ m.

ted in JNUB).

DNA sequence data. *rbcL*, MSK-GA00073 (MT978111), MSK-GA00074 (MT978112), MSK-GA00075 (MT978113); *tufA*, MSK-GA00073 (MT978120), MSK-GA00074 (MT978121), MSK-GA00075 (MT978122); 18S rRNA, MSK-GA00073 (MT978102), MSK-GA00074 (MT978103), MSK-GA00075 (MT978104).

Habit and vegetative morphology. Thallus growing on various substrates such as rocks, shells, small or heavy benthic invertebrates, and cartilaginous macroalga or geniculate corallines in the subtidal (Fig. 3A). Thallus distromatic, foliose, and lobed (Fig. 3B); composed of several blades up to 5–7 one like a blossom; growing 5–6 cm in diameter up to 10 cm (Fig. 3B); attached by numerous minute discoidal plates without a distinct holdfast. Blade light-green to green in color (Fig. 3A). Blade margin slightly ruffled, entire, or slightly uneven without marginal denticulation (Fig. 3C). In the surface view, having polygonal to subspherical cell shape at the upper and middle part of blades, ranging 8–12 × 10–16 wide (Fig. 3D); more and more polygonal shape toward the basal region, ranging 10–22 × 20–40 wide (Fig. 3E). Each cell containing one or two pyrenoids (Fig. 3E), cells irregularly arranged throughout the thallus (Fig. 3D, E). In the transverse section, distromatic thallus composed of entirely cuboidal cells (Fig. 3F–H): 24–32 μm long × 6–15 μm wide, both dorsal and ventral at upper; 30–35 μm long × 6–15 μm wide, dorsal, and 25–30 μm × 6–15 μm wide, ventral, at middle; 40–50 μm long × 10–15 μm wide, dorsal, and 20–30 μm long × 15–25 μm wide, ventral at the base. Blade gradually thickened from upper to base, 48–64 μm at upper, 55–65 μm at middle, 60–80 μm at the base. At the basal blades, sequentially, numerous rhizoids budded off from the ventral-side cells (Fig. 3L); the interwoven rhizoidal bundle growing outside downwardly (Fig. 3M); a minute discoidal plate to attach on the substratum (Fig. 3N).

DISCUSSION

We discovered the unrecorded species, *U. adhaerens*, within the Korean macroalgal biodiversity, showing the interspecific variability between each *Ulva* species, 1.0–3.9% in *rbcL*-3P (Fig. 1) and 4.8–9.3% in *tufA* (Fig. 2). The plastid genes, *rbcL*-3P and *tufA*, which was proposed as DNA barcoding marker to be useful on the family Ulvaceae (Saunders and Kucera, 2010), have contributed to determine species diversity (Kirkendale *et al.*, 2013) and to reveal cryptic species locally (Miladi *et al.*, 2018; Steinhagen *et al.*, 2019). Also, they clarified the type specimens of *U. lactuca* Linnaeus and *U. fenestrata* Postels & Ruprecht (Hughey *et al.*, 2019). The DNA barcoding work, starting in subtidal discoveries from Jeju Island of Korea (Lee *et al.*, 2020), expands the understanding of the spe-

cies diversity for ulvacean taxa.

The reanalysis of ITS, *rbcL* and *tufA* markers from all GenBank dataset of distromatic *Ulva* showed the similar resolution and topology in the phylogenetic analyses and GMYC analyses (Fort *et al.*, 2022). Certainly, there is limitation in *rbcL* marker to delimit species boundaries between some clades, such as *U. ohnoi* and *U. lactuca* or an undescribed *Ulva* (as uploaded as *U. rigida*; this is not real) and *U. lacunculata* because of a small length of those alignment, but this limitation can be improved by using the *tufA* marker (Fort *et al.*, 2022). In addition, the *tufA* and ITS analysis each have limitation to distinguish between *U. aragoënsis* and *U. californica* Wille due to unclearly diverged topology (Krupnik *et al.*, 2018) between *U. ohnoi* and *U. pseudo-ohnoi* (as *U. reticulata* Forsskål) due to little sequence divergence (Fort *et al.*, 2022), respectively. However, *U. aragoënsis* is distinguished from *U. californica* by distinct morphological habit: *U. aragoënsis*, tubular and monostromatic (Cormaci *et al.*, 2014); *U. californica*, various foliose and distromatic (Tanner, 1986). *U. pseudo-ohnoi* (as *U. reticulata* specimens from Philippines) is not conspecific with *U. ohnoi* based on sexual isolation between these species in culture examination (Monotilla *et al.*, 2018).

Matsumoto and Shimada (2015) established *U. adhaerens* dependent on ITS and *rbcL* phylogeny and the diagnosis of rhizoid production and extension between distromatic cell layers reaching to adherence between thalli, and additionally described the presence of Group II intron of *rbcL* region. Recently, the closest species, *U. piritoka*, was newly described from New Zealand based on approximately 0.4% pairwise distance (just five substitutions of over 1149 base pairs), no Group II intron of *rbcL* region unlike *U. adhaerens*, and the geographic distance between Japan and New Zealand (Heesch *et al.*, 2021). The *rbcL* data from two Japanese and three Korean *U. adhaerens* specimens and a *U. piritoka* from New Zealand were separated into two small lineages: one is Korean specimens and *U. piritoka* with poor bootstrap and posterior probability support, the other Japanese type specimens with strong support (Fig. 1). However, although the adherence between thalli was not described in *U. piritoka*, both *U. adhaerens* and *U. piritoka* have distromatic and foliose thallus generally attached on rocks by rhizoidal clumps developing from both of the cell layers (Table 2).

Saunders and Kucera (2010) decided the *rbcL*-3P region as the more useful DNA barcode marker of green macroalga because of the presence of introns than the whole *rbcL* region of some species. The Group II intron of *rbcL* are described in some Ulvophycean taxa, such as *Bryopsis maxima* Okamura ex Segawa, *B. plumosa* (Hudson) C. Agardh, *Caulerpa okamurae* Weber Bosse, *Ca. racemosa* (Forsskål) J. Agardh, *Codium fragile* (Suringar) Hariot and *C. lucassii* Setchell in Bryopsidales (Hanyuda *et al.*, 2000).

Table 2. A comparison of morphological characteristics of *Ulva adhaerens* Matsumoto & Shimada and the closely related *Ulva* species.

Contents	<i>Ulva adhaerens</i> Matsumoto & Shimada (Korea)	<i>Ulva adhaerens</i> Matsumoto & Shimada (Japan)	<i>Ulva piritoka</i> Ngāti Kuri, Heesch & W.A. Nelson	<i>Ulva rigida</i> C. Agardh
Type locality	Japan: Sajima, Yokosuka, Kanagawa prefecture	Japan: Sajima, Yokosuka, Kanagawa prefecture	Tasman Bay, Manawatāwhi, New Zealand	Spain: Cádiz
Habitat	Attached on rocks, coralline algae and shells, intertidal to subtidal	Attached on bed-rocks or macroalgae, middle intertidal	Attached on rock and coralline algae, subtidal	Shallow subtidal up to ca. 8 m deep
Thallus	Distromatic, dorsiventrally lobed, with several blades, blossom-like	Distromatic, foliose, lobed	Distromatic, prostrate	Distromatic or deeply lobed with a short stipe, with a few perforations
Rhizoid production	Present	Present	Present	Undescribed
Adherence	Present at basal by bundles of numerous rhizoids	Present by bundles of rhizoids	Undescribed	Undescribed
Attachment	By scattered clumps made up of numerous rhizoids	Undescribed	By scattered clumps of rhizoids	Distinct, by a single discoid holdfast
Margin	Entire, slightly ruffled, without microscopic teeth	Entire, ruffled, no marginal denticulations	Entire, with ruffled or undulating	Smooth, with rarely simple microscopic teeth/bumps
Color	Light green to green	Light green to green	Light green	Dark green
Thallus size	5–6 cm in diameter	5–6 cm in diameter	1–2 cm in diameter	10–30 cm high up to 40 cm high
Thickness	48–80 µm from upper to basal, up to 120 µm	50–110 µm	45–55 µm at edge, becoming 90–165 µm	56–95 µm from upper to basal
Surface view	Polygonal to sub-roundish, irregularly arranged, 8–22 µm × 10–40 µm wide	Polygonal to sub-roundish, irregularly arranged	Undescribed	Angular to roundish polygonal, irregularly arranged
Pyrenoid	1–2 per cell	1–2 per cell	Undescribed	1–3 per cell, up to 5
Transverse view	Polygonal to rectilinear	Polygonal to rectilinear	Rectangular	Polygonal with angular or rounded corner at apical
Cell size	20–50 µm long × 6–25 µm wide	Undescribed	Undescribed	10–46 µm long × 6–37 µm wide
Reference	This study	Matsumoto and Shimada, 2015	Heesch <i>et al.</i> , 2021	Cornaci <i>et al.</i> , 2014, Hughey <i>et al.</i> , 2022

Table 2. Continued.

Contents	<i>Ulva expansa</i> (Setchell) Setchell & N.L. Gardner	<i>Ulva arasaki</i> M. Chihara	<i>Ulva fenestrata</i> Postel & Ruprecht	<i>Ulva australis</i> Areschoug
Type locality	USA: Monterey, California	Japan: Choshi, Chiba-ken	Russia: Kamchatka	Australia: Port Adelaide
Habitat	Shortly attached, generally free floating and drifting	Epilithic, middle to low intertidal	Attached on substrates, littoral to sublittoral	Attached or free floating
Thallus	Distromatic, orbicular or broadly elongated, without perforation	Distromatic, with perforations, generally simple	Distromatic, lamellar and expanded, whole or dissected, perforated	Distromatic, lobed, with irregular perforations
Rhizoid production	Undescribed	Undescribed	Undescribed	Undescribed
Adherence	Undescribed	Undescribed	Undescribed	Undescribed
Attachment	Undescribed	Distinct, by a single discoid holdfast	Distinct, by a single discoid holdfast	Distinct, by a single discoid holdfast
Margin	Deeply ruffled	Undulate or crisply ruffled, without microscopic teeth	Smooth or wavy, entire	Ruffled, sometimes with slightly microscopic teeth
Color	Pale green	Grass to dark green	Light to dark green, shiny	Deep to pale grass green
Thallus size	Up to 300 cm long, 18–75 cm wide	Mostly 20–50 cm, up to 80 cm	Up to 50 cm in diameter	10–15 cm long × 10–15 cm wide, up to 20 cm
Thickness	60–70 µm at central, 38–45 µm at margin	75–108 µm at central, 42–60 µm at margin	ca. 60 µm	ca. 90 µm up to 160 µm
Surface view	Undescribed	Subrectangular to polygonal with rounded corners, 12–20 µm wide	Undescribed	Roundish rectilinear to pentagonal, up to 20–36 µm wide
Pyrenoid	Undescribed	1–3 per cell, up to 4	Undescribed	1 per cell
Transverse view	Vertically elongated	Subquadrate, vertically elongated at center	Nearly square or slightly elongated	Rectilinear, with bluntly rounded
Cell size	28–30 µm long × 10–12 µm wide	15–45 µm long × 8–27 µm wide	ca. 20 µm long × 16 µm wide	46–65 µm long × 9–25 µm wide
Reference	Setchell and Gardner, 1920	Chihara, 1969, Bae, 2010	Setchell and Gardner, 1920 Zhigadlova and Lopatina, 2020	Kraft <i>et al.</i> , 2010, Lee <i>et al.</i> , 2019

In Ulvaceae, the Group II intron of *rbcL* is described only in *U. adhaerens* (Matsumoto and Shimada, 2015), although we didn't detect this intron as we applied the *rbcL*-3P marker of our Korean *U. adhaerens* specimens. *Ulva piritoka* was reported as a distinct species to *U. adhaerens* due to the hypothesis that the introns of *rbcL* region represents a relatively recent insertion into *U. adhaerens* lineage (Heesch *et al.*, 2021), but it has not been proved because of the limited number of species evaluated. Therefore, it is necessary to demonstrate the correlation between the Group II intron of *rbcL* and the identification of *U. adhaerens* and *U. piritoka*.

The morphological boundaries of each *Ulva* species have been described by thallus shape and thickness, cell size and shape, cell arrangement, chloroplast disposition, and number of pyrenoids (Koeman and van den Hoek, 1981). *Ulva adhaerens*, first reported from the Japanese coast as a new species, was characterized by a small distromatic algae growing singly or in clumps and having lobes adhered to each other by rhizoids (Matsumoto and Shimada, 2015). The Korean specimens exhibit the diagnosis of *U. adhaerens* from Japanese type specimens (Fig. 3I-K, Table 2) as supported by the culture result (Matsumoto and Shimada, 2015). Matsumoto and Shimada (2015) considered that *U. adhaerens* has a close phylogenetic affinity with *U. conglobata*, *U. lactuca* (as *U. fasciata*), *U. australis* (as *U. pertusa*) and *U. tannneri* based on the ball-forming morphology having undulate lobes. They mentioned that the presence or absence of marginal denticulation can be associated with the phylogenetic relationships among the above five species as a heritable characteristic (Hiraoka *et al.*, 2004). In the molecular analyses, *U. adhaerens* belongs to the large monophyletic clade in *Ulva* with several distromatic and foliose species, such as *U. australis*, *U. arasakii*, *U. expansa*, *U. fenestrata* and *U. rigida* (Figs. 1 and 2). The Korean and Japanese *U. adhaerens*, including *U. piritoka*, have the entire margins without marginal denticulation and microscopic teeth or bumps, although *U. rigida* have the simple microscopic teeth or bumps (Table 2). These morphological characteristics support the clade of *U. adhaerens* clade including *U. piritoka* with moderate bootstrap and posterior probability support which is distinguished from the sister taxon of *U. rigida* (Fig. 1).

The habitat and geographical distribution of *U. adhaerens* was first limited in intertidal from the central coastal region facing the Pacific (Matsumoto and Shimada, 2015). Our discovery expands the vertical habitat range of *U. adhaerens* to the subtidal, up to 15 m deep. Efforts to discover new entities of *Ulva* in the subtidal region contribute to widening the ecological knowledge, such as habitat range and environment (Lee *et al.*, 2020). And the geographic distribution can be expanded potentially towards the southern hemisphere through the further molecular

examination using *tufA* marker of the type specimens of *U. adhaerens* and *U. piritoka* with the more detailed morphological observations for the thallus adherence. Since the phylogenetic relationships are still uncertain enough to delimit species boundaries of all *Ulva* members, the continuing investigations of morphological traits and molecular affinities will lead the deep understanding of *Ulva* on species diversity, environmental adaptation and evolutionary history.

ACKNOWLEDGEMENTS

We thank Dr. Kang, J. C., Dr. Yang, M. Y. and members of the Molecular Phylogeny of Marine Algae Laboratory at Jeju National University for helping us to collect samples. This study was supported by the 2022 education, research and student guidance grant funded by Jeju National University.

REFERENCES

- Bae, E.H. 2010. Ulothrichales, Ulvales. In: Algal flora of Korea. Volume 1, Number 1. Chlorophyta: Ulvophyceae: Ulothrichales, Ulvales, Cladophorales, Bryopsidales. Marine green algae. (Bae, E. H., Kim, H.-S., Kwon, C.-J., Hwang, I.-K., Kim, G. H. & Klochkova, T. A. Eds), pp. 7-52. Incheon: National Institute of Biological Resources.
- Chihara, M. 1969. *Ulva arasakii*, a new species of green algae: its life history and taxonomy. Bulletin of the National Science Museum, Tokyo, Series B (Botany) 12:849-862.
- Cormaci, M., G. Furnari and G. Alongi. 2014. Flora marina bentonica del Mediterraneo: Chlorophyta. Bollettino dell'Accademia Gioenia di Scienze Naturali di Catani 47:11-436.
- Fort, A., M. McHale, K. Cascella, P. Potin, M. Rerrineau, P.D. Kerrison, E. da Coasta, R. Calado, M. do Rosário Domingues, I.C. Azevedo, I. Sousa-Pinto, C. Gachon, A. van der Werf, W. de Visser, J.E. Beniers, H. Jansen, M.D. Guiry and R. Sulpice. 2022. Exhaustive reanalysis of barcode sequences from public repositories highlights ongoing misidentifications and impacts taxa diversity and distribution. Mol. Ecol. Resour. 22:86-101.
- Fort, A., M. McHale, K. Cascella, P. Potin, B. Usadel, M.D. Guiry and R. Sulpice. 2021. Foliose *Ulva* species show considerable inter-specific genetic diversity, low intra-specific variation, and rare occurrence of inter-specific hybrid in the wild. J. Phycol. 57:219-233.
- Guiry, M.D. and G.M. Guiry. 2022. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway [Available from: <http://www.algaebase.org>. Accessed on January 30, 2022].
- Hall, T.A. 1999. BioEdit: a user-friendly biological alignment editor and analysis program for Windows 95/98/NT. Nucl-

- ic Acids Symp. Ser. 41:95-98.
- Hanyuda, T., S. Arai and K. Ueda. 2000. Variability in the *rbcL* Introns of Caulerpallean Algae (Chlorophyta, Ulvophyceae). *J. Plant Res.* 113:403-413.
- Hayden, H.S., J. Blomster, C.A. Maggs, P.C. Silva, M.J. Stanhop and R. Waaland. 2003. Linnaeus was right all along: *Ulva* and *Enteromorpha* are distinct genera. *Eur. J. Phycol.* 38:277-294.
- Heesch, S., J.E.S. Broom, K.F. Neill, T.J. Farr, J.L. Dalen and W.A. Nelson. 2009. *Ulva*, *Umbraulva* and *Gemina*: genetic survey of New Zealand taxa reveals diversity and introduced species. *Eur. J. Phycol.* 44:143-154.
- Heesch, S., J.E. Sutherland, S. Ringham and W.A. Nelson. 2021. *Ulva* L. (Ulvales, Chlorophyta) from Manawatāwhi/Three Kings Islands, New Zealand: *Ulva piritoka* Ngāti Kuri, Heesch & W.A. Nelson sp. nov. and records of two non-native species, *U. compressa* and *U. rigida*. *Cryptogam., Algol.* 42:129-140.
- Hiraoka, M., S. Shimada, M. Uenosono and M. Masuda. 2004. A new green-tide-forming alga, *Ulva ohnoi* Hiraoka et Shimada sp. nov. (Ulvales, Ulvophyceae) from Japan. *Phycol. Res.* 52:17-29.
- Hughey, J.R., C.A. Maggs, F. Mineur, C. Jarvis, K.A. Miller, S.H. Shabaka and P.W. Gabrielson. 2019. Genetic analysis of the Linnaean *Ulva lactuca* (Ulvales, Chlorophyta) holotype and related type specimens reveals name misapplications, unexpected origins, and new synonymies. *J. Phycol.* 55:503-508.
- Hughey, J.R., P.W. Gabrielson, C.A. Maggs and F. Mineur. 2022. Genomic analysis of the lectotype specimens of European *Ulva rigida* and *Ulva lacunculata* (Ulvaceae, Chlorophyta) reveals the ongoing misapplication of names. *Eur. J. Phycol.* 52:143-153.
- Hughey, J.R., P.W. Gabrielson, C.A. Maggs, F. Mineur and K.A. Miller. 2021. Taxonomic revisions based on genetic analysis of type specimens of *Ulva conglobata*, *U. laetevierns*, *U. pertusa* and *U. spathulata* (Ulvales, Chlorophyta). *Phycol. Res.* 69:148-153.
- Ichihara, K., K. Miyaji and S. Shimada. 2013. Comparing the low-salinity tolerance of *Ulva* species distributed in different environments. *Phycol. Res.* 61:52-57.
- Kang, J.H., J.E. Jang, J.H. Kim, S.Y. Byeon, S. Kim, S.K. Choi, Y.H. Kang, S.R. Park and H.J. Lee. 2019. Species composition, diversity, and distribution of the genus *Ulva* along the coast of Jeju Island, Korea based on molecular phylogenetic analysis. *PLoS ONE* 14(7):e0219958. <https://doi.org/10.1371/journal.pone.0219958>.
- Kawai, H., T. Hanyuda, I. Mine, S. Takaichi, R. Terada and T. Kitayama. 2020. Morphology and molecular phylogeny of *Umbraulva* spp. (Ulvales, Ulvophyceae), and proposal of *Ryuguphycus* gen. nov. and *R. kuaweuweu* comb. nov. *Eur. J. Phycol.* 56(1). <https://doi.org/10.1080/09670262.2020.1753815>.
- Kirkendale, L., G.W. Saunders and P. Winberg. 2013. A molecular survey of *Ulva* (Chlorophyta) in temperate Australia reveals enhanced levels of cosmopolitanism. *J. Phycol.* 49:69-81.
- Koeman, R.P.T. and C. van den Hoek. 1981. The taxonomy of *Ulva* (Chlorophyceae) in the Netherlands. *Br. Phycol. J.* 16:9-53.
- Kraft, L.G.K., G.T. Kraft and R.F. Waller. 2010. Investigations into southern Australian *Ulva* (Ulvophyceae, Chlorophyta) taxonomy and molecular phylogeny indicate both cosmopolitanism and endemic cryptic species. *J. Phycol.* 46:1257-1277.
- Krupnik, N., G. Paz, J. Douek, E. Lewinsohn, N. Carmel, F. Mineur and C.A. Maggs. 2018. Native, invasive and cryptogenic *Ulva* species from Israeli Mediterranean Sea: risk and potential. *Mediterranean Marine Science* 19:132-146.
- Lee, H.W., E.H. Bae and M.S. Kim. 2020. *Umbraulva yun-seulla* sp. nov. (Ulvaceae, Chlorophyta) from a subtidal habitat of Jeju Island. *Algae* 35:349-359.
- Lee, H.W., J.C. Kang and M.S. Kim. 2019. Taxonomy of *Ulva* causing from Jeju Island, Korea with new species, *U. pseudo-ohnoi* sp. nov. (Ulvales, Chlorophyta). *Algae* 34:253-266.
- Longhane, C.J., L.M. McIvor, F. Rindi, D.B. Stengel and M.D. Guiry. 2008. Morphology, *rbcL* phylogeny and distribution of distromatic *Ulva* (Ulvophyceae, Chlorophyta) in Ireland and southern Britain. *Phycologia* 47:416-429.
- Manhart, J.R. 1994. Phylogenetic analysis of green plant *rbcL* sequences. *Mol. Phylogenet. Evol.* 3:114-127.
- Matsumoto, K. and S. Shimada. 2015. Systematics of green algae resembling *Ulva conglobata*, with a description of *Ulva adhaerens* sp. nov. (Ulvales, Ulvophyceae). *Eur. J. Phycol.* 50:100-111.
- Matsumoto, T., F. Shinozaki, T. Chikuni, A. Yabuki, K. Takishita, M. Kawachi, T. Nakayama, I. Inouye, T. Hashimoto and Y. Inagaki. 2011. Green-colored plastids in the dinoflagellate genus *Lepidodinium* are of Core Chlorophyte Origin. *Protist* 162:268-276.
- Miladi, R., A. Manghisi, S.A. Minicante, G. Genovese, S. Abdelkafi and M. Morabito. 2018. A DNA barcoding survey of *Ulva* (Chlorophyta) in Tunisia and Italy reveals the presence of the overlooked alien *U. ohnoi*. *Cryptogam. Algol.* 39:85-107.
- Monotilla, A.P., T. Nishimura, M. Adachi, Y. Tani, D.B. Largo and M. Hiraoka. 2018. Examination of prezygotic and postzygotic isolating barriers in tropical *Ulva* (Ulvophyceae, Chlorophyta): evidence for ongoing speciation. *J. Phycol.* 54:539-549.
- National Institute of Biological Resources. 2019. National Species List of Korea. I. Plants, Fungi, Algae, Prokaryotes. Designzip. 924 pp. [in Korea].
- National Marine Biodiversity Institute of Korea. 2022. 2022 National List of Marine Species. III. Marine Plants. Namuprint. 99 pp. [in Korea].
- Rambaut, A. 2012. FigTree, tree figure drawing tool v1.4.0. Institute of Evolutionary Biology, University of Edinburgh [Available from: <http://tree.bio.ed.ac.uk/software/figtree/>,

- Accessed on May 30, 2018].
- Ronquist, F. and J.P. Huelsenbeck. 2003. Mr Bayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572-1574.
- Saunders, G.W. and H. Kucera. 2010. An evaluation of *rbcL*, *tufA*, UPA, LSU and ITS as DNA barcode markers for the marine green macroalgae. *Cryptogam. Algol.* 31:487-528.
- Setchell, W.A. and N.L. Gardner. 1920. The marine algae of the Pacific coast of North America. Part II. Chlorophyceae. University of California Publications in Botany 8: 139-374, pls. 9-33.
- Shao, Z., W. Li, H. Guo and D. Duan. 2015. Isolation, expression and characterization of *rbcL* gene from *Ulva prolifera* J. Agardh (Ulvophyceae, Chlorophyta). *J. Ocean Univ. China* 14:1087-1095.
- Shimada, S., M. Hiraoka, S. Nabata, M. Lima and M. Masuda. 2003. Molecular phylogenetic analyses of the Japanese *Ulva* and *Enteromorpha* (Ulvales, Ulvophyceae), with special reference to the free-floating *Ulva*. *Phycol. Res.* 51:99-108.
- Spalding, H.L., K.Y. Conklin, C.M. Smith, C.J. O'Kelly and A.R. Sherwood. 2016. New Ulvaceae (Ulvophyceae, Chlorophyta) from mesophotic ecosystems across the Hawaiian Archipelago. *J. Phycol.* 52:40-53.
- Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688-2690.
- Steinhagen, S., R. Karez and F. Weinberger. 2019. Cryptic, alien and lost species: molecular diversity of *Ulva sensu lato* along the German coasts of the North and Baltic Seas. *Eur. J. Phycol.* 54:466-483.
- Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei and S. Kumar. 2011. MEGA5: Molecular Evolutionary Genetic Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. & Evol.* 28:2731-2739.
- Tanner, C.E. 1986. Investigations of the taxonomy and morphological variation of *Ulva* (Chlorophyta): *Ulva californica* Wille. *Phycologia* 25:510-520.
- Zhigadlova, G.G. and N.A. Lopatina. 2020. [Green algae Chlorophyta algae]. In: Flora and Fauna of Matua Island (middle Kuril Islands): the field guide. Volume 1. Sea. (Sanamyan, K. E. & Sanamyan, N. P. Eds), pp. 468-485. Cherepovets: Intron [in Russian].

Submitted: May 28, 2021

Revised: June 23, 2021

Accepted: July 22, 2022