

FUZZY STABILITY OF AN ADDITIVE-QUADRATIC FUNCTIONAL EQUATION WITH THE FIXED POINT ALTERNATIVE

JEONG PIL SEO^a, SUNGJIN LEE^{b,*} AND REZA SAADATI^c

ABSTRACT. In [41], Th.M. Rassias proved that the norm defined over a real vector space V is induced by an inner product if and only if for a fixed positive integer l

$$2l \left\| \frac{1}{2l} \sum_{i=1}^{2l} x_i \right\|^2 + \sum_{i=1}^{2l} \left\| x_i - \frac{1}{2l} \sum_{j=1}^{2l} x_j \right\|^2 = \sum_{i=1}^{2l} \|x_i\|^2$$

holds for all $x_1, \dots, x_{2l} \in V$. For the above equality, we can define the following functional equation

$$(0.1) \quad 2lf \left(\frac{1}{2l} \sum_{i=1}^{2l} x_i \right) + \sum_{i=1}^{2l} f \left(x_i - \frac{1}{2l} \sum_{j=1}^{2l} x_j \right) = \sum_{i=1}^{2l} f(x_i).$$

Using the fixed point method, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces.

1. INTRODUCTION AND PRELIMINARIES

The stability problem of functional equations originated from a question of Ulam [51] concerning the stability of group homomorphisms. Hyers [13] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [1] for additive mappings and by Th.M. Rassias [40] for linear mappings by considering an unbounded Cauchy difference. A generalization of the Th.M. Rassias theorem was obtained by Găvruta [12] by replacing the unbounded Cauchy difference by a general control function in the spirit of the Th.M. Rassias' approach.

The functional equation

Received by the editors July 06, 2015. Accepted July 21, 2015.

2010 *Mathematics Subject Classification.* 46S40, 46C05, 39B52, 26E50, 47H10.

Key words and phrases. fuzzy Banach space, fixed point, functional equation related to inner product space, Hyers-Ulam stability, quadratic mapping, additive mapping.

*Corresponding author.

$$f(x + y) + f(x - y) = 2f(x) + 2f(y)$$

is called a *quadratic functional equation*. In particular, every solution of the quadratic functional equation is said to be a *quadratic mapping*. A Hyers-Ulam stability problem for the quadratic functional equation was proved by Skof [50] for mappings $f : X \rightarrow Y$, where X is a normed space and Y is a Banach space. Cholewa [8] noticed that the theorem of Skof is still true if the relevant domain X is replaced by an Abelian group. Czerwik [9] proved the Hyers-Ulam stability of the quadratic functional equation. The stability problems of several functional equations have been extensively investigated by a number of authors and there are many interesting results concerning this problem (see [14, 16, 17, 31, 32, 33, 34, 35, 37, 38, 39, 42, 43, 44, 45, 46, 47, 48, 49]).

In [33], Park, Lee and Shin proved that an even mapping $f : V \rightarrow W$ satisfies the functional equation (0.1) if and only if the even mapping $f : V \rightarrow W$ is quadratic. Moreover, they proved the Hyers-Ulam stability of the quadratic functional equation (0.1) in real Banach spaces.

Katsaras [18] defined a fuzzy norm on a vector space to construct a fuzzy vector topological structure on the space. Some mathematicians have defined fuzzy norms on a vector space from various points of view [11, 20, 52]. In particular, Bag and Samanta [2], following Cheng and Mordeson [7], gave an idea of fuzzy norm in such a manner that the corresponding fuzzy metric is of Kramosil and Michalek type [19]. They established a decomposition theorem of a fuzzy norm into a family of crisp norms and investigated some properties of fuzzy normed spaces [3].

We use the definition of fuzzy normed spaces given in [2, 24, 25] to investigate a fuzzy version of the Hyers-Ulam stability for the functional equation (0.1) in the fuzzy normed vector space setting.

Definition 1.1 ([2, 24, 25, 26]). Let X be a real vector space. A function $N : X \times \mathbb{R} \rightarrow [0, 1]$ is called a *fuzzy norm* on X if for all $x, y \in X$ and all $s, t \in \mathbb{R}$,

- (N₁) $N(x, t) = 0$ for $t \leq 0$;
- (N₂) $x = 0$ if and only if $N(x, t) = 1$ for all $t > 0$;
- (N₃) $N(cx, t) = N(x, \frac{t}{|c|})$ if $c \neq 0$;
- (N₄) $N(x + y, s + t) \geq \min\{N(x, s), N(y, t)\}$;
- (N₅) $N(x, \cdot)$ is a non-decreasing function of \mathbb{R} and $\lim_{t \rightarrow \infty} N(x, t) = 1$;
- (N₆) for $x \neq 0$, $N(x, \cdot)$ is continuous on \mathbb{R} .

The pair (X, N) is called a *fuzzy normed vector space*.

The properties of fuzzy normed vector spaces and examples of fuzzy norms are given in [24, 27].

Definition 1.2 ([2, 24, 25, 26]). Let (X, N) be a fuzzy normed vector space. A sequence $\{x_n\}$ in X is said to be *convergent* or *converge* if there exists an $x \in X$ such that $\lim_{n \rightarrow \infty} N(x_n - x, t) = 1$ for all $t > 0$. In this case, x is called the *limit* of the sequence $\{x_n\}$ and we denote it by $N\text{-}\lim_{n \rightarrow \infty} x_n = x$.

Definition 1.3 ([2, 24, 25]). Let (X, N) be a fuzzy normed vector space. A sequence $\{x_n\}$ in X is called *Cauchy* if for each $\varepsilon > 0$ and each $t > 0$ there exists an $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ and all $p > 0$, we have $N(x_{n+p} - x_n, t) > 1 - \varepsilon$.

It is well-known that every convergent sequence in a fuzzy normed vector space is Cauchy. If each Cauchy sequence is convergent, then the fuzzy norm is said to be *complete* and the fuzzy normed vector space is called a *fuzzy Banach space*.

We say that a mapping $f : X \rightarrow Y$ between fuzzy normed vector spaces X and Y is continuous at a point $x_0 \in X$ if for each sequence $\{x_n\}$ converging to x_0 in X , then the sequence $\{f(x_n)\}$ converges to $f(x_0)$. If $f : X \rightarrow Y$ is continuous at each $x \in X$, then $f : X \rightarrow Y$ is said to be *continuous* on X (see [3]).

Let X be a set. A function $d : X \times X \rightarrow [0, \infty]$ is called a *generalized metric* on X if d satisfies

- (1) $d(x, y) = 0$ if and only if $x = y$;
- (2) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (3) $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in X$.

We recall a fundamental result in fixed point theory.

Theorem 1.4 ([4, 10]). Let (X, d) be a complete generalized metric space and let $J : X \rightarrow X$ be a strictly contractive mapping with Lipschitz constant $L < 1$. Then for each given element $x \in X$, either

$$d(J^n x, J^{n+1} x) = \infty$$

for all nonnegative integers n or there exists a positive integer n_0 such that

- (1) $d(J^n x, J^{n+1} x) < \infty$, $\forall n \geq n_0$;
- (2) the sequence $\{J^n x\}$ converges to a fixed point y^* of J ;
- (3) y^* is the unique fixed point of J in the set $Y = \{y \in X \mid d(J^{n_0} x, y) < \infty\}$;
- (4) $d(y, y^*) \leq \frac{1}{1-L} d(y, Jy)$ for all $y \in Y$.

In 1996, G. Isac and Th.M. Rassias [15] were the first to provide applications of stability theory of functional equations for the proof of new fixed point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [5, 6, 27, 29, 30, 36]).

Starting with the paper [24], the stability of some functional equations in the framework of fuzzy normed spaces or random normed spaces has been investigated (see e.g., [21, 22, 23, 24, 25, 26, 27, 28]).

This paper is organized as follows: In Section 2, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for an odd case. In Section 3, we prove the Hyers-Ulam stability of the functional equation (0.1) in fuzzy Banach spaces for an even case.

Throughout this paper, assume that X is a vector space and that (Y, N) is a fuzzy Banach space. Let l be a fixed positive integer.

2. HYERS-ULAM STABILITY OF THE FUNCTIONAL EQUATION (0.1): AN ODD CASE

For a given mapping $f : X \rightarrow Y$, we define

$$Cf(x_1, \dots, x_{2l}) := 2lf\left(\frac{1}{2l}\sum_{i=1}^{2l}x_i\right) + \sum_{i=1}^{2l}f\left(x_i - \frac{1}{2l}\sum_{j=1}^{2l}x_j\right) - \sum_{i=1}^{2l}f(x_i)$$

for all $x_1, \dots, x_{2l} \in X$.

Using the fixed point method, we prove the Hyers-Ulam stability of the functional equation $Cf(x_1, \dots, x_{2l}) = 0$ in fuzzy Banach spaces: an odd case.

Theorem 2.1. *Let $\varphi : X^{2l} \rightarrow [0, \infty)$ and $\psi(x) := \varphi(\underbrace{x, \dots, x}_{l \text{ times}}, \underbrace{0, \dots, 0}_{l \text{ times}})$ be functions such that there exists an $L < 1$ with*

$$\varphi(x_1, \dots, x_{2l}) \leq \frac{L}{2}\varphi(2x_1, \dots, 2x_{2l})$$

for all $x_1, \dots, x_{2l} \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying

$$(2.1) \quad N(Cf(x_1, \dots, x_{2l}), t) \geq \frac{t}{t + \varphi(x_1, \dots, x_{2l})}$$

for all $x_1, \dots, x_{2l} \in X$ and all $t > 0$. Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that

$$(2.2) \quad N(f(x) - A(x), t) \geq \frac{(l - lL)t}{(l - lL)t + \psi(x)}$$

for all $x \in X$ and all $t > 0$.

Proof. Letting $x_1 = \cdots = x_l = x$ and $x_{l+1} = \cdots = x_{2l} = 0$ in (2.1), we get

$$(2.3) \quad N\left(2lf\left(\frac{x}{2}\right) - lf(x), t\right) \geq \frac{t}{t + \varphi(\underbrace{x, \dots, x}_{l \text{ times}}, \underbrace{0, \dots, 0}_{l \text{ times}})} = \frac{t}{t + \psi(x)}$$

for all $x \in X$.

Consider the set

$$S := \{g : X \rightarrow Y\}$$

and introduce the generalized metric on S :

$$d(g, h) = \inf\{\mu \in \mathbb{R}_+ : N(g(x) - h(x), \mu t) \geq \frac{t}{t + \psi(x)}, \forall x \in X, \forall t > 0\},$$

where, as usual, $\inf \phi = +\infty$. It is easy to show that (S, d) is complete. (See the proof of Lemma 2.1 of [22].)

Now we consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := 2g\left(\frac{x}{2}\right)$$

for all $x \in X$.

Let $g, h \in S$ be given such that $d(g, h) = \varepsilon$. Then

$$N(g(x) - h(x), \varepsilon t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. Hence

$$\begin{aligned} N(Jg(x) - Jh(x), L\varepsilon t) &= N\left(2g\left(\frac{x}{2}\right) - 2h\left(\frac{x}{2}\right), L\varepsilon t\right) \\ &= N\left(g\left(\frac{x}{2}\right) - h\left(\frac{x}{2}\right), \frac{L}{2}\varepsilon t\right) \\ &\geq \frac{\frac{Lt}{2}}{\frac{Lt}{2} + \psi\left(\frac{x}{2}\right)} \geq \frac{\frac{Lt}{2}}{\frac{Lt}{2} + \frac{L}{2}\psi(x)} \\ &= \frac{t}{t + \psi(x)} \end{aligned}$$

for all $x \in X$ and all $t > 0$. So $d(g, h) = \varepsilon$ implies that $d(Jg, Jh) \leq L\varepsilon$. This means that

$$d(Jg, Jh) \leq Ld(g, h)$$

for all $g, h \in S$.

It follows from (2.3) that $d(f, Jf) \leq \frac{1}{l}$.

By Theorem 1.4, there exists a mapping $A : X \rightarrow Y$ satisfying the following:

(1) A is a fixed point of J , i.e.,

$$(2.4) \quad A\left(\frac{x}{2}\right) = \frac{1}{2}A(x)$$

for all $x \in X$. Since $f : X \rightarrow Y$ is odd, $A : X \rightarrow Y$ is an odd mapping. The mapping A is a unique fixed point of J in the set

$$M = \{g \in S : d(f, g) < \infty\}.$$

This implies that A is a unique mapping satisfying (2.4) such that there exists a $\mu \in (0, \infty)$ satisfying

$$N(f(x) - A(x), \mu t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$;

(2) $d(J^n f, A) \rightarrow 0$ as $n \rightarrow \infty$. This implies the equality

$$N\text{-}\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right) = A(x)$$

for all $x \in X$;

(3) $d(f, A) \leq \frac{1}{1-L}d(f, Jf)$, which implies the inequality

$$d(f, A) \leq \frac{1}{l - lL}.$$

This implies that the inequality (2.2) holds.

By (2.1),

$$N\left(2^n Cf\left(\frac{x_1}{2^n}, \dots, \frac{x_{2l}}{2^n}\right), 2^n t\right) \geq \frac{t}{t + \varphi\left(\frac{x_1}{2^n}, \dots, \frac{x_{2l}}{2^n}\right)}$$

for all $x_1, \dots, x_{2l} \in X$, all $t > 0$ and all $n \in \mathbb{N}$. So

$$N\left(2^n Cf\left(\frac{x_1}{2^n}, \dots, \frac{x_{2l}}{2^n}\right), t\right) \geq \frac{\frac{t}{2^n}}{\frac{t}{2^n} + \frac{L^n}{2^n}\varphi(x_1, \dots, x_{2l})}$$

for all $x_1, \dots, x_{2l} \in X$, all $t > 0$ and all $n \in \mathbb{N}$. Since $\lim_{n \rightarrow \infty} \frac{\frac{t}{2^n}}{\frac{t}{2^n} + \frac{L^n}{2^n}\varphi(x_1, \dots, x_{2l})} = 1$ for all $x_1, \dots, x_{2l} \in X$ and all $t > 0$,

$$N(CA(x_1, \dots, x_{2l}), t) = 1$$

for all $x_1, \dots, x_{2l} \in X$ and all $t > 0$. Thus $CA(x_1, \dots, x_{2l}) = 0$. Since A is odd, it follows from Lemma 2.1 of [35] that the mapping $A : X \rightarrow Y$ is additive, as desired. \square

Corollary 2.2. Let $\theta \geq 0$ and let p be a real number with $p > 1$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an odd mapping satisfying

$$(2.5) \quad N(Cf(x_1, \dots, x_{2l}), t) \geq \frac{t}{t + \theta \sum_{j=1}^{2l} \|x_j\|^p}$$

for all $x_1, \dots, x_{2l} \in X$ and all $t > 0$. Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(2^p - 2)t}{(2^p - 2)t + 2^p \theta \|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 2.1 by taking

$$\varphi(x_1, \dots, x_{2l}) := \theta \sum_{j=1}^{2l} \|x_j\|^p$$

for all $x_1, \dots, x_{2l} \in X$. Then we can choose $L = 2^{1-p}$ and we get the desired result. \square

Theorem 2.3. Let $\varphi : X^{2l} \rightarrow [0, \infty)$ and $\psi(x) := \varphi(\underbrace{x, \dots, x}_{l \text{ times}}, \underbrace{0, \dots, 0}_{l \text{ times}})$ be functions such that there exists an $L < 1$ with

$$\varphi(x_1, \dots, x_{2l}) \leq 2L\varphi\left(\frac{x_1}{2}, \dots, \frac{x_{2l}}{2}\right)$$

for all $x_1, \dots, x_{2l} \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.1). Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(l - lL)t}{(l - lL)t + L\psi(x)}$$

for all $x \in X$ and all $t > 0$.

Proof. Let (S, d) be the generalized metric space defined in the proof of Theorem 2.1.

Consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := \frac{1}{2}g(2x)$$

for all $x \in X$.

It follows from (2.3) that

$$N(2lf(x) - lf(2x), t) \geq \frac{t}{t + \psi(2x)}$$

for all $x \in X$ and all $t > 0$. Thus

$$N\left(f(x) - \frac{1}{2}f(2x), \frac{2L}{2l}t\right) \geq \frac{2Lt}{2Lt + 2L\psi(x)} = \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. So $d(f, Jf) \leq \frac{L}{l}$.

The rest of the proof is similar to the proof of Theorem 2.1. \square

Corollary 2.4. *Let $\theta \geq 0$ and let p be a real number with $0 < p < 1$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.5). Then $A(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x)$ exists for each $x \in X$ and defines an additive mapping $A : X \rightarrow Y$ such that*

$$N(f(x) - A(x), t) \geq \frac{(2 - 2^p)t}{(2 - 2^p)t + 2^p\theta\|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 2.3 by taking

$$\varphi(x_1, \dots, x_{2l}) := \theta \sum_{j=1}^{2l} \|x_j\|^p$$

for all $x_1, \dots, x_{2l} \in X$. Then we can choose $L = 2^{p-1}$ and we get the desired result. \square

3. HYERS-ULAM STABILITY OF THE FUNCTIONAL EQUATION (0.1): AN EVEN CASE

In this section, using the fixed point method, we prove the Hyers-Ulam stability of the functional equation $Cf(x_1, \dots, x_{2l}) = 0$ in fuzzy Banach spaces: an even case.

Theorem 3.1. *Let $\varphi : X^{2l} \rightarrow [0, \infty)$ and $\psi(x) := \varphi(\underbrace{x, \dots, x}_{l \text{ times}}, \underbrace{0, \dots, 0}_{l \text{ times}})$ be functions such that there exists an $L < 1$ with*

$$\varphi(x_1, \dots, x_{2l}) \leq \frac{L}{4} \varphi(2x_1, \dots, 2x_{2l})$$

for all $x_1, \dots, x_{2l} \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.1). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that

$$(3.1) \quad N(f(x) - Q(x), t) \geq \frac{(l - lL)t}{(l - lL)t + \psi(x)}$$

for all $x \in X$ and all $t > 0$.

Proof. Letting $x_1 = \cdots = x_l = x$ and $x_{l+1} = \cdots = x_{2l} = 0$ in (2.1), we get

$$(3.2) \quad N\left(4lf\left(\frac{x}{2}\right) - lf(x), t\right) \geq \frac{t}{t + \underbrace{\varphi(x, \dots, x)}_{l \text{ times}}, \underbrace{0, \dots, 0}_{l \text{ times}}} = \frac{t}{t + \psi(x)}$$

for all $x \in X$.

Let (S, d) be the generalized metric space defined in the proof of Theorem 2.1.

Now we consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := 4g\left(\frac{x}{2}\right)$$

for all $x \in X$.

Let $g, h \in S$ be given such that $d(g, h) = \varepsilon$. Then

$$N(g(x) - h(x), \varepsilon t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$ and all $t > 0$. Hence

$$\begin{aligned} N(Jg(x) - Jh(x), L\varepsilon t) &= N\left(4g\left(\frac{x}{2}\right) - 4h\left(\frac{x}{2}\right), L\varepsilon t\right) \\ &= N\left(g\left(\frac{x}{2}\right) - h\left(\frac{x}{2}\right), \frac{L}{4}\varepsilon t\right) \geq \frac{\frac{Lt}{4}}{\frac{Lt}{4} + \psi\left(\frac{x}{2}\right)} \\ &\geq \frac{\frac{Lt}{4}}{\frac{Lt}{4} + \frac{L}{4}\psi(x)} = \frac{t}{t + \psi(x)} \end{aligned}$$

for all $x \in X$ and all $t > 0$. So $d(g, h) = \varepsilon$ implies that $d(Jg, Jh) \leq L\varepsilon$. This means that

$$d(Jg, Jh) \leq Ld(g, h)$$

for all $g, h \in S$.

It follows from (3.2) that $d(f, Jf) \leq \frac{1}{l}$.

By Theorem 1.4, there exists a mapping $Q : X \rightarrow Y$ satisfying the following:

(1) Q is a fixed point of J , i.e.,

$$(3.3) \quad Q\left(\frac{x}{2}\right) = \frac{1}{4}Q(x)$$

for all $x \in X$. Since $f : X \rightarrow Y$ is even, $Q : X \rightarrow Y$ is an even mapping. The mapping Q is a unique fixed point of J in the set

$$M = \{g \in S : d(f, g) < \infty\}.$$

This implies that Q is a unique mapping satisfying (3.3) such that there exists a $\mu \in (0, \infty)$ satisfying

$$N(f(x) - Q(x), \mu t) \geq \frac{t}{t + \psi(x)}$$

for all $x \in X$;

(2) $d(J^n f, Q) \rightarrow 0$ as $n \rightarrow \infty$. This implies the equality

$$N\text{-}\lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right) = Q(x)$$

for all $x \in X$;

(3) $d(f, Q) \leq \frac{1}{1-L}d(f, Jf)$, which implies the inequality

$$d(f, Q) \leq \frac{1}{l - lL}.$$

This implies that the inequality (3.1) holds.

The rest of the proof is similar to the proof of Theorem 2.1. □

Corollary 3.2. *Let $\theta \geq 0$ and let p be a real number with $p > 2$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.5). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that*

$$N(f(x) - Q(x), t) \geq \frac{(2^p - 4)t}{(2^p - 4)t + 2^p\theta\|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 3.1 by taking

$$\varphi(x_1, \dots, x_{2l}) := \theta \sum_{j=1}^{2l} \|x_j\|^p$$

for all $x_1, \dots, x_{2l} \in X$. Then we can choose $L = 2^{2-p}$ and we get the desired result. □

Similarly, we can obtain the following. We will omit the proof.

Theorem 3.3. *Let $\varphi : X^{2l} \rightarrow [0, \infty)$ and $\psi(x) := \varphi(\underbrace{x, \dots, x}_{l \text{ times}}, \underbrace{0, \dots, 0}_{l \text{ times}})$ be functions such that there exists an $L < 1$ with*

$$\varphi(x_1, \dots, x_{2l}) \leq 4L\varphi\left(\frac{x_1}{2}, \dots, \frac{x_{2l}}{2}\right)$$

for all $x_1, \dots, x_{2l} \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.1). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{4^n} f(2^n x)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that

$$N(f(x) - A(x), t) \geq \frac{(l - lL)t}{(l - lL)t + \psi(x)}$$

for all $x \in X$ and all $t > 0$.

Corollary 3.4. *Let $\theta \geq 0$ and let p be a real number with $0 < p < 2$. Let X be a normed vector space with norm $\|\cdot\|$. Let $f : X \rightarrow Y$ be an even mapping satisfying $f(0) = 0$ and (2.5). Then $Q(x) := N\text{-}\lim_{n \rightarrow \infty} \frac{1}{4^n} f(2^n x)$ exists for each $x \in X$ and defines a quadratic mapping $Q : X \rightarrow Y$ such that*

$$N(f(x) - Q(x), t) \geq \frac{(4 - 2^p)t}{(4 - 2^p)t + 2^p \theta \|x\|^p}$$

for all $x \in X$ and all $t > 0$.

Proof. The proof follows from Theorem 3.3 by taking

$$\varphi(x_1, \dots, x_{2l}) := \theta \sum_{j=1}^{2l} \|x_j\|^p$$

for all $x_1, \dots, x_{2l} \in X$. Then we can choose $L = 2^{p-2}$ and we get the desired result. \square

REFERENCES

1. T. Aoki: On the stability of the linear transformation in Banach spaces. *J. Math. Soc. Japan* **2** (1950), 64-66.
2. T. Bag & S.K. Samanta: Finite dimensional fuzzy normed linear spaces. *J. Fuzzy Math.* **11** (2003), 687-705.
3. ———: Fuzzy bounded linear operators. *Fuzzy Sets and Systems* **151** (2005), 513-547.
4. L. Cădariu & V. Radu: Fixed points and the stability of Jensen's functional equation. *J. Inequal. Pure Appl. Math.* **4**, no. 1, Art. ID 4 (2003).
5. ———: On the stability of the Cauchy functional equation: a fixed point approach. *Grazer Math. Ber.* **346** (2004), 43-52.
6. ———: Fixed point methods for the generalized stability of functional equations in a single variable. *Fixed Point Theory and Applications* **2008**, Art. ID 749392 (2008).
7. S.C. Cheng & J.M. Mordeson: Fuzzy linear operators and fuzzy normed linear spaces. *Bull. Calcutta Math. Soc.* **86** (1994), 429-436.
8. P.W. Cholewa: Remarks on the stability of functional equations. *Aequationes Math.* **27** (1984), 76-86.
9. S. Czerwik: On the stability of the quadratic mapping in normed spaces. *Abh. Math. Sem. Univ. Hamburg* **62** (1992), 59-64.
10. J. Diaz & B. Margolis: A fixed point theorem of the alternative for contractions on a generalized complete metric space. *Bull. Amer. Math. Soc.* **74** (1968), 305-309.

11. C. Felbin: Finite dimensional fuzzy normed linear spaces. *Fuzzy Sets and Systems* **48** (1992), 239-248.
12. P. Găvruta: A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings. *J. Math. Anal. Appl.* **184** (1994), 431-436.
13. D.H. Hyers: On the stability of the linear functional equation. *Proc. Nat. Acad. Sci. U.S.A.* **27** (1941), 222-224.
14. D.H. Hyers, G. Isac & Th.M. Rassias: *Stability of Functional Equations in Several Variables*. Birkhäuser, Basel, 1998.
15. G. Isac & Th.M. Rassias: Stability of ψ -additive mappings: Applications to nonlinear analysis. *Internat. J. Math. Math. Sci.* **19** (1996), 219-228.
16. S. Jung: *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*. Hadronic Press Inc., Palm Harbor, Florida, 2001.
17. ———: Approximation of analytic functions by Hermite functions. *Bull. Sci. Math.* **133** (2009), 756-764.
18. A.K. Katsaras: Fuzzy topological vector spaces II. *Fuzzy Sets and Systems* **12** (1984), 143-154.
19. I. Kramosil & J. Michalek: Fuzzy metric and statistical metric spaces. *Kybernetika* **11** (1975), 326-334.
20. S.V. Krishna & K.K.M. Sarma: Separation of fuzzy normed linear spaces. *Fuzzy Sets and Systems* **63** (1994), 207-217.
21. D. Mihet: The fixed point method for fuzzy stability of the Jensen functional equation. *Fuzzy Sets and Systems* **160** (2009), 1663-1667.
22. D. Mihet & V. Radu: On the stability of the additive Cauchy functional equation in random normed spaces. *J. Math. Anal. Appl.* **343** (2008), 567-572.
23. A.K. Mirmostafae: A fixed point approach to almost quartic mappings in quasi fuzzy normed spaces. *Fuzzy Sets and Systems* **160** (2009), 1653-1662.
24. A.K. Mirmostafae, M. Mirzavaziri & M.S. Moslehian: Fuzzy stability of the Jensen functional equation. *Fuzzy Sets and Systems* **159** (2008), 730-738.
25. A.K. Mirmostafae & M.S. Moslehian: Fuzzy versions of Hyers-Ulam-Rassias theorem. *Fuzzy Sets and Systems* **159** (2008), 720-729.
26. ———: Fuzzy approximately cubic mappings. *Inform. Sci.* **178** (2008), 3791-3798.
27. M. Mirzavaziri & M.S. Moslehian: A fixed point approach to stability of a quadratic equation. *Bull. Braz. Math. Soc.* **37** (2006), 361-376.
28. A.K. Mirmostafae & M.S. Moslehian: Fuzzy stability of additive mappings in non-Archimedean fuzzy normed spaces. *Fuzzy Sets and Systems* **160** (2009), 1643-1652.
29. C. Park: Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras. *Fixed Point Theory and Applications* **2007**, Art. ID 50175 (2007).

30. ———: Generalized Hyers-Ulam-Rassias stability of quadratic functional equations: a fixed point approach, *Fixed Point Theory and Applications* **2008**, Art. ID 493751 (2008).
31. C. Park, Y. Cho & M. Han: Functional inequalities associated with Jordan-von Neumann type additive functional equations. *J. Inequal. Appl.* **2007**, Art. ID 41820 (2007).
32. C. Park & J. Cui: Generalized stability of C^* -ternary quadratic mappings. *Abstract Appl. Anal.* **2007**, Art. ID 23282 (2007).
33. C. Park, J. Lee & D. Shin: Quadratic mappings associated with inner product spaces. (preprint).
34. C. Park & A. Najati: Homomorphisms and derivations in C^* -algebras. *Abstract Appl. Anal.* **2007**, Art. ID 80630 (2007).
35. C. Park, W. Park & A. Najati: Functional equations related to inner product spaces. *Abstract Appl. Anal.* **2009**, Art. ID 907121 (2009).
36. V. Radu: The fixed point alternative and the stability of functional equations. *Fixed Point Theory* **4** (2003), 91-96.
37. J.M. Rassias: On approximation of approximately linear mappings by linear mappings. *Bull. Sci. Math.* **108** (1984), 445-446.
38. ———: Refined Hyers-Ulam approximation of approximately Jensen type mappings. *Bull. Sci. Math.* **131** (2007), 89-98.
39. J.M. Rassias & M.J. Rassias: Asymptotic behavior of alternative Jensen and Jensen type functional equations. *Bull. Sci. Math.* **129** (2005), 545-558.
40. Th.M. Rassias: On the stability of the linear mapping in Banach spaces. *Proc. Amer. Math. Soc.* **72** (1978), 297-300.
41. ———: New characterizations of inner product spaces. *Bull. Sci. Math.* **108** (1984), 95-99.
42. ———: Problem 16; 2. Report of the 27th International Symp. on Functional Equations. *Aequationes Math.* **39** (1990), 292-293; 309.
43. ———: On the stability of the quadratic functional equation and its applications. *Studia Univ. Babeş-Bolyai* **XLIII** (1998), 89-124.
44. ———: The problem of S.M. Ulam for approximately multiplicative mappings. *J. Math. Anal. Appl.* **246** (2000), 352-378.
45. ———: On the stability of functional equations in Banach spaces. *J. Math. Anal. Appl.* **251** (2000), 264-284.
46. ———: On the stability of functional equations and a problem of Ulam. *Acta Appl. Math.* **62** (2000), 23-130.
47. Th.M. Rassias & P. Šemrl: On the behaviour of mappings which do not satisfy Hyers-Ulam stability. *Proc. Amer. Math. Soc.* **114** (1992), 989-993.
48. ———: On the Hyers-Ulam stability of linear mappings. *J. Math. Anal. Appl.* **173** (1993), 325-338.

49. Th.M. Rassias & K. Shibata: Variational problem of some quadratic functionals in complex analysis. *J. Math. Anal. Appl.* **228** (1998), 234-253.
50. F. Skof: Proprietà locali e approssimazione di operatori. *Rend. Sem. Mat. Fis. Milano* **53** (1983), 113-129.
51. S.M. Ulam: *A Collection of the Mathematical Problems*. Interscience Publ. New York, 1960.
52. J.Z. Xiao & X.H. Zhu: Fuzzy normed spaces of operators and its completeness. *Fuzzy Sets and Systems* **133** (2003), 389-399.

^aOHSANG HIGH SCHOOL, GUMI 730-842, KYEONGSANGBUK DO, KOREA
Email address: `sjp4829@hanmail.net`

^bDEPARTMENT OF MATHEMATICS, DAEJIN UNIVERSITY, KYEONGGI 487-711, KOREA
Email address: `hyper@daejin.ac.kr`

^cDEPARTMENT OF MATHEMATICS, IRAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, TEHRAN, IRAN
Email address: `rsaadati@eml.cc`